



Emendation of *Propionibacterium acnes* subsp. *acnes* (Deiko et al. 2015) and proposal of *Propionibacterium acnes* type II as *Propionibacterium acnes* subsp. *defendens* subsp. nov

McDowell, A., Barnard, E., Liu, J., Li, H., & Patrick, S. (2016). Emendation of *Propionibacterium acnes* subsp. *acnes* (Deiko et al. 2015) and proposal of *Propionibacterium acnes* type II as *Propionibacterium acnes* subsp. *defendens* subsp. nov. *International Journal of Systematic and Evolutionary Microbiology*, 66(12), 5358-5365. Advance online publication. <https://doi.org/10.1099/ijsem.0.001521>

[Link to publication record in Ulster University Research Portal](#)

Published in:
International Journal of Systematic and Evolutionary Microbiology

Publication Status:
Published online: 01/12/2016

DOI:
[10.1099/ijsem.0.001521](https://doi.org/10.1099/ijsem.0.001521)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Title:

Emendation of *Propionibacterium acnes* subsp. *acnes* (Deiko et al. 2015) and proposal of *Propionibacterium acnes* type II as *Propionibacterium acnes* subsp. *defendens* subsp. nov

Running title:

Taxonomic reclassification of *P. acnes* types I and II

Authors:

Andrew McDowell¹, Emma Barnard², Jared Liu², Huiying Li^{2,3}, Sheila Patrick⁴

Affiliations:

¹Northern Ireland Centre for Stratified Medicine, Biomedical Sciences Research Institute, C-TRIC Building, Altnagelvin Area Hospital, University of Ulster, Londonderry, UK.

²Department of Molecular and Medical Pharmacology, Crump Institute for Molecular Imaging, David Geffen School of Medicine, UCLA, California, USA.

³UCLA-DOE Institute for Genomics and Proteomics, Los Angeles, California, USA.

⁴Centre for Infection & Immunity, School of Medicine, Dentistry & Biomedical Sciences, Queen's University, Belfast, UK.

Correspondence:

Dr Andrew McDowell; Email: a.mcdowell@ulster.ac.uk; Telephone +44 (0) 2871 675662 (extn 75662)

Keywords: *Propionibacterium acnes* subsp. *defendens* subsp. nov., type I, type II, Multilocus Sequence Analysis, Genomics, Phenotype

Subject category: New taxa - Actinobacteria

Abstract

Recently, strains of *Propionibacterium acnes* from the type III genetic division have been proposed as *Propionibacterium acnes* subsp. *elongatum* subsp. nov., with strains from the type I and II divisions collectively classified as *Propionibacterium acnes* subsp. *acnes* subsp. nov. Under such a taxonomic re-appraisal, we believe that types I and II should also have their own separate rank of subspecies. In support of this, we describe a polyphasic taxonomic study based on the analysis of publically available multilocus and whole genome sequence datasets, alongside a systematic review of previously published phylogenetic, genomic, phenotypic and clinical data. Strains of types I and II form highly distinct clades based on multilocus sequence analysis (MLSA) and whole genome phylogenetic reconstructions. *In silico* or digital DNA-DNA similarity values also fall within the 70-80% boundary recommended for bacterial subspecies. Furthermore, we see important differences in genome content, including the presence of an active CRISPR/Cas system in type II strains, but not type I, and evidence for increasing linkage equilibrium within the separate divisions. Key biochemical differences include positive tests for β -haemolytic, neuraminidase and sorbitol fermentation activities with type I strains, but not type II. We now emend the description of *Propionibacterium acnes* subsp. *acnes* (Deiko et al. 2015) and propose type II as *Propionibacterium acnes* subsp. *defendens* subsp. nov. The type strain of *Propionibacterium acnes* subsp. *defendens* subsp. nov. is ATCC 11828 (=JCM 6473=CCUG 6369).

Propionibacterium acnes is a Gram-positive anaerobic bacterium and a member of the 'cutaneous' group of human propionibacteria along with *Propionibacterium granulosum*, *Propionibacterium avidum* and *Propionibacterium humerusii*. Although found predominately on the skin, it can also be isolated from the oral cavity and the genitourinary and gastrointestinal tracts (Patrick and McDowell, 2011). While the bacterium is most noted for its association with the inflammatory skin condition acne vulgaris (Lomholt and Kilian, 2010, McDowell et al., 2012, Fitz-Gibbon et al., 2013), there is now a growing recognition that the spectrum of opportunistic infections and clinical conditions to which it may be associated has been underestimated (Tunney et al., 1999, Cohen et al., 2005, Cavalcanti et al., 2011, Eishi, 2013; Barnard et al., 2016).

In the last 10 years, significant advances in our understanding of this bacterium at the population genetic level have been made using single, multilocus and whole genome sequence analyses (McDowell et al., 2005, McDowell et al., 2008, Lomholt and Kilian, 2010, McDowell et al., 2012, Fitz-Gibbon et al., 2013, Tomida et al., 2013, Scholz et al., 2014). Such work has demonstrated the phylogenetically distinct nature of the originally described *P. acnes* serotypes, designated types I and II, and identified a new type, designated type III, which displays an ability to form long filamentous cell structures not seen with types I and II (McDowell et al., 2005, McDowell et al., 2008). These studies have also identified further phylogenetic subdivisions within the type I clade (IA₁, IA₂, IB, IC) which differ in genome content, inflammatory potential, association with disease, production of putative virulence determinants, resistance to antibiotics used in the treatment of acne, as well as biochemical and aggregative properties (Valanne et al., 2005, McDowell et al., 2013, Tomida et al., 2013, Johnson et al., 2016, Scholz et al., 2016).

Very recently, Dekio *et al.* (2015) proposed that *P. acnes* type III be reclassified as *Propionibacterium acnes* subsp. *elongatum* subsp. nov. based on phylogenetic, genomic and phenotypic differences, with strains of type I and II classified as *Propionibacterium acnes* subsp. *acnes* subsp. nov. (Dekio *et al.*, 2015). In bacterial taxonomy, there are currently no clear guidelines for the establishment of subspecies, and the proposal of such essentially remains at the discretion of the researcher. Nevertheless, the proposal of a new bacterial subspecies is normally based on consistent phylogenetic differences and phenotypic variations between groups of strains within a species (Brenner *et al.*, 2000). If the major phylogroups of *P. acnes* are now to be reclassified within a subspecies framework, then strains of types I and II also deserve separate taxonomic ranks of subspecies. In this paper, we describe a polyphasic taxonomic study based on the analysis of publically available multilocus sequence and whole genome datasets, alongside a review of published phylogenetic, genomic, phenotypic and clinical data, that support the division and reclassification of *P. acnes* type I and type II as distinct subspecies. We now emend the description of *Propionibacterium acnes* subsp. *acnes* to include type I only since it contains the type strain, and propose type II as *Propionibacterium acnes* subsp. *defendens* subsp. nov. Type III strains remain as *Propionibacterium acnes* subsp. *elongatum* (hereafter described as type III) as previously proposed (Dekio *et al.*, 2015).

In 2005, we demonstrated that the *P. acnes* serotypes known as types I and II represented highly distinct phylogenetic groups based on sequence analysis of the *recA* housekeeping gene, as well as the putative haemolysin/ FtsJ-like methyltransferase gene *tly* (McDowell *et al.*, 2005). Application of *recA* typing was also central in the identification of strains representing the type III phylogenetic division (McDowell *et al.*, 2008). Since then, two key

MLSA methods based on eight (MLSA₈) and nine protein-encoding genes (MLSA₉) have been described for this bacterium, both based on completely different sets of genetic loci (Lomholt and Kilian, 2010, McDowell et al., 2012). With both independent MLSA schemes we find that types I, II and III form highly distinct clades consistent with the original *recA* and *tly* analysis, and supported by high bootstrap values (Fig. 1). This phylogenetic clustering is also highly congruent with that obtained upon whole genome analysis of 124,731 SNPs in shared or ‘core’ regions of 85 *P. acnes* genomes spanning all the major phylogenetic divisions (Fig. S1); the average p-distance between each of the types based on core region analysis is 0.444 for types I and II, 0.487 for types I and III, and 0.470 for types II and III (Table 1). Twenty six percent of core region SNPs are unique to type I, with 22% unique to type II and 24% unique to type III (Table 1). The genetic distance between types I and II is therefore similar to the distance between type I and type III, and type II and type III. In addition, even though the 16S rRNA gene of *P. acnes* demonstrates a high degree of intra-species sequence identity, the observation of distinct and non-overlapping ribotypes for type I (RT1; RT3; RT4; RT5; RT8; RT16; RT532), type II (RT2; RT6), and type III (RT9) provides further evidence for their different phylogenies (Fitz-Gibbon et al., 2013, Barnard et al., 2016) (Fig S1).

Alongside phylogenetic analyses, previous whole genome typing patterns based on methods such as Random Amplification of Polymorphic DNA (RAPD) and noncoding repeat sequences, as well as the analysis of non-core regions, also support types I and II as highly distinct divisions at the genome level (Perry et al., 2003, Tomida et al., 2013, Hauck et al., 2015). While digital or *in silico* DNA-DNA hybridization values (GGDC 2.0 algorithm) between types I, II and III are above the 70% cut-off value currently used for bacterial species demarcation, thus confirming their membership of the same species, the whole genome relatedness values

are consistent with the proposal that types I and II are also placed in distinct taxonomic ranks in line with that recently proposed for type III (Dekio et al., 2015). Strains representing the different phylogroups within type I (IA₁, IA₂, IB, IC) share high *in silico* DNA-DNA hybridization values of 91-100%, but this drops to 74.1-78.5% when analysed against the type II strains ATCC11828 and JCM18920, and 72.0-72.8% with the type III strain JCM18909 (Dekio et al., 2015). Strains of type II and III share relatedness values of 72.9-73.2% (Dekio et al., 2015). These hybridization values between the major divisions are within the 70-80% similarity boundary recently recommended for bacterial subspecies (Meier-Kolthoff et al., 2014).

Detailed comparative analysis of type I and II whole genome sequences also reveals some salient differences between the divisions. These include specific genomic inversions and insertions present in type II strains, but not type I, which encode genes related to carbohydrate processing and modification, ABC transporters, nickel import, bacitracin resistance and hypothetical proteins (Fig. S2) (McDowell et al., 2013, Scholz et al., 2016). One of the most striking differences relates to the presence in type II strains of clustered regularly interspaced short palindromic repeats (CRISPR)/Cas locus (Brüggemann et al., 2012b, Fitz-Gibbon et al., 2013). In contrast, type I and type III strains contain CRISPR/Cas gene remnants within their genome, indicating deletion of the locus during the evolutionary history of these phylogroups; the deletions are more extensive in type I strains compared to type III. The deletion of the CRISPR/Cas system in type I and type III strains makes these divisions more susceptible to horizontal gene transfer (HGT) and the acquisition of fitness or virulence traits. The observation of such CRISPR/Cas gene remnants has led to the suggestion that the type I and III divisions may constitute younger subpopulations than type II strains which are descended from a more ancient lineage (Brüggemann et al., 2012a, Brüggemann et al.,

2012b). Since $\text{age} = ds/(\text{clock rate} \times 2)$, where ds is the mean number of synonymous substitutions per site and clock rate is the synonymous molecular clock rate, calculation of the ds values for strains currently representing the major type I, II and III divisions may give deeper insights into their relative ages. Interestingly, using the Nei-Gojobori method (Jukes-Cantor) (Nei and Gojobori, 1986) in MEGA v7.0 (Kumar et al., 2016), we observed that the ds value for the entire type I division was slightly higher than type II based on an initial analysis of concatenated MLSA₈ sequence data, while type III values were lower (Table 1). To investigate this further, we examined the shared core-coding regions of 85 *P. acnes* genomes currently available. Multiple sequence alignments were performed using MUSCLE (Edgar, 2004) and the Jukes-Cantor ds values calculated for each pair of sequences in the alignment using the Nei-Gojobori method as implemented in the Bioperl package Bio::Align::DNASTatistics (Stajich, 2002). As before, the resulting ds values obtained for type I (0.008), type II (0.005) and type III (0.001) revealed higher synonymous nucleotide diversity within the large type I clade compared to type II and type III, indicative of an older age. Further studies are therefore required to provide clarity on the series of evolutionary events that have given rise to the emergence and diversity of the current *P. acnes* clades now proposed as subspecies, including the possible diversity-purging effects of periodic selection (Cohan, 2001).

Propionibacterium acnes has a clonal, epidemic population structure and is in linkage disequilibrium, though rates of HGT within the population as a whole are statistically significant (Lomholt and Kilian, 2010, McDowell et al., 2012, McDowell et al., 2013). Previous studies have, however, found that rates of recombination appear to differ throughout the population, and that the association of alleles is less significant when distinct phylogroup

populations are considered (McDowell et al., 2012, McDowell et al., 2013). In particular, we see a drop in the index of association value (I_A) when strains from the type I and II divisions are considered separately, indicating increasing linkage equilibrium within these distinct clusters (McDowell et al., 2013); this can also be observed on a Neighbour-Net split graph based on MLST₈ allelic profile data (Fig. 2). Detailed inspection of MLSA₈ datasets also suggests conjugal transfer and replacement of unusually large chromosomal segments in the genome dynamics of the type I clade, particularly between types IA₂ and IB (Lomholt and Kilian, 2010, McDowell et al., 2012, McDowell et al., 2013). The idea that rates of genetic interchange are more frequent within, but not between, the major divisions suggests increasing sexual isolation which occurs with more genetically divergent organisms (Majewski, 2001). Reduced rates of recombination may also indicate ecological differences since members of the same habitat are more likely to undergo recombination events (sympatric speciation); such population subdivisions can introduce linkage disequilibrium into an analysis if isolates from different niches (Ecotypes) are included (Spratt and Maiden, 1999). Comprehensive analysis of genome differences between the major types does indeed provide potential evidence for distinct environmental challenges within the human host.

Studies by Johnson and Cummins (1972) first revealed types I and II as distinct phenotypes of *P. acnes* based on serological agglutination tests and cell wall sugar analysis; type I strains contain galactose in their cell wall, but this sugar is absent in type II strains which occasionally also contain *meso*-Diaminopimelic acid (DAP) (Table 2). The development of more recent monoclonal antibody typing methods for *P. acnes* have further highlighted differences between the cell wall structures of type I and II, as well as type III, based on the expression of unique antigenic determinants, including those in lipoteichoic acid and adhesin proteins

(Holland et al., 2010, McDowell et al., 2011, Bae et al., 2014). Differences in cell surface hydrophobicity have also been described for types I and II, and upon growth in liquid media, such as protease peptone yeast (PPY) or brain heart infusion (BHI) broth, type II strains form a turbid solution with a slight fine sediment, while strains of type IA and IC can form a large granular sediment or auto-aggregate with a clear solution (Cohen et al., 2005); type IB strains behave as type II in respect to this characteristic. Types I and II can be differentiated from one another and type III based on the analysis of bacterial whole cell proteins by matrix assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) fingerprinting, highlighting further variation at the phenotype level (Nagy et al., 2013, Dekio et al., 2015). Furthermore, differences in the susceptibility of types I and II to bacteriophage infection have also been known for some time (Webster and Cummins, 1978, Liu et al., 2015). The main phylogroups of *P. acnes* share a high degree of similarity with regard to their biochemical phenotype, including traditional tests used to differentiate the bacterium from other 'cutaneous' propionibacteria (Table 2). Notable phylogroup differences, however, include β -haemolytic and neuraminidase activity, as well as sorbitol fermentation, all of which are essentially restricted to the type I division (McDowell et al., 2008, Lomholt and Kilian, 2010, Niazi et al., 2010) (Table 2). The production of lipase also appears much lower amongst type II strains versus those from the type I and III divisions (McDowell et al., 2008, Niazi et al., 2010) (Table 2); we previously described how type II strains have deletions in the TATA box and open reading frame of two candidate lipase genes which may explain this reduced activity (Tomida et al., 2013).

One defining difference between the type I and II phylogroups rests on their association with acne vulgaris. On the basis of both culture and metagenomic analyses, widely disseminated

205 clonal lineages from the type I division have been described in association with acneic skin,
206 but not those from the type II or type III divisions which appear to be associated more with
207 blood, medical device and soft tissue infections (Lomholt and Kilian, 2010, McDowell et al.,
208 2011, McDowell et al., 2012, Fitz-Gibbon et al., 2013, Rollason et al., 2013). Recently, type III
209 strains have also been linked with the depigmenting skin condition progressive macular
210 hypomelanosis (Peterson et al., 2015, Barnard et al., 2016). Interrogation of the *P. acnes*
211 MLST₈ isolate database, which contains information on a large collection of geographically
212 widespread isolates and their clinical source, reveals a statistically significant enrichment
213 overall for strains from the type I clade in acneic versus healthy skin ($p < 0.001$; Fishers exact
214 test, two tailed), while those from the type II clade appear to show no association overall
215 ($p = 0.213$; Fishers exact test). More specifically, associations are found between acneic skin
216 and strains from the type IA₁ clonal complexes CC1 (RT1 and RT532) ($p < 0.01$; Fishers exact
217 test), CC3 (RT1, RT4 and RT5) ($p = 0.043$; Fishers exact test) and CC4 (RT8) ($p = 0.021$; Fishers
218 exact test) (Fig 1 and S1). In a previous study, we found that a globally disseminated clonal
219 lineage with the MLST genotype ST6 (Warwick MLST₇ scheme analysis) or ST1 (MLST₈ analysis)
220 strikingly represented the majority of type IA₁ isolates we analysed from a cohort of patients
221 with acne (McDowell et al., 2011). In contrast, specific type II lineages (RT2 and RT6) belonging
222 to CC72 (MLST₈) appear associated with healthy skin based on metagenomic and culture-
223 based detection (McDowell et al., 2012, Fitz-Gibbon et al., 2013, Johnson et al., 2016). The
224 observation that type II strains, but not those from the type I clade, encode CRISPR/Cas
225 elements may be important in this context, thus preventing the acquisition of genetic loci that
226 may contribute to virulence and acne pathophysiology (Fitz-Gibbon et al., 2013). For example,
227 key type I lineages from CC3 (MLST₈; Fig. 1), believed to be associated with acne contain a
228 novel plasmid with a tight adhesion (Tad) locus and two unique genomic islands, known as

loci 1 and 2, that contain genes proposed to enhance virulence via increased bacterial adhesion and host immune response (Fitz-Gibbon et al., 2013, Tomida et al., 2013, Kasimatis et al., 2013).

To conclude, we now emend the description of *Propionibacterium acnes* subsp. *acnes* (Deiko et al., 2015) as type I only, and propose type II strains as *Propionibacterium acnes* subsp. *defendens* subsp. nov. based on a polyphasic taxonomy approach. The growing number of genomes now becoming available for other propionibacteria will also provide an important opportunity to re-examine the genus.

Emendation of *Propionibacterium acnes* subsp. *acnes* (Deiko et al., 2015).

Description as given by Deiko et al. (2015), emended as follows:

Four phylogenetically distinct *Propionibacterium acnes* subsp. *acnes* (type I) groups have been described, known as type IA₁, IA₂, IB and IC; type IA₂, IB and IC represent phylogenetically tight clusters compared to IA₁. Cells are Gram-positive, nonmotile, non-spore forming, and anaerobic-to-aerotolerant. Colonies appear as lenticular, minute-to-4.0 mm, white, can become tanned, pink or orange in 3 weeks. Growth is most rapid at 30-37°C. Surface colonies on blood agar (horse or rabbit) are punctiform-to-0.5 mm, circular, entire-to-pulvinate, translucent-to-opaque, white-to-gray, glistening. The cell shape after anaerobic culture in broth medium ranges from small plump rods to ellipsoids which tend to occur in pairs joined at a slight angle, and the size is approximately 0.4-to-0.5 by 0.8-to-0.9 µm. In defined medium broth culture, type IA and IC strains form a turbid suspension, while in PPY or BHI broth they form a settled granular sediment with a clear solution. In contrast, type IB strains form a slight fine sediment and turbid solution containing suspended cells. In suitable media with good

growth, the final pH is 4.5-5.0. Generally catalase positive, cultures need to be exposed to air for 1 h before testing. All strains have an absolute requirement for pantothenate, while thiamine, biotin and nicotinamide are stimulatory. Strains are co-haemolytic and variable for β -haemolytic activity and produce a number of extracellular enzymes including ribonuclease, neuraminidase, hyaluronidase, acid phosphatase, lecithinase and lipase. Strains of type IA produce relatively low levels of the putative co-haemolytic Christie-Atkins-Munch-Peterson (CAMP) factor 1, but type IB strains produce an abundance of this protein. The total quantity of acid (especially the proportion of lactic acid) produced from fermentable carbohydrates is highly variable. Cells ferment glucose, but not sucrose or maltose. Lactate is converted to propionate by most strains but only if the initial oxidation-reduction potential of the medium is sufficiently low, or if the initial growth rate is rapid. Sorbitol fermentation is a variable but defining characteristic of type I strains. Gelatin is hydrolysed, and most strains produce indole and reduce nitrate, but esculin is not hydrolysed. The major long chain fatty acid produced in thioglycolate cultures is 13-methyltetradecanoic acid (32-62%) and iso-C15:0 FAME is the predominant cellular fatty acid. Prominent mass ions obtained by MALDI-TOF mass spectrometry are at 3,589 Da and 7,179 Da. Peptidoglycan contains alanine, glutamic acid, glycine and LL-DAP. Cell wall sugars are glucose, mannose and galactose. Strains have been isolated from the human skin, oral cavity and genitourinary tract. Type IA₁ and IC strains are associated with acne vulgaris. The G+C content is ~ 60% based on whole genome sequencing analysis.

Description of *Propionibacterium acnes* subsp. *defendens* subsp. nov.

Propionibacterium acnes subsp. *defendens* (de.fen'dens L. part. adj. *defendens*, defending, guarding, protecting; referring to the fact that strains have an active CRISPR/Cas system which

guards or controls against foreign mobile genetic elements). Description based on McDowell et al. (2008), Niazi et al. (2010), Patrick and McDowell (2011), and Dekio et al. (2015).

Cells are Gram-positive, nonmotile, non-spore forming, and anaerobic-to-aerotolerant. Their cellular and colony morphology is similar to type I cells, but they may appear more coccoid and are most similar to previous descriptions for '*Corynebacterium parvum*' which is a synonym for *P. acnes*. In defined medium broth culture, strains form a slight fine sediment and turbid solution containing suspended cells. In addition to pantothenate, some strains require haem and vitamin K to grow. Biochemical phenotype is similar to type I strains but with some notable differences. Cells are negative for β -haemolysis, and neuraminidase and lipase activity is infrequently found. Abundant levels of CAMP factor 1 are produced; similar to that observed with strains of type IB. Sorbitol fermentation is negative. The predominant cellular fatty acid is iso-C15:0 FAME and prominent mass ions obtained by MALDI-TOF mass spectrometry are 3,628 Da and 7,258 Da. Peptidoglycan contains alanine, glutamic acid, glycine, LL-DAP, and occasionally *meso*-DAP. Cell wall sugars are mannose and glucose, but galactose is not present. Strains have been isolated from the human skin surface, oral cavity and genitourinary tract. Strains are rarely associated with acne vulgaris and some may be associated with skin health and others with opportunistic infection. The G+C content is ~ 60 % based on whole genome sequencing analysis.

The type strain of *Propionibacterium acnes* subspecies *defendens* subsp. nov. is ATCC11828 (=JCM 6473=CCUG 6369) isolated from a subcutaneous abscess (Genbank accession number NC_017550).

295 **Acknowledgments**

296 EB, JL and HL are funded by the NIH grant R01GM099530 from NIGMS awarded to HL. JL is
297 also supported by the Ruth L. Kirschstein National Research Service Award AI007323. This
298 work was also supported by a grant of £11.5M awarded to Professor Tony Bjourson from
299 European Union Regional Development Fund (ERDF) EU Sustainable Competitiveness
300 Programme for N. Ireland; Northern Ireland Public Health Agency (HSC R&D) & Ulster
301 University.

References

- Bae, Y., Ito, T., Iida, T., Uchida, K., Sekine, M., Nakajima, Y., Kumagai, J., Yokoyama, T., Kawachi, H., & other authors (2014). Intracellular *Propionibacterium acnes* infection in glandular epithelium and stromal macrophages of the prostate with or without cancer. *PLoS ONE* **9**: e90324.
- Barnard, E., Liu, J., Yankova, E., Cavalcanti, S. M., Magalhães, M., Li, H., Patrick, S. & McDowell, A. (2016). Strains of the *Propionibacterium acnes* type III lineage are associated with the skin condition progressive macular hypomelanosis. *Sci Rep* **6**, 31968.
- Brenner, D., Stanley, J. & Krieg, N. (2000). Classification of prokaryotic organisms and the concept of bacterial speciation. In: Boone, D. R., Castenholz, W. & Garrity, G. M. (eds.) *Bergey's manual of systematic bacteriology*. 2 ed. New York, NY: Springer.
- Brüggemann, H., Lomholt, H. B. & Kilian, M. (2012a). The flexible gene pool of *Propionibacterium acnes*. *Mob Genet Elements*, **2**, 145-148.
- Brüggemann, H., Lomholt, H. B., Tettelin, H. & Kilian, M. (2012b). CRISPR/cas loci of type II *Propionibacterium acnes* confer immunity against acquisition of mobile elements present in type I *P. acnes*. *PLoS ONE*, **7**, e34171.
- Cavalcanti, S. M., De França, E. R., Lins, A. K., Magalhães, M., De Alencar, E. R. & Magalhães, V. (2011). Investigation of *Propionibacterium acnes* in progressive macular hypomelanosis using real-time PCR and culture. *Int J Dermatol*, **50**, 1347-1352.
- Cohan, F. (2001). Bacterial species and speciation. *Syst Biol*, **50**, 513–524.
- Cohen, R. J., Shannon, B. A., Mcneal, J. E., Shannon, T. & Garrett, K. L. (2005). *Propionibacterium acnes* associated with inflammation in radical prostatectomy specimens: a possible link to cancer evolution? *J Urol*, **173**, 1969-1974.
- Dekio, I., Culak, R., Misra, R., Gaulton, T., Fang, M., Sakamoto, M., Ohkuma, M., Oshima, K., Hattori, M., & other authors (2015). Dissecting the taxonomic heterogeneity within *Propionibacterium acnes*: proposal for *Propionibacterium acnes* subsp. *acnes* subsp. nov. and *Propionibacterium acnes* subsp. *elongatum* subsp. nov. *Int J Syst Evol Microbiol*, **65**, 4776-4787.
- Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**, 1792–1797.
- Eishi, Y. (2013). Etiologic link between sarcoidosis and *Propionibacterium acnes*. *Respir Investig*, **51**, 56-68.
- Fitz-Gibbon, S., Tomida, S., Chiu, B. H., Nguyen, L., Du, C., Liu, M., Elashoff, D., Erfe, M. C., Loncaric, A., & other authors (2013). *Propionibacterium acnes* strain populations in the human skin microbiome associated with acne. *J Invest Dermatol*, **133**, 2152-2160.
- Hauck, Y., Soler, C., G  r  me, P., Vong, R., Macnab, C., Appere, G., Vergnaud, G. & Pourcel, C. (2015). A novel multiple locus variable number of tandem repeat (VNTR) analysis (MLVA) method for *Propionibacterium acnes*. *Infect Genet Evol*, **33**, 233-241.
- Holland, C., Mak, T. N., Zimny-Arndt, U., Schmid, M., Meyer, T. F., Jungblut, P. R. & Br  ggemann, H. (2010). Proteomic identification of secreted proteins of *Propionibacterium acnes*. *BMC Microbiol*, **10**, 230.
- Huson, D. H. & Bryant, D. (2006). Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol*, **23**, 254-267.

343 **Johnson, J. L. & Cummins, C. S. (1972).** Cell wall composition and deoxyribonucleic acid similarities
 344 among the anaerobic coryneforms, classical propionibacteria, and strains of *Arachnia propionica*. *J*
 345 *Bacteriol*, **109**, 1047-1066.

346 **Johnson, T., Kang, D., Barnard, E. & Li, H. (2016).** Strain-Level differences in porphyrin production and
 347 regulation in *Propionibacterium acnes* elucidate disease associations. *mSphere* 1:e00023-15.

348 **Kasimatis, G., Fitz-Gibbon, S., Tomida, S., Wong, M. & Li, H. (2013).** Analysis of complete genomes of
 349 *Propionibacterium acnes* reveals a novel plasmid and increased pseudogenes in an acne associated
 350 strain. *Biomed Res Int* **2013**, 918320

351 **Kumar, S., Stecher, G., & Tamura, K. (2016).** MEGA7: Molecular Evolutionary Analysis version 7.0 for
 352 bigger datasets. *Mol Biol Evol*, **33**, 1870-1874

353 **Lapage, S. P., Sneath, P. H. A., Lessel, E. F., Skerman, V. B. D., Seeliger, H. P. R. & Clark, W. A. (1992).**
 354 *International Code of Nomenclature of Bacteria (1990 revision). Bacteriological Code*, Washington
 355 D.C., ASM Press.

356 **Liu, J., Yan, R., Zhong, Q., Bangayan, N. J., Nguyen, L., Lui, T., Liu, M., Erfe, M.C., Craft, N., & other
 357 authors (2015).** The diversity and host interactions of *Propionibacterium acnes* bacteriophages on
 358 human skin. *ISME J*, **9**, 2078-2093.

359 **Lomholt, H. B. & Kilian, M. (2010).** Population genetic analysis of *Propionibacterium acnes* identifies
 360 a subpopulation and epidemic clones associated with acne. *PLoS One*, **5**, e12277.

361 **Majewski, J. (2001).** Sexual isolation in bacteria. *FEMS Microbiol Lett*, **199**, 161-169.

362 **McDowell, A., Barnard, E., Nagy, I., Gao, A., Tomida, S., Li, H., Eady, A., Cove, J., Nord, C. E. & Patrick,
 363 S. (2012).** An expanded multilocus sequence typing scheme for *Propionibacterium acnes*: investigation
 364 of 'pathogenic', 'commensal' and antibiotic resistant strains. *PLoS One*, **7**, e41480.

365 **McDowell, A., Gao, A., Barnard, E., Fink, C., Murray, P. I., Dowson, C. G., Nagy, I., Lambert, P. A. &
 366 Patrick, S. (2011).** A novel multilocus sequence typing scheme for the opportunistic pathogen
 367 *Propionibacterium acnes* and characterization of type I cell surface-associated antigens. *Microbiology*,
 368 **157**, 1990-2003.

369 **McDowell, A., Nagy, I., Magyari, M., Barnard, E. & Patrick, S. (2013).** The opportunistic pathogen
 370 *Propionibacterium acnes*: insights into typing, human disease, clonal diversification and CAMP factor
 371 evolution. *PLoS One*, **8**, e70897.

372 **McDowell, A., Perry, A. L., Lambert, P. A. & Patrick, S. (2008).** A new phylogenetic group of
 373 *Propionibacterium acnes*. *J Med Microbiol*, **57**, 218-224.

374 **McDowell, A., Valanne, S., Ramage, G., Tunney, M. M., Glenn, J. V., McLorinan, G. C., Bhatia, A.,
 375 Maisonneuve, J. F., Lodes, M., & other authors (2005).** *Propionibacterium acnes* types I and II
 376 represent phylogenetically distinct groups. *J Clin Microbiol*, **43**, 326-334.

377 **Meier-Kolthoff, J. P., Hahnke, R. L., Petersen, J., Scheuner, C., Michael, V., Fiebig, A., Rohde, C.,
 378 Rohde, M., Fartmann, B., & other authors (2014).** Complete genome sequence of DSM 30083(T), the
 379 type strain (U5/41(T)) of *Escherichia coli*, and a proposal for delineating subspecies in microbial
 380 taxonomy. *Stand Genomic Sci*, **9**, 2.

381 **Nagy, E., Urbán, E., Becker, S., Kostrzewa, M., Vörös, A., Hunyadkürti, J. & Nagy, I. (2013).** MALDI-
 382 TOF MS fingerprinting facilitates rapid discrimination of phylotypes I, II and III of *Propionibacterium*
 383 *acnes*. *Anaerobe*, **20**, 20-26.

384 **Nei, M. & Gojobori, T. (1986).** Simple methods for estimating the numbers of synonymous and
 385 nonsynonymous nucleotide substitutions. *Mol Biol Evol*, **3**, 418–426.

- Niazi, S. A., Clarke, D., Do, T., Gilbert, S. C., Mannocci, F. & Beighton, D. (2010). *Propionibacterium acnes* and *Staphylococcus epidermidis* isolated from refractory endodontic lesions are opportunistic pathogens. *J Clin Microbiol*, **48**, 3859-3869.
- Patrick, S. & McDowell, A. (2011). The Propionibacteriaceae. In: Goodfellow, M., Kämpfer, P., Busse, H.-J., Trujillo, M. E., Suzuki, K.-I., Ludwig, W. & Whitman, B. W. B. (eds.) *Bergey's Manual of Systematic Bacteriology*. 2 ed. New York, NY: Springer.
- Perry, A. L., Worthington, T., Hilton, A. C., Lambert, P. A., Stirling, A. J. & Elliott, T. S. (2003). Analysis of clinical isolates of *Propionibacterium acnes* by optimised RAPD. *FEMS Microbiol Lett*, **228**, 51-55.
- Petersen, R., Lomholt, H. B., Scholz, C. F. & Brüggemann, H. (2015) Draft genome sequences of two *Propionibacterium acnes* strains isolated from progressive macular hypomelanosis lesions of human skin. *Genome Announc.* **3**, 6.
- Rollason, J., McDowell, A., Albert, H. B., Barnard, E., Worthington, T., Hilton, A. C., Vernallis, A., Patrick, S., Elliott, T. & Lambert, P. (2013) Genotypic and antimicrobial characterisation of *Propionibacterium acnes* isolates from surgically excised lumbar disc herniations. *Biomed Res Int*, **2013**, 530382.
- Scholz, C. F., Brüggemann, H., Lomholt, H. B., Tettelin, H. & Kilian, M. (2016). Genome stability of *Propionibacterium acnes*: a comprehensive study of indels and homopolymeric tracts. *Sci Rep*, **6**, 20662.
- Scholz, C. F., Jensen, A., Lomholt, H. B., Brüggemann, H. & Kilian, M. (2014). A novel high-resolution single locus sequence typing scheme for mixed populations of *Propionibacterium acnes* *in vivo*. *PLoS One*, **9**, e104199.
- Spratt, B. G. & Maiden, M. C. (1999). Bacterial population genetics, evolution and epidemiology. *Philos Trans R Soc Lond B Biol Sci*, **354**, 701-710.
- Stajich, J.E. (2002). The Bioperl Toolkit: Perl Modules for the Life Sciences. *Genome Res*, **12**, 1611–1618.
- Tomida, S., Nguyen, L., Chiu, B. H., Liu, J., Sodergren, E., Weinstock, G. M. & Li, H. (2013). Pan-genome and comparative genome analyses of *Propionibacterium acnes* reveal its genomic diversity in the healthy and diseased human skin microbiome. *mbio*, **4**, e00003-13.
- Tunney, M. M., Patrick, S., Curran, M. D., Ramage, G., Hanna, D., Nixon, J. R., Gorman, S. P., Davis, R. I. & Anderson, N. (1999). Detection of prosthetic hip infection at revision arthroplasty by immunofluorescence microscopy and PCR amplification of the bacterial 16S rRNA gene. *J Clin Microbiol*, **37**, 3281-3290.
- Valanne, S., McDowell, A., Ramage, G., Tunney, M. M., Einarsson, G. G., O'Hagan, S., Wisdom, G. B., Fairley, D., Bhatia, A., & other authors (2005). CAMP factor homologues in *Propionibacterium acnes*: a new protein family differentially expressed by types I and II. *Microbiology*, **151**, 1369-1379.
- Webster, G. F. & Cummins, C. S. (1978). Use of bacteriophage typing to distinguish *Propionibacterium acne* types I and II. *J Clin Microbiol*, **7**, 84-90.

Table 1. Genetic characteristics of *P. acnes* phylogroups

Genetic Grouping	p-distance (core SNPs)			<i>ds</i> *	% unique core region SNPs
	Type I	Type II	Type III		
Type I	-	0.444	0.487	0.006 ± 0.001	26
Type II	0.444	-	0.470	0.005 ± 0.001	22
Type III	0.487	0.470	-	0.002 ± 0.001	24
Type I, II, III	-	-	-	0.024 ± 0.003	-

*Based on the analysis of concatenated MLSA₈ sequence data using the Nei-Gojobori method (Jukes-Cantor) in MEGA v5.0.

Table 2. Key phenotypic similarities and differences between type I, II and III strains

Characteristic*	Type I	Type II	Type III
Indole production	+	d+	+
Catalase activity	+	+	+
Nitrate reduction	+	+	d+
Gelatin liquefaction	+	+	-
Aesculin Hydrolysis	-	-	-
β -haemolysis (5d at 37°C)	d+	-	-
Neuraminidase	d+	-	-
Lipase	d+	d-	d+
L-pyrrolydonyl arylamidase	d+	d-	-
Pyruvate	d+	+	-
Fermentation of:			
Sorbitol	d+	-	-
Maltose	-	-	-
Sucrose	-	-	-
Glycerol	d+	d+	+
Ribose	d-	d+	-
Cell wall components			
Dermatan sulphate-binding adhesins	d+	-	-
A ₂ pm isomer	LL-	LL- (<i>meso</i>)	ND
Amino acids	Ala, Gly, Glu	Ala, Gly, Glu	ND
Sugars	Galactose, Glucose, Mannose	Glucose Mannose	

+90% isolates are positive; -90% isolates are negative; d+40-89% isolates positive; d-11-39% isolates are positive.

*Key phenotypic characteristics were compiled from the data of one or more of the following publications: McDowell et al. (2005), McDowell et al. (2008), Lomholt and Kilian (2010), Niazi et al. (2010), McDowell et al. (2011), Patrick and McDowell (2011), Dekio et al. (2015).

Figure Legends.

Fig. 1. Minimum evolution phylogenetic tree (MEGA v7.0) (Kumar et al., 2016) of concatenated gene sequences (4253 bp) from all STs currently represented in the MLST₈ database (<http://pubmlst.org/pacnes/>), and covering all major genetic divisions. Sequence input order was randomized, and bootstrapping resampling statistics were performed using 500 data sets. Bootstrap values ($\geq 70\%$) are shown on the arms of the tree. Horizontal bar represents genetic distance. CC= clonal complex.

Fig. 2. Neighbour-net split graph (SplitsTree v4.14.4) of allelic profiles from all STs currently represented in the MLST₈ database (<http://pubmlst.org/pacnes/>), and covering all major genetic divisions (Huson and Bryant, 2006). A distance matrix was generated from the allelic profile data and saved in NEXUS format for input to SplitsTree. Parallelogram formations indicative of recombination/ reticulation events are evident within the major type I and II divisions.

Figure 1




Figure 2

