



Moderating the relationship between diabetes distress and mastery: the role of depression and empowerment

McGuigan, K., Hill, A. J., Coates, V., OKane, M., Thompson, D., Ski, C., & McCay, D. (2021). Moderating the relationship between diabetes distress and mastery: the role of depression and empowerment. *Psychology, Health & Medicine*, 1-10. Advance online publication. <https://doi.org/10.1080/13548506.2021.1894343>

[Link to publication record in Ulster University Research Portal](#)

Published in:

Psychology, Health & Medicine

Publication Status:

Published online: 01/03/2021

DOI:

[10.1080/13548506.2021.1894343](https://doi.org/10.1080/13548506.2021.1894343)

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.



Moderating the relationship between diabetes distress and mastery: the role of depression and empowerment

Karen McGuigan, Alyson Hill, Vivien Coates, Maurice O’Kane, David R Thompson, Chantal F Ski & Deirdre McCay

To cite this article: Karen McGuigan, Alyson Hill, Vivien Coates, Maurice O’Kane, David R Thompson, Chantal F Ski & Deirdre McCay (2021): Moderating the relationship between diabetes distress and mastery: the role of depression and empowerment, Psychology, Health & Medicine, DOI: [10.1080/13548506.2021.1894343](https://doi.org/10.1080/13548506.2021.1894343)

To link to this article: <https://doi.org/10.1080/13548506.2021.1894343>



© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 01 Mar 2021.



Submit your article to this journal [↗](#)



Article views: 220



View related articles [↗](#)



View Crossmark data [↗](#)

Moderating the relationship between diabetes distress and mastery: the role of depression and empowerment

Karen McGuigan ^a, Alyson Hill ^b, Vivien Coates ^{c,d}, Maurice O’Kane ^c,
David R Thompson ^a, Chantal F Ski ^e and Deirdre McCay ^b

^aSchool of Nursing and Midwifery, Queen’s University Belfast, Belfast, UK; ^bSchool of Biomedical Sciences (NICHE), Ulster University, Coleraine, UK; ^cWestern Health & Social Care Trust, Londonderry, UK; ^dSchool of Nursing, Ulster University, Coleraine, UK; ^eIntegrated Care Academy, University of Suffolk, Ipswich, UK

ABSTRACT

Type 2 diabetes is a chronic condition primarily self-managed by the individual. Mastery is a protective factor linked to better control of chronic conditions, effective self-management and improved medication adherence. Mastery appears increasingly important as treatment regimens and self-management demands become more complex and burdensome. Diabetes distress negatively impacts self-management, glycaemic control and treatment adherence. Understanding the relationship between diabetes distress and mastery may provide opportunities to improve condition management and adherence. This relationship may be impacted by other factors affecting the individual’s perceived sense of control over their condition. This study examined the role of diabetes empowerment and depression in the relationship between diabetes distress and mastery. Data were drawn from a randomised controlled trial of 131 adults with type 2 diabetes transitioning to injection therapy. Participants completed measures of diabetes distress, mastery, depression and empowerment. Diabetes distress and depression were negatively associated with mastery, whilst diabetes empowerment was positively associated. A significant interaction effect ($b = .024$, $t(112) = 3.79$, $p = <.005$) confirmed the relationship between diabetes distress and mastery was moderated by depression. Findings highlight the additive deleterious effects of depression. Interventions to improve mastery among those living with type 2 diabetes should address diabetes distress and depression to optimise outcomes.

ARTICLE HISTORY

Received 4 July 2020
Accepted 17 February 2021

KEYWORDS

Type 2 diabetes; mastery; diabetes distress; depression; moderation

Introduction

Type 2 diabetes (T2D) is a condition predominantly managed by the individual (McSharry et al., 2019). To be effective self-managers, those with T2D need to make decisions regarding medication, diet, exercise and blood glucose monitoring (Skinner et al., 2006). Effective self-management is necessary to ensure appropriate glycaemic control and reduce diabetes-related complications (Wilkinson et al., 2014). Treatment advice for T2D advocates lifestyle and dietary change as the first option to

CONTACT Karen McGuigan, Email:  k.mcguigan@qub.ac.uk  School of Nursing and Midwifery, Queen’s University Belfast, 97 Lisburn Road, Belfast BT9 7BL.

© 2021 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

achieve optimal glycaemic control; if control remains poor, oral medications are prescribed, progressing to injection therapy i.e. GLP-1 receptor agonists or insulin (Edelman & Pettus, 2014). However, approximately 1 in 3 express a reluctance to commence prescribed injection therapy, with up to 80% discontinuing or interrupting treatment relatively quickly following initiation (Perez-Nieves et al., 2017, 2016; Peyrot et al., 2012). Poor engagement with injectable therapies corresponds to concerns around the impact of injectables on daily living, treatment complexity and the restrictive nature of injectable regimens (Allen et al., 2017). Individuals who feel they have little control over their T2D and are unable to reach treatment targets report less motivation to manage their condition often irrespective of other factors (Gonzalez et al., 2016).

Mastery is recognised as a health-related protective factor, with higher levels of mastery related to better control of chronic conditions (Roepke & Grant, 2011). Mastery is considered: the extent to which individuals perceive aspects of their lives to be under their control, their ability to manage these aspects in light of the challenges they bring and their capacity to take appropriate action to affect associated positive outcomes (Forgeard & Benson, 2019). Mastery is positively associated with improved health behaviours (O’Kearney et al., 2020). Individuals with higher levels of mastery are more likely to engage in better self-management, adherence to diabetes treatment plans and medication management (Daniel et al., 2001; Roepke & Grant, 2011). Therefore, understanding more about factors which contribute to an individual’s level of mastery, particularly when changes in medication regimen are demanded, may provide insight to improve engagement and adherence.

Whilst mastery may serve as a positive protective factor, other factors can adversely affect adherence and self-management (Assari & Lankarani, 2017; Gonzalez et al., 2016; Linetzky et al., 2017; Spain et al., 2016). Diabetes distress refers to the negative psychological effects of living with diabetes and is linked to poor self-management, poor glycaemic control and difficulties with adherence (Dennick et al., 2017; Linetzky et al., 2017). Due to these link interventions targeting diabetes distress are receiving further attention as potential methods to improve outcomes for those living with diabetes (Hessler et al., 2020). Understanding the impact of diabetes distress on mastery among a population of those transitioning to injection therapy would provide insights for clinicians tasked with improving condition management and adherence among this group. This cannot be explored as a simple direct relationship; other factors should be considered. Depression is often presented as a comorbid condition in those living with T2D and is associated with poorer clinical outcomes, initiation of injectable therapies, medication adherence and motivation which impacts negatively on effective self-management behaviours (Lee, 2015; Nefs et al., 2013). Feelings of disempowerment associated with their diabetes are a further indicator of medication resistance and poorer outcomes (Linetzky et al., 2017).

Mastery is important for effective self-management and may be more potent as management of T2D becomes more complex. Thus, understanding more about mastery and factors that impact this among a T2D population transitioning to injectable therapies has potential to provide additional insights for clinicians and educators aiming to improve engagement with and adherence to injectable therapies. With this in mind the aim of this study is to examine the role of depression and diabetes empowerment as

moderators in the relationship between diabetes distress and mastery among a sample of those with T2D as they transition from oral medications to injectables. Based on the existing literature it was hypothesised that the relationship between diabetes distress and mastery would be moderated by depression and empowerment, with the interaction between patients' distress levels at high levels of depression and low levels of diabetes empowerment reflecting greater reductions in mastery.

Materials and methods

This study used baseline data drawn from a randomised controlled trial (RCT) of a structured diabetes education intervention. Individuals were eligible to participate in the trial if they: had a diagnosis of T2D for one year or more; were aged over 25 years; had HbA_{1c} level of 58–100 mmol/mol; and required transfer from oral hypoglycaemic treatment to injectable therapies as judged by the diabetes healthcare team. Exclusion criteria meant individuals: newly diagnosed with T2D; with HbA_{1c} >100 mmol/mol; undergoing retinal photocoagulation therapy or renal dialysis treatment; in receipt of psychiatric support or clinical psychology input; within 3 months of a major event including MI, stroke or major surgery; within 3 months of diagnosis or treatment of a major coexisting medical condition, were not eligible for inclusion. All participants attended a secondary care diabetes service in Northern Ireland delivered by a consultant-led, multidisciplinary team comprising Diabetologists/Endocrinologists, Diabetes Specialist Nurses and Diabetes Specialist Dietitians. Eligible participants were identified by a member of this multi-disciplinary care team. Participants provided informed consent to participate. Ethical approval was granted by the Office of Research Ethics Committee in Northern Ireland, with governance from the Trust Research Governance Committee prior to the commencement of participant recruitment [15/NI/0091].

Participants

The sample comprised 131 participants diagnosed with T2D transitioning to injectable therapies for effective management of the condition. Mean diabetes duration was 10.4 years (HbA_{1c} M = 70.3 mmol/mol; SD = 12.31). Age ranged from 39 to 85 years (M = 62.3; SD = 8.8); 59.5% of participants were male.

Measures

Once eligible participants were identified and informed consent obtained, participants were asked to complete and return two questionnaires, one providing demographic information (see supplementary online materials for sample characteristics) and the second consisting of the following measures taken at baseline.

Diabetes distress

The Problem Areas in Diabetes Scale (PAID: Polonsky et al., 1995), comprising 20 statements, is a reliable and valid measure of emotional distress specific to diabetes (Huang et al., 2010; Polonsky et al., 1995; Welch et al., 2003). Scored on a 5-point Likert scale (0 = not a problem; 4 = serious problem), scores range from 0 to 100,

with higher scores indicative of higher levels of emotional distress (Welch et al., 1997).

Mastery

The Pearlin Mastery Scale (PMS: Pearlin & Schooler, 1978) is a reliable and valid, unidimensional measure used to assess the extent to which individuals feel they have control over stressful events in their lives (Pearlin et al., 1981; Pearlin & Schooler, 1978; Turner & Noh, 1988). The scale comprises seven items, with higher scores indicative of greater levels of mastery (Brady, 2003).

Depression

The Hospital Anxiety and Depression Scale (HADS: Zigmond & Snaith, 1983) is a reliable and valid measure for assessing anxiety and depression among hospital outpatients, including adults with diabetes (Bjelland et al., 2002; Lloyd et al., 2000). Comprising 14 items (7 items in each subscale) assessing the severity of anxiety and depression; scored on a 4-point Likert scale. Subscale scores range from 0 to 21, with scores of 0–7 considered normal, and scores ≥ 11 moderate to severe (Collins et al., 2009).

Diabetes empowerment

The Diabetes Empowerment Scale – Short Form (DES-SF: Anderson et al., 2003) is a brief, reliable and valid measure assessing overall diabetes-related self-efficacy (Anderson et al. 2003). The scale comprises eight items, scored on a 5-item scale (1 = strongly disagree; 5 = strongly agree). Higher scores are indicative of greater empowerment.

In the current sample, the reliability of the scores on each scale was found to be acceptable with Cronbach's alpha ranging from 0.95 (PAID) to 0.84 (HADSD).

Analysis

The aim of this study is to examine the role of diabetes empowerment (M_1) and depression (M_2) in the relationship between diabetes-specific distress (X) and mastery (Y). To test this, a moderated model (Figure 1) was specified and tested in SPSS using PROCESS (Hayes, 2013). PROCESS 'uses logistic regression-based path analytical framework for estimating direct and indirect effects in simple and multiple moderation models' (Hayes, 2018, p. 1). In PROCESS, R^2 is used to assess model fit (Hayes, 2013).

Results

Model summary

The model was found to be a good fit for the data, with included variables accounting for approximately 54% of the variance in mastery among the group: $R^2 = .539$, $F(5, 112) = 26.22$, $p = <.0001$.




Figure 1. Conceptual and statistical diagram of proposed moderation model.

Main effects

Regression coefficients (Table 1) for the model indicate diabetes distress ($b = -.249, t(5,112) = -3.71, p < .005$), diabetes empowerment ($b = .280, t(5,112) = 3.02, p < .005$) and depression ($b = -.980, t(5,112) = -5.73, p < .005$) are all statistically significant predictors of mastery. The main effects show diabetes distress and depression are negatively associated with mastery. Diabetes empowerment is positively associated with mastery.

Interactions

A significant interaction between diabetes-specific distress and depression was found ($b = .024, t(112) = 3.79, p < .005$), indicating the magnitude of the effect of diabetes distress on mastery depends on level of depression. No significant interaction effect was found for diabetes distress and empowerment. PROCESS provides additional outputs on the change in the variance explained in the outcome due to the addition of the interaction. Results show no significant change in variance explained by mastery due to the addition of the diabetes distress x diabetes empowerment interaction: $F(1,112) = .46,$

Table 1. Model summary: path coefficients for estimated model using PROCESS.

	Coefficient (std error)	T	P	LLCI;ULCI
Constant	30.213 (1.53)	19.71	$p = .000$	27.18;33.25
Diabetes specific distress (X)	-.249 (.07)	-3.71	$p < .001$	-.38;-.12
Diabetes empowerment (M ₁)	.280 (.09)	3.02	$p < .001$.10;.46
Diabetes specific distress x Diabetes empowerment (X M ₁)	.003 (.004)	.68	$p = .500$	-.00;.01
Depression (M ₂)	-.980 (.17)	-5.73	$p < .001$	-1.32;-.64
Diabetes specific distress x Depression (X M ₂)	.024 (.01)	3.79	$p < .001$.01;.04




Figure 2. Relationship between diabetes distress and mastery at different levels of the moderators.

$p = .50$, $\Delta R^2 = .001$. There was a significant increase in variance explained due to the interaction of diabetes distress \times depression: $F(1,112) = 14.40$, $p < .005$, $\Delta R^2 = .06$.

Figure 2 illustrates the interaction between diabetes distress and depression more clearly: increasing levels of distress, at increased levels of depression, are indicative of lower levels of mastery, regardless of the level of diabetes empowerment.

Discussion

This study used data drawn from an RCT of adults living with T2D transitioning to injectable therapies to examine the role of diabetes empowerment and depression in the relationship between diabetes distress and mastery. It was hypothesised that the relationship between diabetes distress and mastery would be moderated by depression and empowerment, in essence individuals experiencing diabetes distress, high levels of depression and low levels of empowerment would report the lowest levels of mastery. Using Hayes' (2013) PROCESS moderation model, this study evidenced that diabetes distress, diabetes empowerment and depression significantly impact levels of mastery. Direct relationships indicated diabetes distress and depression were negatively associated with mastery, in essence in this cohort, as diabetes distress or depression increase, mastery decreases. These relationships are consistent with the existing literature that highlights the negative impact of distress and depression (Dennick et al., 2016; Lee, 2015; Linetzky et al., 2017). Diabetes empowerment was positively associated with mastery, i.e. as diabetes empowerment increases in this grouping, so too does mastery.

The significant interaction between diabetes distress and depression highlights how the negative impact of diabetes distress on mastery is heightened by increasing levels of depression with this interaction creating greater reduction in mastery. Additionally, there was no significant interaction for diabetes empowerment and distress. Further findings confirmed any positive effect of diabetes empowerment on mastery appears to be eroded in the presence of diabetes distress and depression.

The literature evidences the importance of mastery in chronic conditions, positively impacting self-management and treatment adherence (Roepke & Grant, 2011). The results from the current analysis provide insight into the mechanisms by which mastery is impacted, with contributing factors possibly more salient for the cohort involved in this study. Evidence suggests that diabetes distress is chronic, in that if it remains unresolved it can become more pervasive as the condition progresses (Hessler et al., 2020). Those with T2D transitioning to injection therapy are a unique population as they usually have been living and coping with the condition for a longer duration, therefore diabetes distress may be more prominent among this group. The current findings provide the basis for a more comprehensive understanding of the relationship between distress and mastery, recognising the impact of depression which presents as a common diabetes comorbidity (Lee, 2015; Nefs et al., 2013). Unpacking these associations may allow for development of better self-management promotion, strategies and interventions for this population. Indeed, calls for interventions to address diabetes distress to improve self-management and associated outcomes for those with diabetes may not offer effective solutions for this population if efforts are not made to tackle depression also.

There are some limitations with this study which should be recognised. This study utilises baseline data from an RCT, as such provides a snapshot of the population. A lack of longitudinal assessment limits our understanding of the changing nature of distress, depression, empowerment and mastery among this population as they navigate the challenges of injectables. Although a cross-sectional approach is suitable for this type of moderation study, a longitudinal study would have allowed for evidencing of injectable therapy adherence as a covariate in the model. Secondly, the study relies on self-report measures for data collection and therefore may be more influenced by participants' subjective views.

Overall, this study identified the relationship between diabetes distress and mastery is moderated by depression, undermining the impact of diabetes empowerment. These findings have important implications for those involved in the care of those with T2D requiring injectables to achieve optimal glycaemic control. Within the diabetes specialist services, diabetes distress and empowerment may be modulated through specific education to promote understanding about managing the injections and the implications of this additional treatment or through conversations to help resolve individual worries. However, these results indicate depression moderates the relationship between distress and mastery, accordingly it is not enough to simply target distress; if the individual also presents with depression then this must also be treated before mastery improves. As treating depression falls beyond the current remit of the diabetes clinicians and educators, a multi-disciplinary approach may be merited.

Acknowledgments

None

Disclosure statement

No potential conflicts of interest were reported by the authors.

Funding

This work was supported by the HSC Research and Development Division of the Public Health Agency, Northern Ireland (grant number: EAT/4909/13)

ORCID

Karen McGuigan  <http://orcid.org/0000-0002-0607-7936>
Alyson Hill  <http://orcid.org/0000-0003-3134-2794>
Vivien Coates  <http://orcid.org/0000-0002-8069-2961>
Maurice O’Kane  <http://orcid.org/0000-0002-3949-2802>
David R Thompson  <http://orcid.org/0000-0001-8518-6307>
Chantal F Ski  <http://orcid.org/0000-0003-1324-2933>
Deirdre McCay  <http://orcid.org/0000-0002-4184-7481>

References

- Allen, N. A., Zagarins, S. E., Feinberg, R. G., & Welch, G. (2017). Treating psychological insulin resistance in type 2 diabetes. *Journal of Clinical & Translational Endocrinology*, 7, 1–6. <https://doi.org/10.1016/j.jcte.2016.11.005>
- Anderson, R. M., Fitzgerald, J. T., Gruppen, L. D., Funnell, M. M., & Oh, M. S. (2003). The Diabetes Empowerment Scale-Short Form (DES-SF). *Diabetes Care*, 26(5), 1641–1642. <https://doi.org/10.2337/diacare.26.5.1641-a>
- Assari, S., & Lankarani, M. M. (2017). Reciprocal associations between depressive symptoms and mastery among older adults: Black-white differences. *Frontiers in Aging Neuroscience*, 8, 279. <https://doi.org/10.3389/fnagi.2016.00279>
- Bjelland, I., Dahl, A. A., Haug, T. T., & Neckelmann, D. (2002). The validity of the hospital anxiety and depression Scale: An updated literature review. *Journal of Psychosomatic Research*, 52(2), 69–77. [https://doi.org/10.1016/S0022-3999\(01\)00296-3](https://doi.org/10.1016/S0022-3999(01)00296-3)
- Brady, T. J. (2003). Measures of self-efficacy, helplessness, mastery and control. *Arthritis Care & Research*, 49(S5), S147–S164. <https://doi.org/10.1002/art.11413>
- Collins, M. M., Corcoran, P., & Perry, I. J. (2009). Anxiety and depression symptoms in patients with diabetes. *Diabetic Medicine*, 26(2), 153–161. <https://doi.org/10.1111/j.1464-5491.2008.02648.x>
- Daniel, M., Rowley, K. G., Herbert, C. P., O’Dea, K., & Green, L. W. (2001). Lipids and psychosocial status in aboriginal persons with and at risk for Type 2 diabetes: implications for tertiary prevention. *Patient Education and Counseling*, 43(1), 85–95. [https://doi.org/10.1016/S0738-3991\(00\)00153-1](https://doi.org/10.1016/S0738-3991(00)00153-1)
- Dennick, K., Sturt, J., & Speight, J. (2017). What Is Diabetes Distress and How Can We Measure It? A Narrative Review and Conceptual Model. *Journal of Diabetes and Its Complications*, 31(5), 898–911. <https://doi.org/10.1016/j.jdiacomp.2016.12.018>

- Edelman, S., & Pettus, J. (2014). Challenges associated with insulin therapy in type 2 diabetes mellitus. *American Journal of Medicine*, 127(10), S11–6. <https://doi.org/10.1016/j.amjmed.2014.07.003>
- Forgeard, M. J. C., & Benson, L. (2019). Extracurricular involvement and psychological adjustment in the transition from adolescence to emerging adulthood: the role of mastery and creative self-efficacy. *Applied Developmental Science*, 23(1), 41–58. <https://doi.org/10.1080/10888691.2017.1288124>
- Gonzalez, J. S., Tanenbaum, M. L., & Commisariat, P. V. (2016). Psychosocial factors in medication adherence and diabetes self-management: Implications for research and practice. *American Psychologist*, 71(7), 539–551. <https://doi.org/10.1037/a0040388>
- Hayes, A. F. (2013). Introduction to mediation, moderation and conditional process analysis: a regression-based approach. The Guilford Press.
- Hayes, A. F. (2013). *Introduction to mediation, moderation and conditional process analysis: a regression-based approach*. The Guilford Press.
- Hayes, A. F. (2018). *The PROCESS macro for SPSS and SAS*. Retrieved from: on December 11th 2018 <http://www.processmacro.org/index.html>
- Hessler, D., Fisher, L., Polonsky, W., Strycker, L., Parra, J., Bowyer, V., Dedhia, M., & Masharani, U. (2020). There is value in treating elevated levels of diabetes distress: The clinical impact of targeted interventions in adults with Type 1 diabetes. *Diabetic Medicine*, 37(1), 71–74. <https://doi.org/10.1111/dme.14082>
- Huang, M. F., Courtney, M., Edwards, H., & McDowell, J. (2010). Validation of the Chinese version of the problem areas in diabetes (PAID-C) scale. *Diabetes Care*, 33(1), 38–40. <https://doi.org/10.2337/dc09-0768>
- Lee, K. P. (2015). Psychosocial factors associated with psychological insulin resistance in primary care patients in Hong Kong. *Journal of Clinical & Translational Endocrinology*, 2(4), 157–162. <https://doi.org/10.1016/j.jcte.2015.10.001>
- Linetzky, B., Jiang, D., Funnell, M. M., Curtis, B. H., & Polonsky, W. H. (2017). Exploring the role of the patient–physician relationship on insulin adherence and clinical outcomes in Type 2 diabetes: Insights from the MOSAIC study. *Journal of Diabetes*, 9(6), 596–605. <https://doi.org/10.1111/1753-0407.12443>
- Lloyd, C. E., Dyer, P. H., & Barnett, A. H. (2000). Prevalence of symptoms of depression and anxiety in a diabetes clinic population. *Diabetic Medicine*, 17(3), 198–202. <https://doi.org/10.1046/j.1464-5491.2000.00260.x>
- McSharry, J., Dineen, S. F., Humphreys, M., O'Donnell, M., O'Hara, M. C., Smith, S. M., Winkley, K., & Byrne, M. (2019). Barriers and facilitators to attendance at Type 2 diabetes structured education programmes: A qualitative study of educators and attendees. *Diabetic Medicine*, 36(1), 70–79. <https://doi.org/10.1111/dme.13805>
- Nefs, G., Pop, V. J., Denollet, J., & Pouwer, F. (2013). The longitudinal association between depressive symptoms and initiation of insulin therapy in people with Type 2 Diabetes in primary care. *PLoS ONE*, 8(11), e78865. <https://doi.org/10.1371/journal.pone.0078865>
- O'Kearney, E. L., Brown, C. R., Jelinek, G. A., Neate, S. L., Taylor, K. T., Bevans, W., De Livera, A. M., Simpson, Jr., S., Jr, & Weiland, T. J. (2020). Mastery is associated with greater physical and mental health-related quality of life in two international cohorts of people with multiple sclerosis. *Multiple Sclerosis and Related Disorders*, 38, 101481. <https://doi.org/10.1016/j.msard.2019.101481>
- Pearlin, L. I., Menaghan, E. G., Lieberman, M. A., & Mullen, J. T. (1981). The stress process. *Journal of Health and Social Behavior*, 22(4), 337–356. <https://doi.org/10.2307/2136676>
- Pearlin, L. I., & Schooler, I. (1978). The structure of coping. *Journal of Health and Social Behavior*, 19(1), 2–21. <https://doi.org/10.2307/2136319>
- Perez-Nieves, M., Ivanova, J. I., Hadjiyianni, I., Zhao, C., Cao, D., Schmerold, L., Kalirai, S., King, S., DeLozier, A. M., Birnbaum, H. G., & Peyrot, M. (2017). Basal insulin initiation use and experience among people with Type 2 diabetes mellitus with different patterns of persistence: Results from a multi-national survey. *Current Medical Research and Opinion*, 33(10), 1833–1842. <https://doi.org/10.1080/03007995.2017.1341403>

- Perez-Nieves, M., Kabul, S., Desai, U., Ivanova, J. I., Kirson, N. Y., Cummings, A. K., Birnbaum, H. G., Duan, R., Cao, D., & Hadjiyianni, I. (2016). Basal insulin persistence, associated factors, and outcomes after treatment initiation among people with Type 2 diabetes mellitus in the US. *Current Medical Research and Opinion*, 32(4), 669–680. <https://doi.org/10.1185/03007995.2015.1135789>
- Peyrot, M., Barnett, A. H., Meneghini, L. F., & Schumm-Draeger, P. M. (2012). Insulin adherence behaviours and barriers in the multinational global attitudes of patients and physicians in insulin therapy study. *Diabetic Medicine*, 29(5), 682–689. <https://doi.org/10.1111/j.1464-5491.2012.03605.x>
- Polonsky, W. H., Anderson, B. J., Lohrer, P. A., Welch, G., Jacobson, A. M., Aponte, J. E., & Schwartz, C. E. (1995). Assessment of diabetes-related distress. *Diabetes Care*, 18(6), 754–760. <https://doi.org/10.2337/diacare.18.6.754>
- Roepke, S. K., & Grant, I. (2011). Toward a more complete understanding of the effects of personal mastery on cardiometabolic health. *Health Psychology*, 30(5), 615–632. <https://doi.org/10.1037/a0023480>
- Skinner, T. C., Carey, M. E., Craddock, S., Daly, H., Davies, M. J., Doherty, Y., Heller, S., Khunti, K., & Oliver, L., on behalf of the DESMOND collaborative. (2006). Diabetes education and self-management for ongoing and newly diagnosed (DESMOND): process modelling of pilot study. *Patient Education and Counseling*, 64(1-3), 369–377. <https://doi.org/10.1016/j.pec.2006.04.007>
- Spain, C. V., Wirght, J. J., Hahn, M. P. H., Wivel, A., & Martin, A. A. (2016). Self-reported barriers to adherence and persistence to treatment with injectable medications for Type 2 diabetes. *Clinical Therapeutics*, 38(7), 1653–1664. <https://doi.org/10.1016/j.clinthera.2016.05.009>
- Turner, R. J., & Noh, S. (1988). Physical disability and depression: a longitudinal analysis. *Journal of Health and Social Behavior*, 29(1), 23–37. <https://doi.org/10.2307/2137178>
- Welch, G., Jacobson, A., & Polonsky, W. H. (1997). The problem areas in diabetes scale: an evaluation of its clinical utility. *Diabetes Care*, 20(5), 760–766. <https://doi.org/10.2337/diacare.20.5.760>
- Welch, G., Weinger, K., Anderson, B., & Polonsky, W. H. (2003). Responsiveness of the problem areas in diabetes (PAID) questionnaire. *Diabetes Medicine*, 20(1), 69–72. <https://doi.org/10.1046/j.1464-5491.2003.00832.x>
- Wilkinson, A., Whitehead, L., & Ritchie, L. (2014). Factors influencing the ability to self-manage diabetes for adults living with Type 1 or 2 diabetes. *International Journal of Nursing Studies*, 51(1), 111–122. <https://doi.org/10.1016/j.ijnurstu.2013.01.006>
- Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, 67(6), 361–370. <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x>