Cryotherapy and inflammation: evidence beyond the cardinal signs

Bleakley, C. M., & Davison, G. W. (2010). Cryotherapy and inflammation: evidence beyond the cardinal signs. *Physical Therapy Reviews*, 15(6), 430-435. <https://doi.org/10.1179/1743288X10Y.0000000014>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Physical Therapy Reviews

Publication Status:
Published (in print/issue): 01/12/2010

DOI:
[10.1179/1743288X10Y.0000000014](https://doi.org/10.1179/1743288X10Y.0000000014)

Document Version
Publisher's PDF, also known as Version of record

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Cryotherapy and inflammation: evidence beyond the cardinal signs

Chris M. Bleakley¹, Gareth W. Davison²

¹Health and Rehabilitation Sciences Research Institute and ²Sport and Exercise Sciences, Research Institute, University of Ulster, Northern Ireland

Background: Cryotherapy is one of the most popular electro-physical agents used to 'treat' acute inflammation after a soft tissue injury. Much of the clinical rationale for this is based on anecdotal reports, with most clinicians accepting that cryotherapy has an 'anti' inflammatory effect after injury. There have been a number of recent advances towards improving our understanding of the inflammatory process after soft tissue injury.

Objectives: To review the rationale for cryotherapy intervention in the acute phases of soft tissue injury, whilst considering physiological, cellular and molecular models of inflammation.

Methods: Qualitative review of recent evidence.

Results: Research is restricted to animal models, applying various forms of cryotherapy after induced soft tissue injury. Outcomes focus on the effect that cooling has on key physiological, biochemical and molecular inflammatory events including: secondary cell death, white blood cell behaviour, apoptosis, blood flow and oedema formation.

Conclusion: Cryotherapy can have an influence on key inflammatory events at a cellular and physiological level after an acute soft tissue injury. However, the relative benefits of these effects have yet to be fully elucidated and it is difficult to contextualize within a human model. It is important to continue to update our rationale for applying common electro-physical agents such as cryotherapy after acute soft tissue injury, based on contemporary models of inflammation.

Keywords: Cryotherapy, Acute injury, Soft tissue, Inflammation

Introduction

Cryotherapy is one of the simplest and oldest modalities for treating soft tissue injuries such as sprains, contusions, and dislocations. The immediate phase after soft tissue injury is characterized by an acute inflammatory response. This often presents clinically with cardinal signs such as heat, redness, pain and swelling. Few clinicians may look beyond these cardinal signs when providing justification for intervention; and it is commonly accepted that cryotherapy has an 'anti' inflammatory effect after soft tissue injury. Applying a cold agent to a hot and red tissue may seem pragmatic; however, there is not always an obvious link between inflammation visible under the microscope and that clinically apparent and characterized by the original cardinal signs.¹ Paradoxically, recent trends in sports medicine involve delivering growth factors into healing muscle tissue (e.g. via platelet rich plasma or autologous

blood injections)² which seems to lean more towards a pro-inflammatory treatment approach.

Our aim is to review the rationale for cryotherapy intervention in the acute phases of soft tissue injury, whilst considering physiological, cellular and molecular models of inflammation. Where appropriate, relevance to injured human subjects will be assessed, and recommendations for future research provided.

Search Strategy

In January 2010, we undertook a computerized literature search on Medline, EMBASE and Cochrane Central Register of Controlled Trials (CCTR) (via OVID) using nine key words and subject headings relating to cryotherapy. This was supplemented with 'related article' searches on PubMed, and bibliography tracking. Relevant studies were extracted, with exclusions made based on titles, abstracts or full text versions. No restrictions were made on study design or type/mechanism of acute soft tissue injury. Relevant outcomes were any physiological, cellular or molecular measurement associated with inflammation; recorded before injury, and up to one week post-injury. No restrictions were made on the type/dosage of cryotherapy intervention.

Correspondence to: Chris Bleakley, Health and Rehabilitation Sciences Research Institute, School of Health Sciences, University of Ulster, Shore Road, Newtownabbey, Co Antrim, BT37 0QB, Northern Ireland. Email: chrisbleakley@hotmail.com

Table 1 provides details of the cryotherapy interventions and reported outcomes across studies. Qualitative comparisons were made and results were grouped and discussed by outcome.

Secondary Cell Injury

Perhaps the most commonly cited rationale for applying ice after acute soft tissue injury relates to the 'secondary injury model'.³ This is based on the premise that after an initial trauma (e.g. muscle strain or contusion), the patho-physiological events associated with acute inflammation can induce secondary damage

to cells around the injury site. Of particular concern is that this can involve collateral damage to healthy cells not injured in the initial trauma. This phenomenon is known as secondary cell injury, and may be caused by both enzymatic and ischaemic mechanisms.⁴ One of the most important cellular effects associated with cryotherapy is its potential to reduce the metabolic rate of tissue at, and surrounding the injury site. This reduction in metabolic demand may allow the cells to better tolerate the ischaemic environment in the immediate phases after injury thus minimizing the potential for secondary cell injury or death.

Table 1 Details of cryotherapy intervention and reported outcomes

Author	Details of cryotherapy intervention	Directness of cooling	Time after injury of ice initiation	Outcomes (blinded assessor Y/N)
Osterman <i>et al.</i> ⁵	CWI (ice and isotonic saline); duration up to 13 hours (until ATP depletion)	Amputated limb with a single layer of plastic wrap (mean intramuscular temperature: 1.2°C)	Immediately post amputation	ATP/PCr depletion (N)
Sapega <i>et al.</i> ⁶	CWI (isotonic saline at 1, 10, 15 and 22°C); duration 45 minutes, up to 13 hours	Amputated limb with a single layer of plastic wrap (intramuscular temperature as low as 0.5–1°C)	Immediately post amputation	ATP/PCr depletion; pH (N)
Farry <i>et al.</i> ¹⁷	Crushed and compression, 20 minutes × 2	Intact skin	Immediate (likely)	IHA (Y)
Hurme <i>et al.</i> ¹⁸	Cold pack with compression and elevation; 5 minutes every 14 × 4	Intact skin (lowest temperature recorded in deep muscle: 20°C)	Immediate	IHA (Y)
Smith <i>et al.</i> ²²	Ice cylinders; 20 minutes every 6 hours × 3	Intact unshaven skin	Immediate (likely)	Intravital microscopy with MC (Y) Laser Doppler fluxmetry
Curl <i>et al.</i> ²⁹	Ice cylinders; 20 minutes every 6 hours for 2 days	Intact skin	5 minutes	Intravital microscopy with MC Laser fluxmetry (N)
Dolan <i>et al.</i> ³⁰	CWI in 12.8–15.6°C; 30 minutes × 4	CWI to intact shaven limbs	5 minutes	Water displacement (N)
Merrick <i>et al.</i> ⁷	Ice pack with elastic tape; 5 hours	Unshaven intact skin	Immediate (likely)	Biochemical assay (N)
Westermann <i>et al.</i> ¹³	Ice cold saline solution; 1 hour duration	Through MC, muscle surface temperature decreased to 10 ± 2°C	Immediate	Intravital microscopy with MC (N)
Deal <i>et al.</i> ³¹	Cylinder of ice to skin side of MC; 20 minutes	Unshaven intact skin	15 minutes	Intravital fluorescent microscopy with MC (N)
Dolan <i>et al.</i> ³⁰	CWI at 12.8°C; 3 hours, followed by 1 hour rest	CWI intact shaven limbs	5 minutes	Water displacement (N)
Lee <i>et al.</i> ¹⁴	Saline at 3°C; 10 minutes	Cooling directly onto exposed muscle surface	5 minutes	Intravital microscopy (N) Real time laser scanning
Schaser <i>et al.</i> ¹⁵	Saline at 8°C; 20 minutes	Direct to surgically exposed muscle (muscle surface temperature cooled to 10°C)	Immediate (likely)	Intravital microscopy IHA (Y)
Schaser <i>et al.</i> ¹⁶	Saline at 8°C; 6 hours	Shaven intact skin (muscle surface temperature cooled to 10°C)	Immediate (likely)	Intravital microscopy IHA (Y)

Note: CWI, cold water immersion; IHA, immunohistological analysis; MC, microvascular chamber; ATP, adenosine triphosphate; PCr, phosphocreatine.

Immediate (likely): although not stated specifically, it was likely based on the experimental set-up.

Evidence to support secondary injury theory is based largely on studies of limb preservation. Sapega and colleagues^{5,6} used phosphorous-31 nuclear magnetic resonance imaging to monitor cellular metabolism in ischaemic (amputated) cat limbs, stored for up to 10 hours, at a range of temperatures between 22 and 1°C. Limbs were removed at hourly intervals for rescanning; overall, results showed that muscle cells survived better at lower muscle temperatures. This was exemplified by lower levels of adenosine triphosphate (ATP) and phosphocreatine depletion, and lower levels of acidosis, during the period of ischaemia. Of note, these effects appeared to be reversed at more extreme muscle temperature reductions below 5°C. This was attributed to extreme temperature reductions causing inhibition of the calcium pump of the muscle's sarcoplasmic reticulum.⁶

In a more recent and related study, Merrick and colleagues⁷ tried to quantify the effect of cryotherapy on mitochondrial function after injury. Specifically they measured the activity of the mitochondrial enzyme, cytochrome c oxidase, after experimental crush injury; comparing outcomes in cold treated and untreated muscle tissue. Fitting with the 'secondary injury model', 5 hours of continuous cooling with a crushed ice pack inhibited the loss of mitochondrial oxidative function after injury when compared to the untreated controls. Although the model used by Merrick *et al.*⁷ is not directly determining the effects of cryotherapy on the inflammatory process or muscle injury per se, it is the first study to have taken a novel approach to indirectly assess the effects of secondary generated free radicals, and their possible interference with enzymes controlling oxidative phosphorylation (cytochrome c oxidase) and thus ATP production after injury.

White Blood Cells (WBCs)

When muscle or joint injury occurs, phagocytic white cells, such as neutrophils, monocytes, eosinophils and macrophages become activated and dominate the inflammatory response in the early stages. Although these cells have a critical role in healing through their removal of necrotic debris and release of cytokines;⁸ they can also have a negative effect on soft tissue healing after injury.^{8,9} For example, white cell activation results in a series of reactions termed the 'respiratory burst'.¹⁰ These reactions are a source of reactive oxygen species (ROS) such as superoxide (O_2^-), hydrogen peroxide (H_2O_2) and hydroxyl ($OH^\cdot$); and hypochlorous acid ($HOCl$) which is a powerful antibacterial agent. In certain circumstances the production of ROS and antibacterial agents are important immune defense mechanisms; however, they can also be a potentially dangerous mechanism if

inappropriately activated. For example, overproduction of ROS may cause unwanted collateral damage to adjacent tissues and surrounding molecules.¹¹ This may be particularly likely in the event of a closed soft tissue injury such as an ankle sprain, which is not associated with bacteria or infection. Indeed, there is evidence that blocking the respiratory burst, using anti-CD11b antibody (M1/70), produces a three-fold reduction in myofibre damage in an animal model at 24 hours post-injury.¹²

Interestingly, a number of animal models have studied the effect that cryotherapy has on WBC behaviour after soft tissue injury. A popular approach has been to use fluorescent intravital microscopy¹³⁻¹⁶ to observe the effect that ice has on leukocyte activity within the microvasculature. These studies found a clear trend that icing significantly lowered the percentage of both adherent and rolling neutrophils after injury, in comparison to injured untreated tissue. This finding was consistent over the first 24 hours after injury.¹³⁻¹⁶

Other animal models^{7,15-18} have undertaken histological analysis on excised tissue after soft tissue injury. In each case, various staining techniques were used to identify leukocyte sub-types at the injury site. Again each study made comparisons between ice treated, and untreated injured tissue samples. Using an injured ligament model and assessor blinding, Farry *et al.*¹⁷ found that ice treated groups had lower levels of WBCs (polymorphs, lymphocytes and plasma cells) at 48 hours, in comparison to injured contra-lateral untreated limbs. Hurme *et al.*,¹⁸ who also used blinded outcome analysis, found that at various time points post-injury, the ice treated animal tissue had lower levels of erythrocytes (1 hour), neutrophils (6 hours) and macrophages (at 24 hours) in comparison to the untreated control limbs. Although Schaser *et al.*¹⁵ also found cooling decreased neutrophilic granulocyte muscle infiltration, in comparison to control muscle, there were higher levels of macrophages. In a follow-up study¹⁶ using longer periods of cooling (5 hours), tissue analysis at 24 hours post trauma also found lower levels of neutrophilic granulocytes in the cold treated muscle.

Although the examination of adherent and rolling neutrophils following injury and cryotherapy may, in some instances, be beneficial, these models must be developed if we are to further our understanding in this area. It may be more relevant for future research to quantify the amount of direct neutrophil activation that occurs following injury. This approach may allow for estimation as to how much secondary cell and surrounding tissue damage and inflammation will likely occur. A popular marker that is commonly used to determine neutrophil activation is myeloperoxidase.

This is produced by an increase in ROS activity and it has been successfully used in studies looking at free radical production and immune response after stretch injury in animal skeletal muscle.¹⁹

Apoptosis

Apoptosis is a programmed cell death. It is characterized by a cascade of biochemical events cumulating in altered cell morphology and eventual cell death. Although apoptosis is the normal means by which cells die at the end of their life span, its incidence may be affected by soft tissue injury. Higher numbers of apoptotic cells have been recorded around the edges of rotator cuff tears when compared to un-injured control muscles.²⁰ The reasons for this increase have not yet been fully elucidated; however, the accumulation of reactive oxygen species in injured tissue (oxidative stress) could again play a significant role.²¹ Cell survival requires multiple factors, including appropriate proportions of molecular oxygen and various antioxidants. Although most oxidative insults can be overcome by the cell's natural defenses, sustained perturbation of this balance may result in apoptotic cell death.

There is limited evidence from animal models that cryotherapy can reduce the incidence of apoptosis after injury. Westermann *et al.*¹³ found that after chemically induced inflammation, the number of apoptotic muscle cells (quantified by the number with nuclear condensation and fragmentation) was significantly higher in untreated controls, when compared to the cryotherapy group (muscle surface cooled to 10°C). This is an interesting finding as reduced levels of apoptosis may again represent a protective effect of cryotherapy after soft tissue injury. We can only postulate as to the reasons for this finding; however, this may be further evidence that cryotherapy can reduce inflammation and decrease secondary free radical production (from the respiratory burst), thereby causing less interference with important proteins and other cell metabolites that control apoptosis.

Blood Flow and Oedema

Acute soft tissue injury incurs a multitude of changes to the microvasculature. These include: increased vessel diameter;^{16,22} increased cell permeability and macromolecular leakage into the injured tissue;¹⁶ and decreased tissue perfusion.^{13,15,16} Paradoxically, there is clear evidence that ice has a vasoconstrictive response in human tissue based on impedance plethysmography outcomes.^{23,24} Recent studies^{25–28} also confirm that topical cooling has a similar effect on deep tissue haemodynamics, causing significant reductions in capillary blood flow, with facilitated venous capillary outflow in healthy humans. It is important to consider

if cryotherapy can reverse the effects that soft tissue injury has on tissue haemodynamics using injured models. Again much of the evidence in this area is based on animal models.

Using intravital microscopy, some studies found that ice application did not significantly change capillary diameter,^{15,16} arteriole diameter,^{15,22,29} or capillary velocity after injury.^{15,16} In contrast, others found that ice either significantly increased¹⁵ or decreased¹³ arteriole diameter after injury.

There may be clearer patterns associated with venular diameters. Three studies^{14–16} reported smaller venular diameters in ice treated groups in comparison to injured (untreated) controls when measured at both the initial stages^{14,15} and at 24 hours¹⁶ post-injury. In two of these studies,^{14,16} the differences were significant, and in one case, venular diameter in the cold group had returned to pre-injury levels.¹⁶ Although this trend is supported with evidence that iced tissue also had higher levels of venular blood flow velocity in comparison to controls in the immediate stages post-injury,^{13–15} this trend was reversed at 24 hours post-injury.

There is also conflicting evidence on the effect that cryotherapy has on tissue perfusion post-injury. Using fluorescent microscopic assessment of the functional capillary density (length of erythrocyte-perfused capillaries per observation area), three studies^{13,15,16} found that ice application significantly increases tissue perfusion after injury, in comparison to untreated injured controls. Again, in two cases,^{15,16} perfusion was restored to pre-injury levels. In contrast, based on laser Doppler imaging after injury, Curl *et al.*²⁹ found that cooling had little effect on microvascular perfusion.

Using a related outcome measurement, Schaser and colleagues^{15,16} monitored intramuscular pressures in rat limbs after soft tissue injury, randomizing limbs to receive either cold saline, or no intervention. Lower intramuscular pressures [17.7 mm Hg (SD: 4.7)] were recorded at 1.5 hours post-injury in the cooling group (treated with 20 minutes of saline cooling), when compared to untreated controls [19.2 mm Hg (SD: 3.1)]. Their follow-up study also found that longer periods of muscle cooling (5 hours) was associated with lower intramuscular pressures¹⁸ (95% CI: 5.5 mm Hg) in comparison to untreated controls [26 (95% CI: 1.9 mm Hg)], at 24 hours post-injury.

Limitation and Future Study

There are a number of shortcomings associated with the current evidence base, particularly when we try to relate these findings to the injured human subject. Primarily, much of the research relies on animal models, applying various forms of cryotherapy after an induced 'crush type' soft tissue injury. Furthermore,

in the majority of cases, the severity of the injury was controlled to avoid excessive haemorrhage or damage to major blood vessels,^{14,22,30,31} and is therefore not applicable to more serious soft tissue injuries presenting clinically. Similarly, the majority of studies have used a muscle contusion model, which on a mechanical, anatomical and pathological level, is different to a stretch type muscle injury or a ligament sprain.

The use of amputated or excised tissues samples has obvious drawbacks, when compared to models using perfused tissue. We must also consider that many of the *in vivo* models in this area used heavy anaesthesia on the animals throughout the experiment. Anaesthesia can alter tissue perfusion by as much as 38%,²² which could clearly confound findings in an inflammatory study. Furthermore, an induced injury on an anaesthetized animal means that the muscle will be in a relaxed state. If the muscle is contracted at the time of contusion injury, which is usually the case in the sporting environment, we might expect a significantly different impact response and force displacement.^{32,33}

It is clear that cryotherapy can have an effect on key inflammatory events at a cellular and physiological level. However, the relative benefits of these effects have yet to be fully elucidated and it is currently difficult to contextualize within a clinical model. Most of the work completed to date has focused on the animal model, perhaps due to its relative ease with regard to obtaining tissue. We must consider that the temperature reductions reported in animal tissues are extremely low (intra-muscular temperatures of 1–10°C), and usually obtained within 15 minutes of injury. These tissue temperatures are not only difficult to replicate in a human model, but practically, cooling is usually initiated hours or days after injury, particularly within randomized controlled studies. Notwithstanding this, there needs to be more emphasis placed on the effects of cryotherapy on inflammation and muscle injury using a human approach and model. Much work needs to be done using an array of peripheral and muscle specific markers to determine and quantify the inflammatory process *per se*. Considerable new information and knowledge within cryotherapy may be obtained by directly examining inflammatory related markers such as high sensitive C-reactive protein, tumour necrosis factor alpha, nuclear factor kappa B and interleukin molecules such as interleukin-6. Investigators should also be encouraged to examine markers of cellular oxidative stress in order to determine the relationship between primary and secondary free radical production (caused by soft tissue injury) and inflammation with and without cryotherapy application.

Considerations must also be given to the type of soft tissue affected. Muscle, ligaments and tendon tissue may also have different levels of tolerance when faced with post-injury ischaemia, and some may be more at risk of secondary cell injury and collateral damage. We must also clarify whether the potential benefits of cryotherapy are restricted to reducing or preventing secondary injury in cells not initially damaged by primary trauma, or if they could also target retardation of primary injury progression, i.e. rescuing cells that were involved in primary injury but not initially destroyed.⁴ This could also have considerable implications for clinical management of soft tissue injury.

References

- 1 Scott A, Khan KM, Roberts CR, Cook JL, Duronio V. What do we mean by the term 'inflammation'? A contemporary basic science update for sports medicine. *Br J Sports Med* 2004;**38**(3):372–80.
- 2 Creaney L, Hamilton B. Growth factor delivery methods in the management of sports injuries: the state of play. *Br J Sports Med* 2008;**42**(5):314–20.
- 3 Knight KL. Cryotherapy in sports injury management. Champaign, IL: Human Kinetics; 1995. p. 3–98.
- 4 Merrick MA. Secondary injury after musculoskeletal trauma: a review and update. *J Athl Train* 2002;**37**(2):209–17.
- 5 Osterman AL, Heppenstall RB, Sapega AA, Katz M, Chance B, Sokolow D. Muscle ischaemia and hypothermia: a bioenergetic study using ³¹Phosphorous Nuclear Magnetic Resonance Spectroscopy. *J Trauma* 1984;**24**(9):811–7.
- 6 Sapega AA, Heppenstall RB, Sokolow DP, Graham TJ, Maris JM, Ghosh AK, *et al.* The bioenergetics of preservation of limbs before replantation. The rationale for intermediate hypothermia. *J Bone Joint Surg Am* 1988;**70**(10):1500–13.
- 7 Merrick MA, Rankin JM, Andres FA, Hinman CL. A preliminary examination of cryotherapy and secondary injury in skeletal muscle. *Med Sci Sports Exerc* 1999;**31**:1516–21.
- 8 Tidball JG. Inflammatory processes in muscle injury and repair. *Am J Physiol Regul Integr Comp Physiol* 2005;**288**(2):R345–53.
- 9 Toumi H, Best TM. The inflammatory response: friend or foe for muscle injury? *Br J Sports Med* 2003;**37**(4):284–6.
- 10 Blake DR, Allen RE, Lunec J. Free radicals in biological systems – a review orientated to inflammatory processes. *Br Med Bull* 1987;**43**(2):371–85.
- 11 Martinez-Cayuela M. Oxygen free radicals and human disease. *Biochimie* 1995;**77**(3):147–61.
- 12 Brickson S, Ji LL, Schell K, Olabisi R, St Pierre Schneider B, Best TM. M1/70 attenuates blood-borne neutrophil oxidants, activation and myofiber damage following stretch injury. *J Appl Physiol* 2003;**95**(3):969–76.
- 13 Westermann S, Vollmar B, Thorlacius H, Menger MD. Surface cooling inhibits tumor necrosis factor- α -induced microvascular perfusion failure, leukocyte adhesion, and apoptosis in the striated muscle. *Surgery* 1999;**126**:881–9.
- 14 Lee H, Natsui H, Akimoto T, Yanagi K, Ohshima N, Kono I. Effects of cryotherapy after contusion using real-time intravital microscopy. *Med Sci Sports Exerc* 2005;**37**(7):1093–8.
- 15 Schaser KD, Stover JF, Melcher I, Lauffer A, Haas NP, Bail HJ, *et al.* Local cooling restores microcirculatory hemodynamics after closed soft-tissue trauma in rats. *J Trauma* 2006;**61**:642–9.
- 16 Schaser KD, Disch AC, Stover JF, Lauffer A, Bail HJ, Mittlmeier T. Prolonged superficial local cryotherapy attenuates microcirculatory impairment, regional inflammation, and muscle necrosis after closed soft tissue injury in rats. *Am J Sports Med* 2007;**35**(1):93–102.
- 17 Farry PJ, Prentice NG, Hunter AC, Wakelin CA. Ice treatment of injured ligaments: an experimental model. *N Z Med J* 1980;**91**(651):12–4.
- 18 Hurme T, Rantanen J, Kalimo H. Effects of early cryotherapy in experimental skeletal muscle injury. *Scand Med Sci Sports* 1993;**3**:46–51.

- 19 Brickson S, Hollander J, Corr DT, Ji LL, Best TM. Oxidant production and immune response after stretch injury in skeletal muscle. *Med Sci Sports Exerc* 2001;**33**(12):2010–5.
- 20 Yuan J, Murrell GA, Wei AQ, Wang MX. Apoptosis in rotator cuff tendonopathy. *J Orthop Res* 2002;**20**(6):1372–9.
- 21 Yuan J, Murrell GA, Trickett A, Wang MX. Involvement of cytochrome c release and caspase-3 activation in the oxidative stress-induced apoptosis in human tendon fibroblasts. *Biochim Biophys Acta* 2003;**1641**(1):35–41.
- 22 Smith TL, Curl WW, Paterson Smith B, Holden MB, Wise T, Marr A, *et al.* New skeletal muscle model for the longitudinal study of alterations in microcirculation following contusion and cryotherapy. *Microsurgery* 1993;**14**:487–93.
- 23 Weston M, Taber C, Casagrande L, Cornwall M. Changes in local blood volume during cold gel pack application to traumatized ankles. *J Orthop Sports Phys Ther* 1994;**19**(4):197–9.
- 24 Karunakara RG, Lephart SM, Pincivero DM. Changes in forearm blood flow during single and intermittent cold application. *J Orthop Sports Phys Ther* 1999;**29**(3):177–80.
- 25 Knobloch K, Grasemann R, Spies M, Vogt PM. Midportion Achilles tendon microcirculation after intermittent combined cryotherapy and compression with cryotherapy alone: a randomized trial. *Am J Sports Med* 2008;**36**(11):2128–38.
- 26 Knobloch K, Kraemer R, Lichtenberg A, Jagodzinski M, Gosling T, Richter M, *et al.* Microcirculation of the ankle after Cryo/Cuff application in healthy volunteers. *Int J Sports Med* 2006;**27**(3):250–5.
- 27 Knobloch K, Grasemann R, Jagodzinski M, Richter M, Zeichen J, Krettek C. Changes of Achilles midportion tendon microcirculation after repetitive simultaneous cryotherapy and compression using a Cryo/Cuff. *Am J Sports Med* 2006;**34**(12):1953–9.
- 28 Knobloch K, Grasemann R, Spies M, Vogt PM. Intermittent KoldBlue cryotherapy of 3 × 10 min changes mid-portion Achilles tendon microcirculation. *Br J Sports Med* 2007;**41**(6):e4.
- 29 Curl WW, Smith BP, Marr A, Rosencrance E, Holden M, Smith TL. The effect of contusion and cryotherapy on skeletal muscle microcirculation. *J Sports Med Phys Fitness* 1997;**37**(4):279–86.
- 30 Dolan MG, Mychaskiw AM, Mattacola CG, Mendel FC. Effect of cool-water immersion and high voltage electric stimulation for 3 continuous hours on acute edema in rats. *J Athl Train* 2003;**38**(4):325–9.
- 31 Deal DN, Tipton J, Rosencrance E, Curl WW, Smith TL. Ice reduces edema. A study of microvascular permeability in rats. *J Bone Joint Surg Am* 2002;**84-A**(9):1573–8.
- 32 Garrett WE Jr, Safran MR, Seaber AV, Glisson RR, Ribbeck BM. Biomechanical comparison of stimulated and nonstimulated skeletal muscle pulled to failure. *Am J Sports Med* 1987;**15**:448–54.
- 33 Crisco JJ, Hentel KD, Jackson WO, Goehner K, Jokl P. Maximal contraction lessens impact response in a muscle contusion model. *J Biomech* 1996;**29**:1291–6.