Current evidence linking nutrition with brain health in ageing

Moore, K., O'Shea, M., Hughes, C., Hoey, L., Ward, M., & McNulty, H. (2017). Current evidence linking nutrition with brain health in ageing. *Nutrition Bulletin*, 42(1), 61-68. <https://doi.org/10.1111/nbu.12250>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Nutrition Bulletin

Publication Status:
Published (in print/issue): 09/02/2017

DOI:
[10.1111/nbu.12250](https://doi.org/10.1111/nbu.12250)

Document Version
Author Accepted version

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:

This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:

1. Replace (Ins) Tool – for replacing text.

Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.

standard framework for the analysis of microeconomic behavior. Nevertheless, it also led to the development of a new class of strategic form games. The number of competitors in the industry is that the structure of the game is a main component. At the micro level, are exogenous variables and important works on entry by firms (M. Henceforth) we open the 'black b

2. Strikethrough (Del) Tool – for deleting text.

Strikes a red line through text that is to be deleted.

How to use it

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.

there is no room for extra profits as mark-ups are zero and the number of firms (set) values are not determined by Blanchard and Kiyotaki (1987), perfect competition in general equilibrium of aggregate demand and supply in the classical framework assuming monopoly between an exogenous number of firms

3. Add note to text Tool – for highlighting a section to be changed to bold or italic.

Highlights text in yellow and opens up a text box where comments can be entered.

How to use it

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark-ups consistent with the VAR evidence

sation by Markov and Bell on the number of competitors and the impact on the demand-

4. Add sticky note Tool – for making notes at specific points in the text.

Marks a point in the proof where a comment needs to be highlighted.

How to use it

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.

standard and supply shocks. Most of the number of competitors and the impact on the demand-

5. Attach File Tool – for inserting large amounts of text or replacement figures.

Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it

- Click on the [Attach File](#) icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.

END

6. Drawing Markups Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks.

How to use it

- Click on one of the shapes in the Drawing Markups section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.

Current evidence linking nutrition with brain health in ageing

1 K. Moore, M. O'Shea and H. McNulty

2 Northern Ireland Centre for Food and Health, Ulster University Coleraine, Coleraine, UK

Abstract

Cognitive dysfunction and depression are significant problems of ageing with major health and socio-economic impacts; therefore, preventing or delaying the onset of these disorders should be a public health priority. In particular, there is a need to identify modifiable factors that could be targeted to promote better brain health in ageing. Epidemiological studies indicate a protective role for certain dietary patterns, in particular the Mediterranean diet, and for specific nutrients, including *n*-3 polyunsaturated fatty acids, polyphenols and B vitamins. However, the evidence to date from randomised controlled trials is generally inconsistent, although there is clearer evidence to support a role for folate and related B vitamins in slowing the progression of cognitive decline and possibly reducing the risk of depression in ageing. Future studies incorporating new technologies offer much promise for the development of effective nutrition strategies that could reduce the risk of cognitive and mental disorders and improve quality of life in our ageing population.

Keywords: ageing, B vitamins, cognitive function, depression, fatty acids, nutrition

Introduction

The global population is ageing with the number of those aged 60 years and over predicted to reach two billion by 2050 (United Nations Department of Economic and Social Affairs/Population Division 2009). An estimated 46.8 million people are living with dementia worldwide, and this figure is predicted to double in the next 20 years, with associated costs to the global economy currently estimated at over \$818 billion/year (Prince *et al.* 2015). Depression is a leading cause of disability, currently costing approximately £7.5 billion annually in England alone, and projected to increase by 67% by 2026 (National Collaborating Centre for Mental Health 2010). Given the significant impact of these conditions, there is a need to identify

modifiable factors that could be targeted to promote better brain health in ageing populations. Epidemiological evidence supports a role for certain dietary factors in brain health, opening up new potential avenues for prevention of dementia and mental illness in ageing (Panza *et al.* 2008; Rechenberg 2016).

The ageing brain

Ageing affects the brain from a cellular to a functional level. The brain has a high metabolic rate; therefore, oxidative stress and inflammation are common in ageing neural tissue (Bishop *et al.* 2010). On average, brain size decreases with age in later life and although there is no loss of cortical volume, there is white matter atrophy (Silverman *et al.* 1997). The rate of this atrophy is known to be a marker of cognitive decline (Fox *et al.* 1999) and white matter lesions in ageing are linked with late onset depression (Wang *et al.* 2014). Neurodegeneration is the loss of neuronal

Correspondence: Prof. Helene McNulty, Northern Ireland Centre for Food and Health, Ulster University Coleraine, Coleraine BT52 1SA, UK.

3 E-mail: h.mculty@ulster.ac.uk

processes and nerve cells. This can lead to neurodegenerative disorders which are characterised by selective loss of neurons in the motor, sensory and cognitive systems. This affects memory, cognition, language, personality and mood and can have sporadic and/or hereditary origins (Gelb 2016).

Mild cognitive impairment can be described as a rate of cognitive decline greater than expected for an individual's age and education level but that does not interfere notably with activities of daily life (Gauthier *et al.* 2006). Of note, however, is that an estimated 50% of those with mild cognitive impairment will go on to develop dementia within 5 years (Gauthier *et al.* 2006). Dementia is characterised by progressive deterioration of multiple cognitive domains leading to impaired daily functioning, with Alzheimer's disease being the most common cause of dementia in older adults (Prince *et al.* 2014). The World Health Organisation (2014) defines mental health as a state of 'complete physical, mental and social wellbeing and not merely the absence of disease or infirmity'; mental illness includes depression and anxiety as common disorders. A number of modifiable risk factors for declining cognitive and mental health have been identified, including dietary patterns and specific nutrients (Smith 2008; Stahl *et al.* 2014; Gallagher *et al.* 2016; Kennedy 2016; Rechenberg 2016). Ageing itself is associated with a reduction in the quantity of food consumed, altered metabolism and decreased absorption of nutrients, all of which can increase the risk of malnutrition (Wakimoto & Block 2001). Therefore, nutritional approaches to prevent or slow the progression of cognitive decline are of much current research interest.

Foods and brain health

Increasing evidence implicates certain dietary patterns, such as high intake of fruit and vegetables (Kang *et al.* 2005) and fish consumption (Barberger-Gateau *et al.* 2007), as being beneficial to brain health. Compliance with a Mediterranean diet (typically characterised by higher intakes of fruit, vegetables, wholegrains, olive oil and fish) has been associated with lower risk of Alzheimer's disease and slower progression of symptoms from mild cognitive impairment to Alzheimer's disease (Scarmeas *et al.* 2006; Solfrizzi *et al.* 2010; Singh *et al.* 2014). Adherence to a Mediterranean diet also has been associated with a protective effect against depression (Psaltopoulou *et al.* 2013). These findings were recently confirmed in a study of 4470 participants, aged from 45 to 79 years, which found

significantly lower depression scores in those with greater adherence to a Mediterranean diet (Veronese *et al.* 2016). Although there is some biological basis for the protective role of the Mediterranean diet in brain health, the findings do not prove causality as the studies to date have predominantly been observational.

Specific nutrients and brain health

Fatty acids

The fatty acid composition of brain membrane is directly affected by the diet, and this has focused attention on the role of dietary fatty acids in brain health. There is evidence that long-chain *n*-3 polyunsaturated fatty acids, eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) have potential benefits in cognitive and mental health. The suggested mechanism for the effects of these fatty acids is via their antithrombotic and anti-inflammatory properties (Gillette-Guyonnet *et al.* 2013) and their potential interactions with neurotransmitters via phospholipid metabolism and regulation of signal transduction (Grosso *et al.* 2014a). Regular consumption of dietary *n*-3 polyunsaturated fatty acids has been associated with reduced risk of cognitive decline and reduced development of incident dementia in epidemiological studies (Jicha & Markesbery 2010). Likewise, higher dietary DHA has been associated with lower incidence of dementia (Schaefer *et al.* 2006; Lopez *et al.* 2011). Randomised controlled trials (RCTs) of *n*-3 polyunsaturated fatty acid supplementation have, however, shown somewhat inconsistent results with regard to cognitive function (Freund-Levi *et al.* 2006; Quinn *et al.* 2010; Yurko-Mauro *et al.* 2010). Furthermore, the relationship between fatty acids and cognitive outcomes is complex since higher intakes of *n*-6 fatty acids, irrespective of *n*-3 polyunsaturated fatty acid intake, seem to negate any of the aforementioned beneficial effects of *n*-3 polyunsaturated fatty acids (Jicha & Markesbery 2010). One meta-analysis of RCTs intervening with supplemental *n*-3 polyunsaturated fatty acids concluded that the beneficial effects on brain health may be confined only to patients with mild cognitive impairment (Mazereeuw *et al.* 2012); however, this interpretation requires further investigation.

In two recent meta-analysis of RCTs, EPA supplementation was found to benefit patients with a diagnosis of depression; however, in those with non-clinical depression, the findings were far less clear

(Grosso *et al.* 2014b; Hallahan *et al.* 2016). A recent Cochrane review concluded that there was insufficient evidence to support increased *n*-3 polyunsaturated fatty acid intake as a treatment for depression, highlighting the need for further investigation in this area (Appleton *et al.* 2015).

Protein and carbohydrates

The role of dietary protein in cognitive function and mental health has not been extensively studied in ageing populations. Lower verbal memory scores were, however, observed in older people with lower dietary protein intakes (Goodwin *et al.* 1983). Additionally higher dietary protein intake was found to be positively correlated with non-verbal learning and verbal memory (Koehler *et al.* 1997) and reduced risk of mild cognitive impairment or dementia (Roberts *et al.* 2012). One RCT investigating the effects of dietary protein from red meat on cognitive function in older adults is ongoing (Daly *et al.* 2015).

The association between carbohydrates and cognitive function is unclear because available evidence is scarce (Power *et al.* 2015), with a Cochrane review identifying only one RCT in older adults (Ooi *et al.* 2011). While more research has focused on carbohydrates and depression, the available evidence is somewhat conflicting. One study of community-dwelling older adults found that those with depressive symptoms had a diet with a higher glycaemic index (GI) and glycaemic load (GL) (Mwamburi *et al.* 2011). A prospective investigation also reported that a high GI diet was associated with an increased risk of depression (Gangwisch *et al.* 2015). Contrary to these findings, research in institutionalised older adults reported that those suffering from depression had diets with a lower GL (Aparicio *et al.* 2013). Given the inconsistencies within the evidence base in this area, there is clearly a need for further well-designed studies.

Polyphenols and vitamins

Inflammation is thought to be involved in the neurodegenerative cascade resulting in cognitive decline (Gorelick 2010), and evidence suggests a protective anti-inflammatory role for flavonoids against cognitive decline and Alzheimer's disease (Commenges *et al.* 2000; Letenneur *et al.* 2007; Schaffer *et al.* 2012). In one large prospective study of 5395 older adults in the Netherlands, polyphenol intake was associated with a reduced risk of dementia in smokers (Engelhart *et al.* 2002). Likewise, dietary polyphenol intake was also

associated with cognitive performance in 2574 middle-aged French participants (Kesse-Guyot *et al.* 2012). Of greater note, new evidence from a recent RCT of healthy older adults reported that intervention with high doses of cocoa-flavanol enhanced dentate gyrus function, a brain area essential for learning and memory, which was measured by magnetic resonance imaging (MRI) and cognitive tests (Brickman *et al.* 2014). In the case of depression, animal studies have shown anti-depressant effects of numerous polyphenols, postulated to be related to their anti-inflammatory and neurotransmitter modulation roles (Pathak *et al.* 2013). Although positive effects of cocoa polyphenols on mood were reported in one small study of 87 middle-aged participants (Pase *et al.* 2013), the evidence from human studies is generally scarce and expert groups have called for further co-ordinated research in this area (Ward & Pasinetti 2016).

Oxidative stress is thought to be a major contributor to neurodegeneration and depression; thus, antioxidants have received particular attention in the study of older adults. Although large prospective studies have failed to demonstrate a protective effect of either dietary or supplemental ascorbic acid on cognitive function in older adults (Luchsinger *et al.* 2003; Laurin *et al.* 2004; Devore *et al.* 2010), the studies had significant limitations including the lack of a biomarker measure of vitamin C status. With regard to mental health, an early RCT with an antioxidant supplement containing vitamins A, E and C in 205 participants reported substantial increases in plasma ascorbic acid concentrations and correspondingly better mood outcomes after one year (Smith *et al.* 1999). Likewise, more recent cross-sectional and prospective studies have reported lower dietary intakes or plasma concentrations of ascorbic acid among older adults with depression (Merrill *et al.* 2008; Hamer *et al.* 2011; Payne *et al.* 2012). More robust RCTs, which include both biomarker and dietary intake measures, are required to further explore the role of vitamin C in mental health.

There is extensive literature in the area of vitamin D and the ageing brain. Of note, a 7-year follow-up study of 498 women aged 75 years and older identified significantly lower risk of Alzheimer's disease for those participants in the highest quintile of dietary vitamin D intake at baseline (Annweiler *et al.* 2012). In a recent meta-analysis of observational studies, patients suffering from Alzheimer's disease were found to have significantly lower serum vitamin D concentrations compared with matched cognitively healthy

controls (Annweiler *et al.* 2013). However, combined supplementation with vitamin D and calcium was not found to decrease the risk of dementia or cognitive decline in a RCT involving 4143 older women from the US (Rossom *et al.* 2012). In the case of depression, large cross-sectional ($n = 2598$) and prospective ($n = 2839$) studies have linked lower serum vitamin D status with an increased risk of depression (Williams *et al.* 2014; Brouwer-Brolsma *et al.* 2016). The available evidence from intervention studies, however, has not confirmed a causative relationship for vitamin D in relation to the risk of depression (Li *et al.* 2014; van den Berg *et al.* 2016).

B vitamins and cognition in ageing

The role of folate and related B vitamins in brain health in ageing, and related mechanisms involving one-carbon metabolism, is an area of active research worldwide. Within one-carbon metabolism, folate, along with vitamins B₁₂ and B₆, and riboflavin are cofactors in DNA synthesis and repair, amino acid metabolism and methylation reactions, including the remethylation of homocysteine to methionine and subsequent generation of S-adenosylmethionine. The latter metabolite is essential for neurotransmitter synthesis and thus vital for brain function and activity (McGarel *et al.* 2015).

A number of observational studies have shown that lower status of folate, vitamin B₁₂ and vitamin B₆ (and/or higher concentrations of homocysteine) are associated with cognitive deficit in ageing (Smith & Refsum 2016). RCTs have addressed the hypothesis that optimising B vitamin status by supplementing with folate alone, or in combination with vitamins B₁₂ and/or B₆, will delay cognitive decline in older adults. Available RCTs in older adults that include intervention with high-dose folic acid, vitamin B₁₂ and vitamin B₆ over 2 years or more have shown not only improved cognitive performance (Durga *et al.* 2007; Smith *et al.* 2010; de Jager *et al.* 2012; Douaud *et al.* 2013) but, in those incorporating MRI, also a reduced rate of brain atrophy (Durga *et al.* 2007; Smith *et al.* 2010; de Jager *et al.* 2012; Douaud *et al.* 2013). The RCT evidence is not entirely conclusive, however, with one notable trial from New Zealand failing to observe a significant beneficial effect of folic acid, vitamin B₁₂ and vitamin B₆ on cognition (McMahon *et al.* 2006), possibly owing to the inclusion of participants with generally high baseline folate status and thus unlikely to benefit from intervention aimed at improving status. One recent and rather controversial meta-

analysis in this area concluded that neither folic acid nor vitamin B₁₂ had a beneficial effect on cognition in older adults (Clarke *et al.* 2014). The article was widely criticised at the time of publication, mainly as a result of the inclusion criteria used to select the trials for investigation, and thus, the findings are not generally accepted by the scientific community in this area (Garrard & Jacoby 2015; Smith *et al.* 2015). It is clear that further well-designed RCTs are needed, especially those targeting participants with low B vitamin status as they are likely to benefit the most from increasing B vitamin concentrations to achieve better cognitive health in ageing.

The role of B vitamins in depressive disorders has not received as much interest as in cognitive function. However, one meta-analysis of 19 observational studies reported a significant relationship between low folate status and risk of depression (Gilbody *et al.* 2007). Low dietary intake and status of vitamin B₁₂ have also been linked with an increased risk of depression (Reynolds 2006; Kim *et al.* 2008; Ng *et al.* 2009; Sánchez-Villegas *et al.* 2009; Robinson *et al.* 2011; Moorthy *et al.* 2012). RCTs of B vitamin supplementation as an adjunct to anti-depressant medications (Coppens & Bailey 2000; Almeida *et al.* 2014) or alone (Almeida *et al.* 2010; Walker *et al.* 2010) have provided mixed results and no clear conclusions, partly because of major methodological differences among studies. Reviews of the available evidence linking B vitamins with depression have concluded that folate and vitamin B₁₂ may have roles in the longer term management of this condition (Taylor *et al.* 2004; Almeida *et al.* 2015).

Use of novel technologies in nutrition and brain research

Validated questionnaire-based assessments are the most common means of investigating cognitive and mental health outcomes and indeed are most informative in regard to the effect of nutrition on behaviour (Macready *et al.* 2010). However, in order to study the role of nutrition in brain function, the emerging use of brain imaging techniques in recent years provides an objective and highly robust means of assessing brain function and response (de Jager & Kovatcheva 2010). A number of such techniques are available and have been reviewed in detail elsewhere (Sizonenko *et al.* 2013). Magnetic resonance imaging (MRI) is a structural imaging technique that provides detailed pictures of brain tissue (white and grey matter, blood vessels and bone) using magnetic fields

and radiofrequency pulses. Functional MRI (fMRI) uses blood oxygen level-dependant imaging to visualise changes in blood flow, in order to identify areas of neural activity within the brain. Electroencephalography (EEG) and magnetoencephalography (MEG) are two techniques for functional brain imaging that record electric and magnetic activities of the brain at the scalp and have the highest temporal resolution compared to other imaging techniques.

In recent years, these imaging techniques have been utilised in nutrition research. One notable study, referred to earlier in this review, effectively used MRI to confirm the beneficial effects of B vitamins on cognition in older adults over a 2-year intervention period (Smith *et al.* 2010). Another study used fMRI to demonstrate higher brain activation in specific regions of the brain in participants who consumed a nutritionally balanced breakfast (Akitsuki *et al.* 2011). EEG has also been used to investigate the effects of diet on brain function, with one recent report showing improved memory and functional connectivity in the delta band in response to Souvenaid®, a nutritional supplement, in mild Alzheimer-type patients (Ritchie *et al.* 2014). In recent years, MEG has been approved by the US Food and Drug Administration (FDA) and is being used for clinical and research purposes, including investigating cognitive dysfunction, Alzheimer's disease and depression (Maestú *et al.* 2008; Cheng *et al.* 2012; de Haan *et al.* 2012; Kurita *et al.* 2016). The application of these new technologies in the field of nutrition, in combination with clinical and questionnaire-based assessments, provides much potential for robust investigation in future studies, furthering knowledge and discovery, in an effort to reduce the burden of declining brain health in ageing.

Conclusions

Undoubtedly, nutrition has an important role in preserving cognitive and mental health and thus improving quality of life in older age. The impact of specific nutritional factors on brain health in ageing is an area of active investigation worldwide. Emerging evidence implicates subclinical deficiencies of certain nutrients in cognitive decline and poor mental health in older adults; however, the threshold for nutrient levels to prevent or delay declining brain health is still unknown. If the findings of studies described in this review, which show promise in relation to B vitamins, *n*-3 polyunsaturated fatty acids and polyphenols, are confirmed, a public health strategy to improve status of these key nutrients may help to achieve better

cognitive and mental health in ageing. Future studies incorporating imaging techniques offer a robust basis for confirming effective nutrition interventions that could reduce the risk of cognitive and mental decline in ageing and the related burden on health services.

Acknowledgements

The PhD studentship of K.M. was funded by the Northern Ireland Department for Employment and Learning.

Conflicts of interest

The authors declare no conflict of interest.

References

- Akitsuki Y, Nakawaga S, Sugiura M *et al.* (2011) Nutritional quality of breakfast affects cognitive function: an fMRI study. *Neuroscience and Medicine* 2: 192–7.
- Almeida OP, Marsh K, Alfonso H *et al.* (2010) B-vitamins reduce the long-term risk of depression after stroke: the VITATOPS-DEP trial. *Annals of Neurology* 68: 503–10.
- Almeida OP, Ford AH, Hirani V *et al.* (2014) B vitamins to enhance treatment response to antidepressants in middle-aged and older adults: results from the B-VITAGE randomised, double-blind, placebo-controlled trial. *British Journal of Psychiatry* 205: 450–7.
- Almeida OP, Ford AH & Flicker L (2015) Systematic review and meta-analysis of randomized placebo-controlled trials of folate and vitamin B12 for depression. *International Psychogeriatrics* 27: 727–37.
- Annweiler C, Rolland Y, Schott AM *et al.* (2012) Higher vitamin D dietary intake is associated with lower risk of Alzheimer's disease: a 7-year follow-up. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences* 67: 1205–11.
- Annweiler C, Llewellyn D & Beauchet O (2013) Low serum vitamin D concentrations in Alzheimer's disease: a systematic review and meta-analysis. *Journal of Alzheimer's Disease* 33: 659–74.
- Aparicio A, Robles F, López-Sobaler A *et al.* (2013) Dietary glycaemic load and odds of depression in a group of institutionalized elderly people without antidepressant treatment. *European Journal of Nutrition* 52: 1059–66.
- Appleton KM, Sallis HM, Perry R *et al.* (2015) Omega-3 fatty acids for depression in adults. *The Cochrane Database of Systematic Reviews* (11): CD004692.
- Barberger-Gateau P, Raffaitin C, Letenneur L *et al.* (2007) Dietary patterns and risk of dementia: the Three-City cohort study. *Neurology* 69: 1921–30.
- van den Berg KS, Marijnissen RM, van den Brink RHS *et al.* (2016) Vitamin D deficiency, depression course and mortality: longitudinal results from the Netherlands Study on Depression in Older persons (NESDO). *Journal of Psychosomatic Research* 83: 50–6.
- Bishop NA, Lu T & Yankner BA (2010) Neural mechanisms of ageing and cognitive decline. *Nature* 464: 529–35.

- Brickman AM, Khan UA, Provenzano FA *et al.* (2014) Enhancing dentate gyrus function with dietary flavanols improves cognition in older adults. *Nature Neuroscience* 17: 1798–803.
- Brouwer-Brolsma E, Dhonukshe-Rutten R, van Wijngaarden J *et al.* (2016) Low vitamin D status is associated with more depressive symptoms in Dutch older adults. *European Journal of Nutrition* 55: 1525–34.
- Cheng C, Wang P, Hsu W *et al.* (2012) Inadequate inhibition of redundant auditory inputs in Alzheimer's disease: an MEG study. *Biological Psychology* 89: 365–73.
- Clarke R, Bennett D, Parish S *et al.* (2014) Effects of homocysteine lowering with B vitamins on cognitive aging: meta-analysis of 11 trials with cognitive data on 22,000 individuals. *American Journal of Clinical Nutrition* 100: 657–66.
- Commenges D, Scotet V, Renaud S *et al.* (2000) Intake of flavonoids and risk of dementia. *European Journal of Epidemiology* 4: 357.
- Coppen A & Bailey J (2000) Enhancement of the antidepressant action of fluoxetine by folic acid: a randomised, placebo controlled trial. *Journal of Affective Disorders* 60: 121–30.
- Daly RM, Gianoudis J, Prosser M *et al.* (2015) The effects of a protein enriched diet with lean red meat combined with a multi-modal exercise program on muscle and cognitive health and function in older adults: study protocol for a randomised controlled trial. *Trials* 16: 1.
- Devore EE, Grodstein F, van Rooij F *et al.* (2010) Dietary antioxidants and long-term risk of dementia. *Archives of Neurology* 67: 819–25.
- Douaud G, Nichols TE, Smith SM *et al.* (2013) Preventing Alzheimer's disease-related gray matter atrophy by B-vitamin treatment. *Proceedings of the National Academy of Sciences of the United States of America* 110: 9523–8.
- Durga J, van Boxtel M, Schouten PJ *et al.* (2007) Articles: effect of 3-year folic acid supplementation on cognitive function in older adults in the FACIT trial: a randomised, double blind, controlled trial. *The Lancet* 369: 208–16.
- Engelhart MJ, Geerlings MI, Ruitenberg A *et al.* (2002) Dietary intake of antioxidants and risk of Alzheimer disease. *The Journal of the American Medical Association* 287: 3223–9.
- Fox NC, Schill RI, Crum WR *et al.* (1999) Correlation between rates of brain atrophy and cognitive decline in AD. *Neurology* 52: 1687–9.
- Freund-Levi Y, Eriksdotter-Jönghagen M, Cederholm T *et al.* (2006) Omega-3 fatty acid treatment in 174 patients with mild to moderate Alzheimer disease: OmegAD study: a randomized double-blind trial. *Archives of Neurology* 63: 1402–8.
- Gallagher D, Kiss A, Lancot K *et al.* (2016) Depressive symptoms and cognitive decline: a longitudinal analysis of potentially modifiable risk factors in community dwelling older adults. *Journal of Affective Disorders* 190: 235–40.
- Gangwisch J, Hale L, Garcia L *et al.* (2015) High glycemic index diet as a risk factor for depression: analyses from the Women's Health Initiative. *American Journal of Clinical Nutrition* 102: 454–63.
- Garrard P & Jacoby R (2015) B-vitamin trials meta-analysis: less than meets the eye. *The American Journal of Clinical Nutrition* 101: 414–5.
- Gauthier S, Reisberg B, Zaudig M *et al.* (2006) Mild cognitive impairment. *Lancet* 367: 1262–70.
- Gelb D (2016) *Introduction to Clinical Neurology*, 5th edn. Oxford University Press: New York.
- Gilbody S, Lightfoot T & Sheldon T (2007) Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity. *Journal of Epidemiology and Community Health* 7: 631.
- Gillette-Guyonnet S, Secher M & Vellas B (2013) Nutrition and neurodegeneration: epidemiological evidence and challenges for future research. *British Journal of Clinical Pharmacology* 75: 738–55.
- Goodwin JS, Goodwin JM & Garry PJ (1983) Association between nutritional status and cognitive functioning in a healthy elderly population. *Journal of the American Medical Association* 249: 2917–21.
- Gorelick PB (2010) Role of inflammation in cognitive impairment: results of observational epidemiological studies and clinical trials. *Innate Inflammation and Stroke* 1207: 155–62.
- Grosso G, Galvano F, Marventano S *et al.* (2014a) Omega-3 fatty acids and depression: scientific evidence and biological mechanisms. *Oxidative Medicine and Cellular Longevity* 2014: 313570.
- Grosso G, Pajak A, Marventano S *et al.* (2014b) Role of omega-3 fatty acids in the treatment of depressive disorders: a comprehensive meta-analysis of randomized clinical trials. *PLoS One* 9: e96905.
- de Haan W, van ~~DE~~ *DF*, van Koene T *et al.* (2012) Disrupted modular brain dynamics reflect cognitive dysfunction in Alzheimer's disease. *NeuroImage* 59: 3085–93.
- Hallahan B, Ryan T, Davis JM *et al.* (2016) Efficacy of omega-3 highly unsaturated fatty acids in the treatment of depression. *British Journal of Psychiatry* 209: 192–201.
- Hamer M, Bates CJ & Mishra GD (2011) Depression, physical function, and risk of mortality: national diet and nutrition survey in adults older than 65 years. *American Journal of Geriatric Psychiatry* 19: 72–8.
- de Jager CA & Kovatcheva A (2010) Summary and discussion: methodologies to assess long-term effects of nutrition on brain function. *Nutrition Reviews* 68 (Suppl 1): S53–8.
- de Jager C, Oulhaj A, Jacoby R *et al.* (2012) Cognitive and clinical outcomes of homocysteine-lowering B-vitamin treatment in mild cognitive impairment: a randomized controlled trial. *International Journal of Geriatric Psychiatry* 27: 592–600.
- Jicha GA & Markesbery WR (2010) Omega-3 fatty acids: potential role in the management of early Alzheimer's disease. *Clinical Interventions in Aging* 5: 45–61.
- Kang JH, Ascherio A & Grodstein F (2005) Fruit and vegetable consumption and cognitive decline in aging women. *Annals of Neurology* 57: 713–20.
- Kennedy DO (2016) B vitamins and the brain: mechanisms, dose and efficacy—a review. *Nutrients* 8: 2.
- Kesse-Guyot E, Fezeu L, Andreeva VA *et al.* (2012) Total and specific polyphenol intakes in midlife are associated with cognitive function measured 13 years later. *The Journal of Nutrition* 142: 76–83.
- Kim JM, Stewart R, Kim SW *et al.* (2008) Predictive value of folate, vitamin B-12 and homocysteine levels in late-life depression. *British Journal of Psychiatry* 192: 268–74.
- Koehler KM, La Rue A, Wayne SJ *et al.* (1997) Nutritional status and cognitive functioning in a normally aging sample: a 6-y reassessment. *American Journal of Clinical Nutrition* 65: 20–9.

- Kurita S, Takei Y, Maki Y *et al.* (2016) Magnetoencephalography study of the effect of attention modulation on somatosensory processing in patients with major depressive disorder. *Psychiatry and Clinical Neurosciences* 70: 116–25.
- Laurin D, Masaki KH, Foley DJ *et al.* (2004) Midlife dietary intake of antioxidants and risk of late-life incident dementia: the Honolulu-Asia Aging Study. *American Journal of Epidemiology* 159: 959–67.
- Letenneur L, Proust-Lima C, Le Gouge A *et al.* (2007) Flavonoid intake and cognitive decline over a 10-year period. *American Journal of Epidemiology* 165: 1364–71.
- Li G, Mbuagbaw L, Samaan Z *et al.* (2014) Efficacy of vitamin d supplementation in depression in adults: a systematic review. *Journal of Clinical Endocrinology & Metabolism* 99: 757–67.
- Lopez LB, Kritiz-Silverstein D & Barrett-Connor E (2011) High dietary and plasma levels of the omega-3 fatty acid docosahexaenoic acid are associated with decreased dementia risk: the Rancho Bernardo study. *The Journal of Nutrition, Health & Aging* 15: 25–31.
- Luchsinger JA, Tang M, Shea S *et al.* (2003) Antioxidant vitamin intake and risk of Alzheimer disease. *Archives of Neurology* 60: 203–8.
- Macready AL, Butler LT, Kennedy OB *et al.* (2010) Cognitive tests used in chronic adult human randomised controlled trial micronutrient and phytochemical intervention studies. *Nutrition Research Reviews* 23: 200–29.
- Maestú F, Campo P, Del Río D *et al.* (2008) Increased biomagnetic activity in the ventral pathway in mild cognitive impairment. *Clinical Neurophysiology* 119: 1320–7.
- Mazereeuw G, Lancot KL, Chau SA *et al.* (2012) Effects of omega-3 fatty acids on cognitive performance: a meta-analysis. *Neurobiology of Aging* 33: 1482.e17–1482.e29.
- McGarel C, Pentieva K, Strain JJ *et al.* (2015) Emerging roles for folate and related B-vitamins in brain health across the lifecycle. *The Proceedings of the Nutrition Society* 74: 46–55.
- McMahon J, Green T, Skeaff C *et al.* (2006) A controlled trial of homocysteine lowering and cognitive performance. *New England Journal of Medicine* 354: 2764–72.
- Merrill RM, Taylor P, Aldana SG (2008) Coronary Health Improvement Project (CHIP) is associated with improved nutrient intake and decreased depression. *Nutrition (Burbank, Los Angeles County, Calif.)* 24: 314–21.
- Moorthy D, Peter I, Scott TM *et al.* (2012) Status of vitamins B-12 and B-6 but not of folate, homocysteine, and the methylenetetrahydrofolate reductase C677T polymorphism are associated with impaired cognition and depression in adults. *The Journal of Nutrition* 142: 1554–60.
- Mwamburi DM, Liebson E, Folstein M *et al.* (2011) Depression and glycemic intake in the homebound elderly. *Journal of Affective Disorders* 132: 94–8.
- National Collaborating Centre for Mental Health (2010) *The Treatment and Management of Depression in Adults (Updated Edition)*, National Clinical Practice Guideline 90. The British Psychological Society and The Royal College of Psychiatrists: Leicester.
- Ng TP, Feng L, Niti M *et al.* (2009) Folate, vitamin B12, homocysteine, and depressive symptoms in a population sample of older Chinese adults. *Journal of the American Geriatrics Society* 57: 871–6.
- Ooi CP, Loke SC, Yassin Z *et al.* (2011) Carbohydrates for improving the cognitive performance of independent-living older adults with normal cognition or mild cognitive impairment. *Cochrane Database of Systematic Reviews* (Online) (4): CD007220.
- Panza F, Solfrizzi V, Capurso C *et al.* (2008) Lifestyle-related factors in predementia and dementia syndromes. *Expert Review of Neurotherapeutics* 8: 133–58.
- Pase MP, Scholey AB, Pipingas A *et al.* (2013) Cocoa polyphenols enhance positive mood states but not cognitive performance: a randomized, placebo-controlled trial. *Journal of Psychopharmacology* 27: 451–8.
- Pathak L, Agrawal Y & Dhir A (2013) Natural polyphenols in the management of major depression. *Expert Opinion on Investigational Drugs* 22: 863–80.
- Payne ME, Steck SE, George RR *et al.* (2012) Fruit, vegetable, and antioxidant intakes are lower in older adults with depression. *Journal of the Academy of Nutrition and Dietetics* 112: 2022–7.
- Power SE, O'Connor EM, Ross RP *et al.* (2015) Dietary glycaemic load associated with cognitive performance in elderly subjects. *European Journal of Nutrition* 54: 557–68.
- Prince M, Knapp M, Guerchet M *et al.* (2014) *Dementia UK: Second Edition - Overview*. Alzheimer's Society: London.
- Prince M, Wimo M, Guerchet M *et al.* (2015) *World Alzheimer Report 2015. The Global Impact of Dementia. An Analysis of Prevalence, Incidence, Cost and Trends*. Alzheimer's Disease International (ADI): London.
- Psaltopoulou T, Sergentanis TN, Panagiotakos DB *et al.* (2013) Mediterranean diet, stroke, cognitive impairment, and depression: a meta-analysis. *Annals of Neurology* 74: 580–91.
- Quinn JF, Raman R, Thomas RG *et al.* (2010) Docosahexaenoic acid supplementation and cognitive decline in Alzheimer disease: a randomized trial.
- Rechenberg K (2016) Nutritional interventions in clinical depression. *Clinical Psychological Science* 4: 144–62.
- Reynolds E (2006) Vitamin B12, folic acid, and the nervous system. *The Lancet. Neurology* 5: 949–60.
- Ritchie CW, Bajwa J, Coleman G *et al.* (2014) Souvenaid®: a new approach to management of early Alzheimer's disease. *The Journal of Nutrition, Health & Aging* 18: 291–9.
- Roberts R, Roberts L, Geda Y *et al.* (2012) Relative intake of macronutrients impacts risk of mild cognitive impairment or dementia. *Journal of Alzheimer's Disease* 32: 329–39.
- Robinson DJ, O'Lunaigh C, Tehee E *et al.* (2011) Associations between holotranscobalamin, vitamin B12, homocysteine and depressive symptoms in community-dwelling elders. *International Journal of Geriatric Psychiatry* 26: 307–13.
- Rossum RC, Margolis KL, Espeland MA *et al.* (2012) Calcium and vitamin D supplementation and cognitive impairment in the Women's health initiative. *Journal of the American Geriatrics Society* 60: 2197–205.
- Sánchez-Villegas A, Doreste J, Schlatter J *et al.* (2009) Association between folate, vitamin B6 and vitamin B12 intake and depression in the SUN cohort study. *Journal of Human Nutrition & Dietetics* 22: 122–33.
- Scarmeas N, Stern Y, Tang M *et al.* (2006) Mediterranean diet and risk for Alzheimer's disease. *Annals of Neurology* 59: 912–21.
- Schaefer EJ, Bongard V, Beiser AS *et al.* (2006) Plasma phosphatidylcholine docosahexaenoic acid content and risk of

- dementia and Alzheimer disease: the Framingham Heart Study. *Archives of Neurology* 63: 1545–50.
- Schaffer S, Asseburg H, Kuntz S *et al.* (2012) Effects of polyphenols on brain ageing and Alzheimer's disease: focus on mitochondria. *Molecular Neurobiology* 46: 161–78.
- Silverman W, Wisniewski HM, Bobinski M *et al.* (1997) Frequency of stages of Alzheimer-related lesions in different age categories. *Neurobiology of Aging* 18: 377–9.
- Singh B, Parsaik AK, Knopman DS *et al.* (2014) Association of Mediterranean diet with mild cognitive impairment and Alzheimer's disease: a systematic review and meta-analysis. *Journal of Alzheimer's Disease* 39: 271–82.
- Sizonenko SV, Babiloni C, de Bruin E *et al.* (2013) Brain imaging and human nutrition: which measures to use in intervention studies? *The British Journal of Nutrition* 110 (Suppl 1): S1–30.
- Smith D (2008) The Worldwide challenge of the dementias: a role for B vitamins and homocysteine. *Food and Nutrition Bulletin* 29: S143.
- Smith AD & Refsum H (2016) Homocysteine, B vitamins, and cognitive impairment. *Annual Review of Nutrition* 36: 211–39.
- Smith AP, Clark RE, Nutt DJ *et al.* (1999) Vitamin C, mood and cognitive functioning in the elderly. *Nutritional Neuroscience* 2: 249–56.
- Smith AD, de Jager CA, Whitbread P *et al.* (2010) Homocysteine-lowering by b vitamins slows the rate of accelerated brain atrophy in mild cognitive impairment: a randomized controlled trial. *PLoS One* 5: 1–10.
- Smith A, de Jager C, Refsum H *et al.* (2015) Homocysteine lowering, B vitamins, and cognitive aging. *American Journal of Clinical Nutrition* 101: 415–6.
- Solfrizzi V, Scafato E, Capurso C *et al.* (2010) Metabolic syndrome and the risk of vascular dementia: the Italian Longitudinal Study on Ageing. *Journal of Neurology, Neurosurgery and Psychiatry* 81: 433–40.
- Stahl ST, Albert SM, Dew MA *et al.* (2014) Coaching in healthy dietary practices in at-risk older adults: a case of indicated depression prevention. *The American Journal of Psychiatry* 171: 499–505.
- Taylor MJ, Carney SM, Goodwin GM *et al.* (2004) Folate for depressive disorders: systematic review and meta-analysis of randomized controlled trials. *Journal of Psychopharmacology (Oxford, England)*, 18: 251–6.
- United Nations Department of Economic and Social Affairs/Population Division (2009) *World Population Prospects The 2008 revision-Highlights*. United Nations: New York.
- Veronese N, Stubbs B, Noale M *et al.* (2016) Adherence to the Mediterranean diet is associated with better quality of life: data from the Osteoarthritis Initiative. *American Journal of Clinical Nutrition* 104: 1403–9.
- Wakimoto P, Block G (2001) Dietary intake, dietary patterns, and changes with age: an epidemiological perspective. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 56 Spec No 2: 65–80.
- Walker JG, Mackinnon AJ, Batterham P *et al.* (2010) Mental health literacy, folic acid and vitamin B12, and physical activity for the prevention of depression in older adults: randomised controlled trial. *The British Journal of Psychiatry: The Journal of Mental Science* 197: 45–54.
- Wang L, Leonards CO, Sterzer P *et al.* (2014) White matter lesions and depression: a systematic review and meta-analysis. *Journal of Psychiatric Research* 56: 56–64.
- Ward L & Pasinetti GM (2016) Recommendations for development of botanical polyphenols as “natural drugs” for promotion of resilience against stress-induced depression and cognitive impairment. *NeuroMolecular Medicine* 18: 487–95.
- Williams J, Sink K, Toozé J *et al.* (2014) Low 25-hydroxyvitamin d concentrations predict incident depression in well-functioning older adults: the health, aging, and body composition study. *Journals of Gerontology Series A-Biological Sciences and Medical Sciences* 70: 757–63.
- World Health Organisation (2014) *Mental health: a state of well-being*. World Health Organisation. Available at: http://www.who.int/features/factfiles/mental_health/en/
- Yurko-Mauro K, McCarthy D, Rom D *et al.* (2010) Beneficial effects of docosahexaenoic acid on cognition in age-related cognitive decline. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association* 6: 456–64.

Author Query Form

Journal: NBU
Article: 12250

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please confirm that given names (red) and surnames/family names (green) have been identified correctly.
2	AUTHOR: Please check that authors and their affiliations are correct.
3	AUTHOR: Please check and provide the corresponding author's job position.
4	AUTHOR: The term "polyunsaturated fatty acids" should not be abbreviated as per style sheet. Hence "PUFA" has been changed to "polyunsaturated fatty acids" in all occurrences. Kindly check and approve the edits made.
5	AUTHOR: Please provide the familyname for the author DF for reference Haan et al. (2012).
6	AUTHOR: Please provide if anything missing in the rereference "Quinn et al. 2010".
7	AUTHOR: Please provide accessed date, month and year for reference "World Health Organisation (2014)".