Entrapment of *Lactobacillus casei* ATCC393 in the viscous matrix of Pistacia terebinthus resin for functional myzithra cheese manufacture

Vasiliki, S., Terpou, A., Bosnea, L., Kanellaki, M., & Singh - Nee Nigam, P. (2018). Entrapment of *Lactobacillus casei* ATCC393 in the viscous matrix of Pistacia terebinthus resin for functional myzithra cheese manufacture. *LWT - Food Science and Technology*, 89, 441-448. <https://doi.org/10.1016/j.lwt.2017.11.015>

[Link to publication record in Ulster University Research Portal](#)

Published in:
LWT - Food Science and Technology

Publication Status:
Published (in print/issue): 01/03/2018

DOI:
[10.1016/j.lwt.2017.11.015](https://doi.org/10.1016/j.lwt.2017.11.015)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Accepted Manuscript

Entrapment of *L. casei* ATCC393 in the viscous matrix of *Pistacia terebinthus* resin for functional myzithra cheese manufacture

Vasiliki Schoina, Antonia Terpou, Loulouda Bosnea, Maria Kanellaki, Poonam S. Nigam

PII: S0023-6438(17)30835-6

DOI: [10.1016/j.lwt.2017.11.015](https://doi.org/10.1016/j.lwt.2017.11.015)

Reference: YFSTL 6644

To appear in: *LWT - Food Science and Technology*

Received Date: 19 June 2017

Revised Date: 6 November 2017

Accepted Date: 9 November 2017

Please cite this article as: Schoina, V., Terpou, A., Bosnea, L., Kanellaki, M., Nigam, P.S., Entrapment of *L. casei* ATCC393 in the viscous matrix of *Pistacia terebinthus* resin for functional myzithra cheese manufacture, *LWT - Food Science and Technology* (2017), doi: 10.1016/j.lwt.2017.11.015.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Entrapment of *L. casei* ATCC393 in the viscous matrix of *Pistacia*
terebinthus resin for functional myzithra cheese manufacture**

Vasiliki Schoina¹, Antonia Terpou^{1*}, Loulouda Bosnea^{1,2}, Maria Kanellaki¹, Poonam S.
Nigam³.

1. Food Biotechnology Group, Department of Chemistry, University of Patras, GR-26500, Patras, Greece.

2. National Agricultural Research Foundation, Dairy Research Institute, Katsikas, 45221 Ioannina, Greece.

3. School of Biomedical Sciences, University of Ulster, Cromore Rd., Coleraine BT62 1SA, Northern Ireland, United Kingdom.

* Corresponding author

Dr. Antonia Terpou

E-mail : aterpou@upatras.gr

Tel.: +30-2610997123

Abstract

Pissa Paphos, a natural mastic resin (*Pistacia terebinthus*) was evaluated as an encapsulating and matrix-forming material for the immobilisation of the probiotic bacterium *Lactobacillus casei* ATCC 393. The immobilized biocatalyst was added as an adjunct for the production of functional myzithra cheese. In total, four myzithra cheeses were manufactured: a. cheese with *L. casei* cells entrapped in a *P. terebinthus* matrix (Pissa Paphos) b. cheese with free *L. casei* cells and *P. terebinthus* resin, c. myzithra cheese with free *L. casei* cells without the resin and d. traditional myzithra cheese. *P. terebinthus* resin provided antimicrobial properties by suppressing the growth of fungi/yeasts in myzithra cheese during refrigerated storage. On the contrary, the presence of the resin did not affect the cell counts of the probiotic microorganism which maintained in high populations (above 9 log CFU g⁻¹) during hole storage period. Additionally, the viscous matrix of the resin seems to confer a protective effect on entrapped *L. casei* cells since higher populations were observed over free cells. Moreover, all myzithra cheeses with incorporated resin were characterized by an exceptional mastic gum aroma and pleasant coherent texture which indicates the product's high commercialization potential.

Keywords: *Pistacia terebinthus* resin; probiotics; encapsulation; terpenes; myzithra cheese.

Chemical compounds studied in this article: α -pinen (PubChem CID: 6654), β -pinen (PubChem CID: 14896), α -terpineol (PubChem CID: 442501), 4-terpineol (PubChem CID: 11230), eukalyptol (PubChem CID: 2758), terpinolene (PubChem CID: 11463), myrtenol (PubChem CID: 10582), pinocarveol (PubChem CID: 88297),

48 3-carene (PubChem CID: 26049), o-cymene (PubChem CID: 10703), verbenol
49 (PubChem CID: 61126).

50

1. Introduction

Resins are plant products that exude naturally (surface resins) or can be obtained by incision or infection (internal resins). They are insoluble in water but soluble in organic solvents (Dell & McComb, 1979). The most popular resins origin from Pistacia plants (*Pistacia lentiscus*, *Pistacia terebinthus*) which native to the Mediterranean region from Morocco and Portugal to Greece, Turkey and Syria (Rauf et al., 2017). Most of the plant parts including fruits, fruit fatty oil and resin were used as food and traditional medicine in the region since ancient times (Lardos, 2006).

Pissa Paphos, the mastic gum obtained from *Pistacia terebinthus* L. (*Anacardiaceae* family) tree, grows mainly on dry rock slopes and hill sides or in pine forests of Cyprus especially in Paphos and Limassol district. The tree's aromatic resin, called by the locals "*Paphos pissa*" and/ or "*pissa Pafitiki*" has a significant contribution to the local economy (Lardos, 2006). Over the years, different parts of *P. terebinthus* tree have been reported to provide several ethnopharmacological utilizations such as an antiseptic, diuretic, anti-inflammatory, antipyretic, antibacterial and antiviral agent, for wound treatments, eczema, burns and stomach-aches (Rauf et al., 2017; Topçu et al., 2007).

Lately, consumers awareness has focused on safe and high-quality food products, leading the food companies and related industries to implement novel methods for the production of functional foods (Bogue, Collins, & Troy, 2017). One major category of functional foods is probiotic dairy products. Probiotics are lactic acid bacteria that when presented as live microbial supplements can confer a beneficially affect to the host by improving its intestinal microbial balance (Fuller & Gibson, 1998). It has been established that a minimum level of probiotic lactic acid bacteria ($10^6 \sim 10^7$ CFU g⁻¹) contained viable in dairy products is necessary for

improving human health (Shori, 2015). Thus, scientists have recently targeted on the development of novel methods that will enhance probiotic viability in dairy products like cell immobilisation, microencapsulation, addition of prebiotics, drying (freeze-drying, spray-drying) etc. (Bosnea, Moschakis, Nigam, & Biliaderis, 2017). Techniques like immobilisation and microencapsulation of probiotics, even though represent a great challenge, have been established in order to improve the survival of probiotic bacteria giving promising results since they increased survival rates and stability of probiotics during fermentation processing and storage (Champagne, Ross, Saarela, Hansen, & Charalampopoulos, 2011; Terpou, Bekatorou, Kanellaki, Koutinas, & Nigam, 2017). Also, cell encapsulation in dairy fermentation is a rapidly expanding research area because of its attractive technical and economic advantages compared to the conventional free cell systems (Bosnea, Moschakis, et al., 2017; Morales & Ruiz, 2016).

In addition, synthetic chemical additives are used as preservatives (antimicrobials, antioxidants and anti-browning) to ensure the products self-life and safety or as flavor enforcements for improvement of products characteristics (Carocho, Barreiro, Morales, & Ferreira, 2014). However, many studies have confirmed that the excessive consumption of synthetic food additives is related with gastrointestinal, respiratory, dermatological and neurological adverse reactions and as a result consumers avoid the consumption of such products (Caleja et al., 2016; Carocho et al., 2014). Therefore, an alternative solution for enhancement of self-life and safety of food products is the use of natural additives. Mastic gum and its essential oils are very promising food additives since numerous recent studies have demonstrated its antimicrobial and flavoring effect (Aksoy, Duran, & Koksall, 2006; Daifas et al., 2004; Paraschos et al., 2011; Schoina et al., 2014). Moreover, the use of

mastic gum (*Pistacia lentiscus*) as immobilisation support has recently been proposed by Morkhade (2017) as a successful microencapsulating and matrix-forming material for sustained drug release..

Thus, the aim of the present study was to examine the capability of *Pistacia terebinthus* resin as a probiotic microencapsulation matrix and its use as an adjunct for functional myzithra cheese making. The main targets of the study were to investigate *Pistacia terebinthus* resin as a natural encapsulation matrix for the probiotic bacterial strain *Lactobacillus casei* ATCC 393 (Saxami et al., 2012) and the effects on probiotic cell viability in myzithra cheeses during 30 days of storage (4°C) and product's shelf-life and finally the influence on the aromatic profile of the produced myzithra cheeses.

2. Materials and methods

2.1 *Pistacia terebinthus* resin for probiotic cell encapsulation

The probiotic Gram⁺ *Lactobacillus* bacterial strain *Lactobacillus casei* ATCC 393 (DSMZ, Braunschweig, Germany) was used for the microencapsulation process. The probiotic *L. casei* ATCC 393 was selected according to its *in vitro* and *in vivo* studies of the microbial survival in GI tract, adhesion to the intestine and modulation of the intestinal microflora in rats (Saxami et al., 2012).

L. casei cells were grown at 37°C in de Man-Rogosa-Sharpe (MRS) liquid medium (LabM, UK) for 48-72 h. Wet biomass was harvested by centrifugation (Sigma 3K12, Bioblock Scientific, France) at 5000 rpm for 10 min. The cultivated wet biomass was introduced in MRS liquid medium along with small particles of sterile freeze-dried *Pistacia terebinthus* resin (pissa Paphos'), as described previously by Antonia Terpou et al. (2017). Freeze drying was important for immobilisation performance since the structure opens and creates holes where the probiotic LAB cells can be entrapped. The system was placed in an incubator at 37°C and agitated periodically for immobilisation to be achieved (approximately 48h). When immobilisation bioprocess was performed (glucose in the liquid culture was <1 g/L), the fermented liquid was decanted *Pistacia terebinthus* resin with immobilized probiotic cells was washed twice with sterile Ringer's ¼ solution targeting the removal of any free cells (Schoina et al., 2015). All media were autoclaved at 120°C at 1–1.5 atm for 15 min prior to use.

2.2 Verification of encapsulation of probiotic cells by scanning electron microscopy and microbiological analysis

Pieces of *Pistacia terebinthus* resin (pissa Paphos) in comparison with pieces of the encapsulated biocatalyst were coated with gold in a Balzers SCD 004 Sputter coater (Bal-Tec, Schalksmühle, Germany) for 2 min. The samples were examined in a JSM-6300 scanning electron microscope (JEOL, Tokyo, Japan), operated at an accelerating voltage of 20kV. Scanning electron micrographs were obtained in order to investigate probiotic cell encapsulation in *Pistacia terebinthus* matrix.

The counts of the entrapped *L. casei* cells into a specific amount of the resins' viscous matrix were determined. In particular, 1 g of encapsulated biocatalyst was added to 9 mL of ringer solution followed by shaking in a homogenizer for 210 s. Existing probiotic colonies of the encapsulated biocatalyst were identified by enumeration in MRS agar medium (37 °C, 48 - 72 h). To demonstrate the complete detachment of encapsulated *L. casei* cells from the resins matrix, the first homogenate solution is poured out and new ringers' solution is added. Subsequently, the mixture is shaken in a homogenizer for 210 s and the resulting liquid is tested until no growth of *L. casei* colonies is detected.

2.3 *Myzithra* cheese production

Myzithra is a traditional Greek whey cheese produced by heating cheese whey at 88–92 °C under continuous stirring for 40–45 min in order to obtain the protein remaining from cheese production (Litopoulou-Tzanetaki & Tzanetakis, 2011).

Sweet cheese whey (0.54% fat, 1.67% total protein, 5.35% lactose and pH 6.4) derived as an industrial by-product of Graviera hard cheese production (A.VI.GAL SA-Achaia milk industry) (Bozoudi et al., 2016) was used for *myzithra* cheese making as described previously by Litopoulou-Tzanetaki and Tzanetakis (2011) with small modifications. Specifically, cheese whey was gradually heated under continuous

stirring with a rate of 1,5 °C/ min up to 75°C and when small curd particles of whey proteins were formed, the temperature was increased to 90 °C with a rate of 1 °C/ min for 45 min in total. Acidification of whey to pH 5.2 was performed by the addition of 10% lactic acid before heating (Pappas & Voutsinas, 2009). A very thin layer of coagulum was formed on the surface of the whey after 45 min of heating at 90 °C and then stirring was reduced and finally stopped. The formed curd was transferred gradually by a perforated ladle into a sterile perforated fabric and hung from a pole in a ventilated room at room temperature (18~22 °C) for 3 h in order to drain.

The drained curd was divided equivalent particles and four types of myzithra cheeses were prepared: (C) control myzithra cheese; (F) myzithra cheese with free probiotic cells (ME) myzithra cheese with encapsulated probiotic cells and (MF) myzithra cheese with *Pistacia terebinthus* resin and free probiotic cells. In the case of myzithra cheeses produced by the adjunct probiotic culture, the curd was cooled after heating at 37°C and homogenously 1g of free *L. casei* cells (F) or 0.5g of encapsulated *L. casei* cells in *P. terebinthus* resin (ME) or 0.5g of *P. terebinthus* resin with 1g of free *L. casei* cells (MF) were added respectively per 100g of cheese curd. All myzithra cheeses were placed into different sterile paper containers and stored for 30 days at 4°C.

2.4 Microbiological analysis of myzithra cheese

Representative 10g portions of myzithra cheese samples were obtained at various time intervals (1st, 5th, 7th, 14th, 30th storage day at 4°C) and blended with 90 mL of sterile trisodium citrate (2% w/v) solution and mixed in a stomacher (Bagmixer 400, Model VW, Interscience). The solution was then subjected to serial dilutions of 9mL of Ringer solution ¼ strength. Enumeration of viable cell counts of total aerobic

mesophilic bacteria, lactococci, lactobacilli, enterobacteria, coliforms, yeasts & fungi and staphylococci were performed in triplicate by pour plating 0.1 mL or 1mL of appropriate dilutions on the selective media for each species and according to instructions of the manufacturer as described previously by A. Terpou et al. (2017). Cell counts were expressed as log of mean colony-forming units.

For the enumeration of *L. casei* cells MRS-V agar (0.1% vancomycin) was prepared which according to Tharmaraj and Shah (2003) would disintegrate the adjunct probiotic strain and *L. bulgaricus ssp. delbrueckii* cell counts. The growth potential of *L. bulgaricus* was expected in higher number than any lactobacilli naturally occurring microflora since the whey used as raw material for myzithra cheese production was not sterilized and was obtained after Graviera cheese manufacture. Nowadays, in industrial dairies, cheese milk is at first pasteurized and then used with the addition of a starter mesophilic or thermophilic culture for Graviera cheese manufacture (Litopoulou-Tzanetaki & Tzanetakis, 2011). In this case Graviera cheese was manufactured by the addition of yogurt culture (*L. bulgaricus ssp. delbrueckii* and *S. thermophilus*) and rennet enzyme thus, the presence of *L. bulgaricus* cells can be expected. Cell counts were expressed as % of *L. casei* viability during 30 days of refrigerated storage.

2.5 Physicochemical analysis

The pH values of cheese whey during cheese production and of myzithra cheese during storage were measured using a digital pH meter by direct immersion of the electrode (EPI-BION SENTRON pH-System 1001).

Myzithra cheese samples (20 g each) were macerated with warm water (40 °C) to produce a total volume of 210 mL and then each sample was filtered and used for

the identification of total acidity and lactose concentration. A quantity of 25mL from the above filtrate was used for titration with 0,1 N NaOH and phenolphthalein indicator. Total acidity was determined to the official method by AOAC International (2000) and expressed as lactic acid content. Lactose was determined by high performance liquid chromatography, using a Shimadzu chromatograph with a SCR-101 N stainless steel column, a LC-9A pump, a CTO-10A oven at 60 °C and a RID-6A refractive index detector as described previously by A. Terpou et al. (2017). Lactose concentrations were calculated using standard curves.

2.6 Solid phase microextraction gas chromatography–mass spectrometry analysis

Samples of myzithra cheese with adjunct free (F) or adjunct microencapsulated (ME) *L. casei* cells were studied for terpenoid content as an aroma profile indicator using SPME GC–MS analysis and compared with control myzithra cheese samples (C). For the analysis, myzithra cheese samples (7.0 g each) from the 1st day of storage (4°C) were introduced into a 20mL headspace vial fitted with a Teflon-lined septum and sealed with an aluminum crimp seal. Through the seal a syringe needle (Supelco, Bellefonte, PA, USA) was inserted. The container was then thermostated for 5min at 60 °C with the syringe closed and when the temperature was stable the syringe was introduced at the gas area of the vial for 45 min at 60 °C.

The absorbed volatile analytes were then analyzed by GC–MS (Shimadzu GC-17A, MS QP5050, capillary column Supelco CO Wax-10 60 m, 0.32 mm i.d., 0.25 µm film thickness) as described previously by A. Terpou et al. (2017). The identification of the absorbed terpenoid content, presented % in total area of hydrocarbons, was performed by comparing the retention times with those of authentic compounds, by mass spectra of the authentic compounds generated in the

laboratory, by mass spectra obtained from NIST107, NIST21 and SZTERP libraries and by determining Kovats' retention indexes compared with those reported in the literature. Kovats' retention indexes (KI) were determined by injection of a standard mixture containing the homologous series of normal alkanes (C7–C32) in pure hexane under exactly the same experimental conditions, as described above.

2.7 *Myzithra* cheese sensory evaluation

Sensory evaluation of cheese is necessary in order to determine the influence of cheese composition on sensory characteristics, eating quality and consumers acceptability. Sensory evaluation was carried out by 10 laboratory members, priority trained, using locally approved protocols. Samples were tested by two different laboratories (5 members each) and all members were from different parts of the country. Consumers selection criteria were between 20 ~ 45 years of age, and frequent users of cheese (>once a week). The questions asked and procedure of cheese testing were identical for the two laboratories. *Myzithra* cheese samples from the 1st storage day (4°C) were placed into equivalent amounts of 5 × 5 cm and served at room temperature (18–22 °C). This procedure was chosen as consumers normally will consume cheese directly from refrigerator. Sensory analysis was carried out in panel booths conforming to international standards (International standard, 2007). The samples were coded by a different 3-digital number each and were served in a randomized order while the panel was asked to evaluate all *myzithra* cheeses (C, F, ME, MF) on a 0–10 scale (the higher the number the greater the intensity) based on saltiness, acidity, bitterness, sweetness, chewiness, cheese odor, mastic odor and overall acceptability. Data from both laboratories were handled as one data set during statistical analysis. The results are presented as a star chart of the product's attributes.

261

262 2.8 *Experimental design and statistical analysis*

263 Myzithra cheese production and analysis was carried out in triplicate and results are
264 presented as mean values $\pm$ standard deviation. All experiments were designed and
265 analyzed statistically by ANOVA. Significant differences among results (coefficients,
266 ANOVA tables and significance) which were computed using SPSS v.8.5.

3. Results and Discussion

3.1 Rational

Whey cheeses are prepared by denaturation and precipitation of whey proteins (α -lactalbumin, β -lactoglobulin) achieved by heating whey effluent at approximately at 85 °C. In Greece traditionally, whey cheeses like myzithra cheese are consumed as table cheeses. They have high nutritional value, low fat and salt content and have good organoleptic characteristics and are therefore great vehicles for incorporation of probiotics (Papaioannou, Chouliara, Karatapanis, Kontominas, & Savvaidis, 2007).

However, freshly made whey cheeses have a pH value of greater than 6, high moisture content and a low salt concentration and are therefore considered to be extremely sensitive to microbial deterioration (Hough, Puglieso, Sanchez, & da Silva, 1999) with very short shelf-life reaching up to 7 days under aerobic conditions (Samelis, Kakouri, Rogga, Savvaidis, & Kontominas, 2003).

In contrast, *Pistacia terebinthus* resin has been proved to confer an antimicrobial effect to dairy products due to its high terpenoid characteristics (Schoina et al., 2014). Moreover, immobilisation in various natural supports has been proved to enhance the viability of probiotic cells by a protective film that is formed by each immobilisation support protecting cells against the acidic environment of dairy products (Bosnea, Kopsahelis, Kokkali, Terpou, & Kanellaki, 2017; Antonia Terpou et al., 2017; A. Terpou et al., 2017). Thus, *Pistacia terebinthus* resin was assessed as encapsulation matrix for *Lactobacillus casei* cells and incorporated in myzithra cheese.

3.2 The probiotic encapsulated biocatalyst

Electron micrographs reinsured encapsulation of the probiotic cells within the viscous

matrix of *Pistacia terebinthus* (Fig. 1). The average encapsulation yield obtained in the present study was reported by experiments carried out on the encapsulated biocatalyst presenting an average of 1.56 g of *L. casei* cells successfully encapsulated in 100 g of *Pistacia terebinthus* (data not shown). More analytically, in each 5g of the resin 7.8% of the initial probiotic culture is encapsulated (data not shown). Subsequently, 4 log CFU of *L. casei* cells were proved to be encapsulated in each gram of *Pistacia terebinthus*' viscus matrix.

3.3 Effect of the encapsulated biocatalyst on growth of foodborne pathogens, cheese microflora, and spoilage microorganisms

Myzithra cheese samples were tested for their microbial stability as a shelf-life indicator, through refrigerated storage for 30 days (Table 1). No coliforms, enterobacteria or *Staphylococcus aureus* were detected in any myzithra cheese. There was observed a significant amount of yeast and fungi in all myzithra cheeses from the 1st storage day ($2.6 \sim 2.7 \log \text{CFU g}^{-1}$) which in the case of control cheeses (C) increased significantly until the 30th storage day ($4.7 \log \text{CFU g}^{-1}$). On the other hand, yeast and fungi showed a sharp decrease in the case of cheese samples with incorporated *Pistacia terebinthus* resin either added as an encapsulation support (ME) or as an adjunct (MF). Myzithra cheese which is usually consumed within 7 days from production and does not have yeast and fungi as a naturally occurring microflora. When high numbers of the prementioned microorganisms are detected the cheese is most likely accompanied by a bad odor and cannot be consumed.

The addition of the probiotic strain, either free or encapsulated, affected significantly ($P < 0.05$) the total lactobacilli counts of myzithra cheese after the 1st storage day (F, ME, MF), compared to control cheese (C). Moreover, in all myzithra

samples was detected a significant amount of Lactococci ($2.9 \sim 2.0 \log \text{CFU g}^{-1}$) that did not significantly differ among cheese samples during 30 days of storage. Their presence in cheeses may occur due to non-starter lactic acid bacteria and cross-contamination during cheese production (Kalogridou-Vassiliadou, Tzanetakis, & Litopoulou-Tzanetaki, 1994).

3.4 Growth capacity of the adjunct probiotic strain during refrigerated storage

Cheese presents a good vehicle for the delivery of probiotics in the intestine, while the maintenance of viable probiotic cell counts at high level by the end of expiration date is most crucial in such products in order to confer most health benefits. In this vein, experiments were carried out in order to evaluate the effect of *Pistacia terebinthus* resin and myzithra cheese storage conditions on viability of *L. casei* cells. Figure 2 shows the % viability of *L. casei* added either as a free culture, as an encapsulated culture in *Pistacia terebinthus* resin or as a free culture along with resin particles, during storage at 4°C for 30 days. All myzithra cheeses in which *L. casei* was incorporated (F, ME, MF), was observed a high count of viable cell (10^9CFU g^{-1} , data not shown) during storage. This result indicated the probiotic character of the products.

More specifically, the survival rates of encapsulated *L. casei* was 8.2% by the end of storage period while in contrast *L. casei* free cells along with resins incorporated particles were reduced down to 4%. The lowers viability rates were observed in the case of free *L. casei* cells which were reduced down to 8.2%. These results occurred most likely due to the absence of *P. terebinthus* resin and its antimicrobial effects allowing foodborne microorganisms like yeast and fungi (Table 1) to grow at the expense of probiotic cells. Since myzithra cheeses with encapsulated

biocatalyst showed the higher survival rates during storage we can assume that encapsulation acts protectively to probiotic cells against storage conditions and cheese environment.

3.5 Physicochemical characteristics of myzithra cheeses

During storage, lactose concentration, pH values and total acidity were determined for all myzithra cheeses and the results are presented in Table 2. In most whey cheeses prepared by the incorporation of lactic acid bacteria there has been observed a higher content of total acidity and a parallel with pH decrease, compared to whey cheeses prepared with the traditional recipe (Madureira et al., 2008; Madureira et al., 2015). As expected, a continuous increase of total acidity was observed during 30 storage days and ranged in acceptable levels for all myzithra cheese products (Anifantakis, 1991; Kalogridou-Vassiliadou et al., 1994). The total acidity of the whey cheeses was affected by the adjunct probiotic culture. Specifically, myzithra cheeses with *L. casei* culture (free or encapsulated) showed an acidity significantly higher ($P < 0.05$) compared to control myzithra cheeses prepared with no additional culture. In particular, total acidity increased from 0.3 to 0.5 g of lactic acid per 100 g of cheese in myzithra samples free *L. casei* cells (F) and an increase from 0.3 to 0.6 g of lactic acid / 100 g of cheese in the case of myzithra samples with encapsulated *L. casei* cells (ME). No significant differences were observed in myzithra cheese samples with free *L. casei* cells (F) and cheese samples with *Pistacia terebinthus* resin and free *L. casei* cells (MF). In parallel with total acidity increase, there was observed a pH reduction most likely as a result of continuous growth of microorganisms during refrigerated storage as it can be reinsured by microbiological analysis (Table 1). Regarding the pH of traditional whey cheese produced without an adjunct culture the presented values

are significantly higher (Pappa, Samelis, Kondyli, & Pappas, 2016) than those of whey cheeses like myzithra produced with the incorporation *L. casei*.

Lactose accumulation was observed in all cheese samples and ranged in usual levels of commercial myzithra cheeses (Kaminarides, 2015), while no significant differences were observed within the samples. Nevertheless, lactose content was lower in myzithra cheeses produced with either free or encapsulated probiotic cells compared to traditionally made myzithra cheese. This trend was expected due to the presence of live probiotic cells during storage (Kourkoutas et al., 2006; A. Terpou et al., 2017).

3.6 Effect of *Pistacia terebinthus* resin on volatile by-products of myzithra cheese

The majority of volatiles, especially ones that define a characteristic aroma to cheese products, were identified in myzithra cheeses with encapsulated biocatalyst (ME) by SPME/ GC-MS analysis and are presented as % of the total area of hydrocarbons' content (Table 3). A large number of terpenoid compounds, predominantly α -pinene (Table 3), appear in the produced myzithra cheeses with encapsulated *L. casei* cells (ME), due solely to added *Pistacia terebinthus* resin (pissa Paphos). Additionally, in Figure 3A is highlighted the plethora of terpenoid compounds of myzithra cheese with encapsulated biocatalyst (ME) presented on the chromatograph of headspace analysis. Concerning total terpenoid content (Figure 3B), total mono-terpenes were found to be 90.9% and total oxygenated mono-terpenes 9.1%. Monoterpenes have been reported as potential antimicrobial agents, as antiviral agents, antifungal agents and as potential antioxidants while they can also be used as ingredients of soaps, perfumes, and food additives (Armaka, Papanikolaou, Sivropoulou, & Arsenakis, 1999; Prates et al., 1998; Ruberto & Baratta, 2000; Zhang et al., 2016). Figure 3B shows the distribution

of terpenoid components of myzithra cheese with encapsulate biocatalyst (0.5/ 100 g of product), in which the higher content of monoterpenes was detected as is α -pinene (84.5%). These results were mostly expected as essential oils of three *Pistacia* species compile mainly of α -pinene, β -pinene, limonene and α -terpineol (Duru et al., 2003). Likewise, the significant influence of α -pinene in myzithra cheeses with encapsulated biocatalyst indicates the reins' antimicrobial effects against spoilage microorganisms (Kivrak et al., 2009) resulting to a possibly extended shelf-life of functional myzithra cheeses. Moreover, apart from α -pinene the detected α -terpineol (1.3%), eucalyptol (0.3%) and linalool (0.2%) can provide antibacterial and antioxidant effects to produced cheeses (Zengin & Baysal, 2014). In addition, according Park et al. (2012) linalool and α -terpineol provides an antimicrobial effect against periodontopathic and cariogenic bacteria. Thus, the antibacterial, antifungal and antioxidant activities of such monoterpenes result to the conclusion that *P. terebinthus* resin can be used as a natural preservative of food products with good organoleptic characteristics.

Another important influence of the incorporated resin with encapsulated probiotic cells refers to improved aromatic profile of myzithra cheeses. The numerous monoterpenes originating from the incorporated resin are well known for their contribution to the aromatic profile of products and are very often used as additives in food production (Prates et al., 1998). Most of detected terpenes are characterized by exceptional aromatic characteristics and can contribute to flavor of produced cheese due to their low threshold value. For example, α -pinene is known for its pine odor, 3-carene (0.2%) its sweet lemon odor and linalool (0.2%) for its sweet floral odor (Curioni & Bosset, 2002; Vichi et al., 2007). Apart from their floral and fruity aromas, monoterpenes are also considered important compounds because of their ability to reduce the effects of unpleasant odors caused by phenolic compounds or

short chain fatty acids (Curioni & Bosset, 2002).

The results indicated that the plethora of terpenoid compounds were detected in myzithra cheese due to the incorporation of *P. terebinthus* resin. In addition, by GCMS/ SPME analysis has highlighted that the use of *P. terebinthus* resin as encapsulation support leads to exceptional aromatic characteristics of produced myzithra cheeses, which is also in agreement with the sensory evaluation results.

3.7 *Myzithra* cheese sensory evaluation

Sensory evaluation of myzithra cheese samples is presented in Figure 4. The results of sensory evaluation of myzithra cheese samples are presented in Figure 4. In most cases, no significant differences were observed between myzithra cheeses produced by adjunct *P. terebinthus* resin with either free or encapsulated probiotic cells. However, samples prepared without the present of the resin (C, F) presented significantly ($P < 0.05$) lower values in all cases compared to myzithra cheese samples in which *P. terebinthus* resin was added as an adjunct. These findings indicated the high industrialization potential of the proposed technology since sensory evaluation showed high consumers preference referring to all novel myzithra cheeses with incorporated *Pistacia terebinthus* resin.

4. Conclusions

The production of a novel functional whey cheese by the adjunct encapsulated *L. casei* in *Pistacia terebinthus* resin (pissa Paphos) was assessed in the present study. The probiotic cells were successfully encapsulated in the viscous matrix of the resin retaining its viability despite the resins' antimicrobial properties. The obtained results showed that encapsulation favored the viability of *L. casei* in refrigerated storage

while the antimicrobial properties of the resin resulted in shelf-life extension of the produced myzithra cheeses. In the present study, was highlighted the potential use of pissa Paphos as a lactobacilli microencapsulation support, as an antimicrobial additive and as an aromatic enhancement material indicating its future use in nutraceutical and food industry.

5. Acknowledgements

Schoina V. would like to thank the State Scholarships Foundation (IKY) for the financial support in the frame of her PhD thesis.

6. References

- Aksoy, A., Duran, N., & Koksall, F. (2006). In vitro and in vivo antimicrobial effects of mastic chewing gum against *Streptococcus mutans* and *mutans streptococci*. Archives of Oral Biology, 51(6), 476-481. doi: <https://doi.org/10.1016/j.archoralbio.2005.11.003>
- Anifantakis, E. M. (1991). Greek Cheeses: A Tradition of Centuries: National Dairy Committee of Greece.
- Armaka, M., Papanikolaou, E., Sivropoulou, A., & Arsenakis, M. (1999). Antiviral properties of isoborneol, a potent inhibitor of herpes simplex virus type 1. Antiviral Res, 43(2), 79-92.
- Bogue, J., Collins, O., & Troy, A. J. (2017). Chapter 2 - Market analysis and concept development of functional foods - Bagchi, Debasis. In S. Nair (Ed.), Developing New Functional Food and Nutraceutical Products (pp. 29-45). San Diego: Academic Press.
- Bosnea, L. A., Kopsahelis, N., Kokkali, V., Terpou, A., & Kanellaki, M. (2017). Production of a novel probiotic yogurt by incorporation of *L. casei* enriched fresh apple pieces, dried raisins and wheat grains. Food and Bioproducts Processing, 102, 62-71. doi: [10.1016/j.fbp.2016.11.010](https://doi.org/10.1016/j.fbp.2016.11.010)
- Bosnea, L. A., Moschakis, T., Nigam, P. S., & Biliaderis, C. G. (2017). Growth adaptation of probiotics in biopolymer-based coacervate structures to enhance cell viability. LWT - Food Science and Technology, 77, 282-289. doi: [10.1016/j.lwt.2016.11.056](https://doi.org/10.1016/j.lwt.2016.11.056)
- Bozoudi, D., Pavlidou, S., Kotzamanidis, C., Georgakopoulos, P., Torriani, S.,

- 473 Kondyli, E., . . . Litopoulou-Tzanetaki, E. (2016). "Graviera Naxou and Graviera
474 Kritis Greek PDO cheeses: Discrimination based on microbiological and
475 physicochemical criteria and volatile organic compounds profile". *Small Ruminant*
476 *Research*, 136, 161-172. doi: <https://doi.org/10.1016/j.smallrumres.2016.01.022>
- 477 Caleja, C., Barros, L., Antonio, A. L., Carcho, M., Oliveira, M. B. P. P., & Ferreira, I.
478 C. F. R. (2016). Fortification of yogurts with different antioxidant preservatives: A
479 comparative study between natural and synthetic additives. *Food Chemistry*, 210, 262-
480 268. doi: <https://doi.org/10.1016/j.foodchem.2016.04.114>
- 481 Carcho, M., Barreiro, M. F., Morales, P., & Ferreira, I. C. F. R. (2014). Adding
482 Molecules to Food, Pros and Cons: A Review on Synthetic and Natural Food
483 Additives. *Comprehensive Reviews in Food Science and Food Safety*, 13(4), 377-399.
484 doi: 10.1111/1541-4337.12065
- 485 Champagne, C. P., Ross, R. P., Saarela, M., Hansen, K. F., & Charalampopoulos, D.
486 (2011). Recommendations for the viability assessment of probiotics as concentrated
487 cultures and in food matrices. *International Journal of Food Microbiology*, 149(3),
488 185-193. doi: [10.1016/j.ijfoodmicro.2011.07.005](https://doi.org/10.1016/j.ijfoodmicro.2011.07.005)
- 489 Chemists, A. o. O. A. (2000). Official methods of analysis of the AOAC: Association
490 of Official Analytical Chemists.
- 491 Curioni, P. M. G., & Bosset, J. O. (2002). Key odorants in various cheese types as
492 determined by gas chromatography-olfactometry. *International Dairy Journal*, 12(12),
493 959-984. doi: [10.1016/S0958-6946\(02\)00124-3](https://doi.org/10.1016/S0958-6946(02)00124-3)
- 494 Daifas, D. P., Smith, J. P., Blanchfield, B., Sanders, G., Austin, J. W., & Koukoutisis,
495 J. (2004). Effects of mastic resin and its essential oil on the growth of proteolytic

- 496 *Clostridium botulinum*. International Journal of Food Microbiology, 94(3), 313-322.
497 doi: <https://doi.org/10.1016/j.ijfoodmicro.2004.01.017>
- 498 Dell, B., & McComb, A. J. (1979). Plant Resins—Their Formation, Secretion and
499 Possible Functions. Advances in Botanical Research, 6, 277-316. doi:
500 [http://dx.doi.org/10.1016/S0065-2296\(08\)60332-8](http://dx.doi.org/10.1016/S0065-2296(08)60332-8)
- 501 Duru, M. E., Cakir, A., Kordali, S., Zengin, H., Harmandar, M., Izumi, S., & Hirata, T.
502 (2003). Chemical composition and antifungal properties of essential oils of three
503 *Pistacia* species. Fitoterapia, 74(1–2), 170-176. doi: [https://doi.org/10.1016/S0367-](https://doi.org/10.1016/S0367-326X(02)00318-0)
504 [326X\(02\)00318-0](https://doi.org/10.1016/S0367-326X(02)00318-0)
- 505 Fuller, R., & Gibson, G. R. (1998). Probiotics and prebiotics: microflora management
506 for improved gut health. Clinical Microbiology and Infection, 4(9), 477-480. doi:
507 <http://dx.doi.org/10.1111/j.1469-0691.1998.tb00401.x>
- 508 Gardeli, C., Papageorgiou, V., Mallouchos, A., Kibouris, T., & Komaitis, M. (2008).
509 Essential oil composition of *Pistacia lentiscus* L. and *Myrtus communis* L.: Evaluation
510 of antioxidant capacity of methanolic extracts. Food Chemistry, 107(3), 1120-1130.
511 doi: <https://doi.org/10.1016/j.foodchem.2007.09.036>
- 512 Goodner, K. L. (2008). Practical retention index models of OV-101, DB-1, DB-5, and
513 DB-Wax for flavor and fragrance compounds. LWT - Food Science and Technology,
514 41(6), 951-958. doi: <https://doi.org/10.1016/j.lwt.2007.07.007>
- 515 Högnadóttir, Á., & Rouseff, R. L. (2003). Identification of aroma active compounds in
516 orange essence oil using gas chromatography–olfactometry and gas chromatography–
517 mass spectrometry. Journal of Chromatography A, 998(1–2), 201-211. doi:
518 [https://doi.org/10.1016/S0021-9673\(03\)00524-7](https://doi.org/10.1016/S0021-9673(03)00524-7)

- 519 Hough, G., Puglieso, M. L., Sanchez, R., & da Silva, O. M. (1999). Sensory and
 520 Microbiological Shelf-Life of a Commercial Ricotta Cheese. *Journal of Dairy*
 521 *Science*, 82(3), 454-459. doi: [http://dx.doi.org/10.3168/jds.S0022-0302\(99\)75253-7](http://dx.doi.org/10.3168/jds.S0022-0302(99)75253-7)
- 522 International standard ISO 8589: Sensory analysis: General guidance for the design of
 523 test rooms. International Organization for Standardization (2007).
- 524 Kalogridou-Vassiliadou, D., Tzanetakis, N., & Litopoulou-Tzanetaki, E. (1994).
 525 Microbiological and physicochemical characteristics of 'Anthotyro', a Greek
 526 traditional whey cheese. *Food Microbiology*, 11(1), 15-19. doi:
 527 <https://doi.org/10.1006/fmic.1994.1003>
- 528 Kaminarides, S. (2015). A modified form of Myzithra cheese produced by substituting
 529 the fresh cheese whey by dried whey protein concentrate and ovine milk and cream.
 530 *Small Ruminant Research*, 131, 118-122. doi:
 531 <https://doi.org/10.1016/j.smallrumres.2015.07.020>
- 532 Kivrak, İ., Duru, M. E., Öztürk, M., Mercan, N., Harmandar, M., & Topçu, G. (2009).
 533 Antioxidant, anticholinesterase and antimicrobial constituents from the essential oil
 534 and ethanol extract of *Salvia potentillifolia*. *Food Chemistry*, 116(2), 470-479. doi:
 535 <http://dx.doi.org/10.1016/j.foodchem.2009.02.069>
- 536 Kourkoutas, Y., Bosnea, L., Taboukos, S., Baras, C., Lambrou, D., & Kanellaki, M.
 537 (2006). Probiotic cheese production using *Lactobacillus casei* cells immobilized on
 538 fruit pieces. [Article]. *Journal of Dairy Science*, 89(5), 1439-1451.
- 539 Lardos, A. (2006). The botanical materia medica of the Iatrosophikon—A collection
 540 of prescriptions from a monastery in Cyprus. *Journal of Ethnopharmacology*, 104(3),
 541 387-406. doi: <https://doi.org/10.1016/j.jep.2005.12.035>

- 542 Lee, S.-J., Umamo, K., Shibamoto, T., & Lee, K.-G. (2005). Identification of volatile
543 components in basil (*Ocimum basilicum* L.) and thyme leaves (*Thymus vulgaris* L.)
544 and their antioxidant properties. Food Chemistry, 91(1), 131-137. doi:
545 <https://doi.org/10.1016/j.foodchem.2004.05.056>
- 546 Litopoulou-Tzanetaki, E., & Tzanetakis, N. (2011). Microbiological characteristics of
547 Greek traditional cheeses. Small Ruminant Research, 101(1–3), 17-32. doi:
548 <https://doi.org/10.1016/j.smallrumres.2011.09.022>
- 549 Madureira, A. R., Soares, J. C., Pintado, M. E., Gomes, A. M. P., Freitas, A. C., &
550 Malcata, F. X. (2008). Sweet whey cheese matrices inoculated with the probiotic
551 strain *Lactobacillus paracasei* LAFTI® L26. Dairy Science & Technology, 88(6),
552 649-665. doi: 10.1051/dst:2008026
- 553 Madureira, A. R., Soares, J. C., Pintado, M. E., Gomes, A. M. P., Freitas, A. C., &
554 Xavier Malcata, F. (2015). Effect of the incorporation of salted additives on probiotic
555 whey cheeses. Food Bioscience, 10, 8-17. doi:
556 <https://doi.org/10.1016/j.fbio.2015.01.004>
- 557 Mallouchos, A., Paul, L., Bekatorou, A., Koutinas, A., & Komaitis, M. (2007).
558 Ambient and low temperature winemaking by immobilized cells on brewer's spent
559 grains: Effect on volatile composition. Food Chemistry, 104(3), 918-927. doi:
560 <https://doi.org/10.1016/j.foodchem.2006.12.047>
- 561 Morales, M. E., & Ruiz, M. A. (2016). 16 - Microencapsulation of probiotic cells:
562 applications in nutraceutic and food industry - Grumezescu, Alexandru Mihai
563 Nutraceuticals (pp. 627-668): Academic Press.
- 564 Morkhade, D. M. Evaluation of gum mastic (*pistacia lentiscus*) as a

- 565 microencapsulating and matrix forming material for sustained drug release. Asian
 566 Journal of Pharmaceutical Sciences. doi: <https://doi.org/10.1016/j.ajps.2017.05.002>
- 567 Papaioannou, G., Chouliara, I., Karatapanis, A. E., Kontominas, M. G., & Savvaidis,
 568 I. N. (2007). Shelf-life of a Greek whey cheese under modified atmosphere
 569 packaging. International Dairy Journal, 17(4), 358-364. doi:
 570 <https://doi.org/10.1016/j.idairyj.2006.04.001>
- 571 Pappa, E. C., Samelis, J., Kondyli, E., & Pappas, A. C. (2016). Characterisation of
 572 Urda whey cheese: Evolution of main biochemical and microbiological parameters
 573 during ripening and vacuum packaged cold storage. International Dairy Journal, 58,
 574 54-57. doi: <https://doi.org/10.1016/j.idairyj.2015.12.016>
- 575 Pappas, C., & Voutsinas, L. (2009). Effect of acidification of sheep's cheese whey
 576 with organic acids on the retention of protein and on the quality of Myzithra cheese.
 577 Journal of Dairy Research, 55(2), 247-254. doi: 10.1017/S0022029900026078
- 578 Paraschos, S., Magiatis, P., Gousia, P., Economou, V., Sakkas, H., Papadopoulou, C.,
 579 & Skaltsounis, A.-L. (2011). Chemical investigation and antimicrobial properties of
 580 mastic water and its major constituents. Food Chemistry, 129(3), 907-911. doi:
 581 <https://doi.org/10.1016/j.foodchem.2011.05.043>
- 582 Park, S.-N., Lim, Y. K., Freire, M. O., Cho, E., Jin, D., & Kook, J.-K. (2012).
 583 Antimicrobial effect of linalool and α -terpineol against periodontopathic and
 584 cariogenic bacteria. Anaerobe, 18(3), 369-372. doi:
 585 <http://dx.doi.org/10.1016/j.anaerobe.2012.04.001>
- 586 Prates, H. T., Santos, J. P., Waquil, J. M., Fabris, J. D., Oliveira, A. B., & Foster, J. E.
 587 (1998). Insecticidal activity of monoterpenes against *Rhyzopertha dominica* (F.) and

- 588 *Tribolium castaneum* (Herbst). Journal of Stored Products Research, 34(4), 243-249.
 589 doi: [http://dx.doi.org/10.1016/S0022-474X\(98\)00005-8](http://dx.doi.org/10.1016/S0022-474X(98)00005-8)
- 590 Rauf, A., Patel, S., Uddin, G., Siddiqui, B. S., Ahmad, B., Muhammad, N., . . . Hadda,
 591 T. B. (2017). Phytochemical, ethnomedicinal uses and pharmacological profile of
 592 genus *Pistacia*. Biomedicine & Pharmacotherapy, 86, 393-404. doi:
 593 <https://doi.org/10.1016/j.biopha.2016.12.017>
- 594 Ruberto, G., & Baratta, M. T. (2000). Antioxidant activity of selected essential oil
 595 components in two lipid model systems. Food Chemistry, 69(2), 167-174. doi:
 596 [http://dx.doi.org/10.1016/S0308-8146\(99\)00247-2](http://dx.doi.org/10.1016/S0308-8146(99)00247-2)
- 597 Samelis, J., Kakouri, A., Rogga, K. J., Savvaidis, I. N., & Kontominas, M. G. (2003).
 598 Nisin treatments to control *Listeria monocytogenes* post-processing contamination on
 599 Anthotyros, a traditional Greek whey cheese, stored at 4°C in vacuum packages. Food
 600 Microbiology, 20(6), 661-669. doi: [https://doi.org/10.1016/S0740-0020\(03\)00018-2](https://doi.org/10.1016/S0740-0020(03)00018-2)
- 601 Saxami, G., Ypsilantis, P., Sidira, M., Simopoulos, C., Kourkoutas, Y., & Galanis, A.
 602 (2012). Distinct adhesion of probiotic strain *Lactobacillus casei* ATCC393 to rat
 603 intestinal mucosa. Anaerobe, 18(4), 417-420. doi:
 604 <http://dx.doi.org/10.1016/j.anaerobe.2012.04.002>
- 605 Schoina, V., Terpou, A., Angelika-Ioanna, G., Koutinas, A., Kanellaki, M., & Bosnea,
 606 L. (2014). Use of *Pistacia terebinthus* resin as immobilization support for
 607 *Lactobacillus casei* cells and application in selected dairy products. Journal of Food
 608 Science and Technology. doi: 10.1007/s13197-014-1627-9
- 609 Schoina, V., Terpou, A., Angelika-Ioanna, G., Koutinas, A., Kanellaki, M., & Bosnea,
 610 L. (2015). Use of *Pistacia terebinthus* resin as immobilization support for

- 611 *Lactobacillus casei* cells and application in selected dairy products. Journal of Food
 612 Science and Technology, 52(9), 5700-5708. doi: 10.1007/s13197-014-1627-9
- 613 Shiratsuchi, H., Shimoda, M., Minegishi, Y., & Osajima, Y. (1993). Isolation and
 614 identification of volatile flavor compounds in nonfermented coarse-cut sausage.
 615 Flavor as a quality factor of nonfermented sausage. 1. Journal of Agricultural and
 616 Food Chemistry, 41(4), 647-652. doi: 10.1021/jf00028a027
- 617 Shori, A. B. (2015). The potential applications of probiotics on dairy and non-dairy
 618 foods focusing on viability during storage. Biocatalysis and Agricultural
 619 Biotechnology, 4(4), 423-431. doi: <https://doi.org/10.1016/j.bcab.2015.09.010>
- 620 Terpou, A., Bekatorou, A., Kanellaki, M., Koutinas, A. A., & Nigam, P. (2017).
 621 Enhanced probiotic viability and aromatic profile of yogurts produced using wheat
 622 bran (*Triticum aestivum*) as cell immobilization carrier. Process Biochemistry, 55, 1-
 623 10. doi: <https://doi.org/10.1016/j.procbio.2017.01.013>
- 624 Terpou, A., Gialleli, A. I., Bosnea, L., Kanellaki, M., Koutinas, A. A., & Castro, G. R.
 625 (2017). Novel cheese production by incorporation of sea buckthorn berries
 626 (*Hippophae rhamnoides L.*) supported probiotic cells. [Article]. LWT - Food Science
 627 and Technology, 79, 616-624. doi: 10.1016/j.lwt.2016.11.021
- 628 Tharmaraj, N., & Shah, N. P. (2003). Selective Enumeration of *Lactobacillus*
 629 *delbrueckii ssp. bulgaricus*, *Streptococcus thermophilus*, *Lactobacillus acidophilus*,
 630 *Bifidobacteria*, *Lactobacillus casei*, *Lactobacillus rhamnosus*, and *Propionibacteria*.
 631 Journal of Dairy Science, 86(7), 2288-2296. doi: [https://doi.org/10.3168/jds.S0022-](https://doi.org/10.3168/jds.S0022-0302(03)73821-1)
 632 0302(03)73821-1
- 633 Topçu, G., Ay, M., Bilici, A., Sarıkürkcü, C., Öztürk, M., & Ulubelen, A. (2007). A

- 634 new flavone from antioxidant extracts of *Pistacia terebinthus*. Food Chemistry,
635 103(3), 816-822. doi: <https://doi.org/10.1016/j.foodchem.2006.09.028>
- 636 Vichi, S., Riu-Aumatell, M., Mora-Pons, M., Guadayol, J. M., Buxaderas, S., &
637 López-Tamames, E. (2007). HS-SPME coupled to GC/MS for quality control of
638 *Juniperus communis* L. berries used for gin aromatization. Food Chemistry, 105(4),
639 1748-1754. doi: <https://doi.org/10.1016/j.foodchem.2007.03.026>
- 640 Zengin, H., & Baysal, A. H. (2014). Antibacterial and antioxidant activity of essential
641 oil terpenes against pathogenic and spoilage-forming bacteria and cell structure-
642 activity relationships evaluated by SEM microscopy. Molecules, 19(11), 17773-
643 17798. doi: [10.3390/molecules191117773](https://doi.org/10.3390/molecules191117773)
- 644 Zhang, Z., Yang, T., Mi, N., Wang, Y., Li, G., Wang, L., & Xie, Y. (2016). Antifungal
645 activity of monoterpenes against wood white-rot fungi. International Biodeterioration
646 & Biodegradation, 106, 157-160. doi: <http://dx.doi.org/10.1016/j.ibiod.2015.10.018>

Figure captions

Figure 1. Electron micrographs of *Pistacia terebinthus* resin surface (A), *L. casei* ATCC 393 cells in *Pistacia terebinthus* encapsulating matrix (B, C).

Figure 2. *L. casei* % viability during refrigerated storage (4°C) for 30 days.

Figure 3. SPME GC–MS spectra (full scan mode chromatogram) of myzithra cheese with adjunct *L. casei* cells encapsulated in *Pistacia terebinthus* resin (ME).

Figure 4. Sensory evaluation of produced myzithra chesses from the 1st day of storage at 4°C.

Table 1. Myzithra cheese microbial population (log CFU g⁻¹ ±) during 30 days of storage (4°C).

Myzithra cheese	Storage time (days)	Total aerobic counts	Lactococci	Lactobacilli	Yeasts & fungi	Staphylococci
C	1	3.78 ^{±0.19}	2.80 ^{±0.14}	2.97 ^{±0.15}	2.68 ^{±0.08}	1.98 ^{±0.10}
	5	4.30 ^{±0.21}	2.85 ^{±0.15}	2.66 ^{±0.13}	3.30 ^{±0.12}	1.92 ^{±0.09}
	7	4.65 ^{±0.23}	2.70 ^{±0.14}	2.57 ^{±0.13}	3.62 ^{±0.13}	2.14 ^{±0.11}
	14	4.75 ^{±0.24}	2.75 ^{±0.14}	2.85 ^{±0.15}	4.42 ^{±0.12}	2.21 ^{±0.11}
	30	5.78 ^{±0.29}	2.78 ^{±0.14}	2.74 ^{±0.19}	4.73 ^{±0.12}	2.25 ^{±0.11}
F	1	3.30 ^{±0.17}	2.83 ^{±0.15}	9.97 ^{±0.17}	2.60 ^{±0.08}	2.11 ^{±0.11}
	5	4.26 ^{±0.21}	2.85 ^{±0.19}	9.28 ^{±0.16}	2.90 ^{±0.10}	1.70 ^{±0.09}
	7	4.83 ^{±0.24}	2.18 ^{±0.16}	9.81 ^{±0.18}	2.04 ^{±0.10}	nd
	14	4.50 ^{±0.23}	2.23 ^{±0.16}	9.32 ^{±0.17}	2.95 ^{±0.10}	nd
	30	5.00 ^{±0.25}	2.58 ^{±0.18}	9.16 ^{±0.12}	2.80 ^{±0.09}	nd
ME	1	3.78 ^{±0.19}	2.90 ^{±0.15}	9.96 ^{±0.10}	2.74 ^{±0.09}	1.98 ^{±0.10}
	5	4.99 ^{±0.24}	2.52 ^{±0.17}	9.87 ^{±0.19}	1.80 ^{±0.09}	1.02 ^{±0.07}
	7	5.00 ^{±0.25}	2.48 ^{±0.18}	10.8 ^{±0.12}	1.63 ^{±0.09}	nd
	14	5.12 ^{±0.25}	2.20 ^{±0.16}	10.71 ^{±0.10}	1.26 ^{±0.08}	nd
	30	5.21 ^{±0.26}	2.85 ^{±0.14}	10.78 ^{±0.17}	1.00 ^{±0.05}	nd
MF	1	3.30 ^{±0.17}	2.69 ^{±0.14}	9.85 ^{±0.18}	2.70 ^{±0.09}	2.01 ^{±0.10}
	5	4.22 ^{±0.21}	2.20 ^{±0.16}	9.42 ^{±0.17}	1.84 ^{±0.12}	1.00 ^{±0.09}
	7	5.00 ^{±0.25}	2.11 ^{±0.20}	9.34 ^{±0.15}	1.36 ^{±0.11}	nd
	14	5.02 ^{±0.24}	2.06 ^{±0.15}	9.54 ^{±0.18}	1.11 ^{±0.11}	nd
	30	5.30 ^{±0.27}	2.01 ^{±0.15}	9.46 ^{±0.14}	1.05 ^{±0.07}	nd

*Variation within treatments is not greater than 10% in all cases.

Table 2. pH, total acidity and lactose content of myzithra whey cheeses during refrigerated (4°C) storage for 30 days.

myzithra cheese	Storage time (days)	pH	Total acidity (g lactic acid/ 100 g cheese)	Lactose (g /100 g cheese)
C	1	6.47 ^{±0.04}	0.19 ^{±0.01}	3.78 ^{±0.08}
	5	6.45 ^{±0.05}	0.20 ^{±0.01}	3.80 ^{±0.08}
	7	6.47 ^{±0.05}	0.20 ^{±0.01}	3.79 ^{±0.07}
	14	6.44 ^{±0.04}	0.21 ^{±0.01}	3.80 ^{±0.07}
	30	6.43 ^{±0.04}	0.21 ^{±0.01}	3.78 ^{±0.04}
F	1	6.34 ^{±0.04}	0.30 ^{±0.02}	3.79 ^{±0.05}
	5	6.22 ^{±0.03}	0.38 ^{±0.02}	3.66 ^{±0.06}
	7	6.16 ^{±0.04}	0.40 ^{±0.02}	3.63 ^{±0.07}
	14	6.08 ^{±0.04}	0.48 ^{±0.03}	3.60 ^{±0.06}
	30	6.00 ^{±0.04}	0.48 ^{±0.03}	3.57 ^{±0.04}
ME	1	6.37 ^{±0.04}	0.31 ^{±0.02}	3.78 ^{±0.07}
	5	6.17 ^{±0.04}	0.50 ^{±0.03}	3.67 ^{±0.06}
	7	6.00 ^{±0.03}	0.58 ^{±0.04}	3.55 ^{±0.05}
	14	5.94 ^{±0.03}	0.60 ^{±0.04}	3.53 ^{±0.07}
	30	5.92 ^{±0.04}	0.63 ^{±0.04}	3.51 ^{±0.07}
MF	1	6.33 ^{±0.04}	0.29 ^{±0.02}	3.78 ^{±0.04}
	5	6.19 ^{±0.04}	0.28 ^{±0.02}	3.67 ^{±0.04}
	7	6.21 ^{±0.04}	0.40 ^{±0.03}	3.63 ^{±0.07}
	14	6.04 ^{±0.03}	0.44 ^{±0.03}	3.61 ^{±0.06}
	30	6.03 ^{±0.03}	0.47 ^{±0.03}	3.59 ^{±0.06}

*Variation within treatments is not greater than 10% in all cases.

Table 3. Terpenoid content (%) of myzithra cheese samples (C, F, ME) from the 1st day of storage (4°C) identified by SPME GC/MS.

Compound name	ID*	KI	KI from literature	C	F	ME
Monoterpenes						
α -pinene	KI, MS	1025	1017 ^c 1020 ^f 1019 ^b	Nd	Nd	83.5 ^{±0.17}
camphene	KI, MS	1063	1053 ^c 1080 ^h 1063 ^a	Nd	Nd	0.8 ^{±0.05}
β -pinene	KI, MS	1105	1113 ^d 1108 ^b	Nd	Nd	2.2 ^{±0.13}
3-carene	KI, MS	1138	1114 ^e 1141 ^f	Nd	Nd	0.6 ^{±0.10}
β -myrcene	KI, MS	1152	1157 ^c 1152 ^a 1158 ^h	Nd	Nd	0.6 ^{±0.11}
2-carene	KI, MS	1164	1164 ^a	Nd	Nd	0.2 ^{±0.05}
D-limonen	KI, MS	1180	1188 ^e 1198 ^b	Nd	Nd	1.3 ^{±0.14}
Beta-phellandrene	KI, MS	1188	1188 ^a	Nd	Nd	0.1 ^{±0.03}
o-cymene	KI, MS	1250	1250 ^a	Nd	Nd	0.7 ^{±0.14}
Oxygenated Monoterpenes						
Eucalyptol	KI, MS	1196	1196 ^a	Nd	Nd	0.3 ^{±0.12}
Terpinolene	KI, MS	1259	1271 ^g	Nd	Nd	3.5 ^{±0.13}
Camphelanol	KI, MS	1482	1482 ^a	Nd	Nd	0.1 ^{±0.05}
Linalool	KI, MS	1531	1531 ^a	Nd	Nd	0.2 ^{±0.05}
Bornyl acetate	KI, MS	1571	1574 ^g	Nd	Nd	0.8 ^{±0.12}
4-terpineol	KI, MS	1593	1602 ^b 1593 ^d	Nd	Nd	0.3 ^{±0.14}
Pinocarveol	KI, MS	1651	1651 ^a	Nd	Nd	0.4 ^{±0.14}
Verbenol	KI, MS	1671	1671 ^a	Nd	Nd	0.3 ^{±0.05}
α -terpineol	KI, MS	1688	1691 ^g	Nd	Nd	1.3 ^{±0.12}
Melilotol	KI, MS	1783	1783 ^a	Nd	Nd	0.2 ^{±0.03}
Myrtenol	KI, MS	1787	1789 ^g	Nd	Nd	0.2 ^{±0.05}
p-cymene-8-ol	KI, MS	1842	1846 ^g	Nd	Nd	0.7 ^{±0.13}

*ID: Method of identification, KI = tentative identification by Kovats retention index in accordance with literature [^a: Schoina et al. (2014) ^b: Shiratsuchi, Shimoda, Minegishi, and Osajima (1993), ^c: Gardeli, Papageorgiou, Mallouchos, Kibouris, and Komaitis (2008), ^d: Högnadóttir and Rouseff (2003), ^e: Mallouchos, Paul, Bekatorou, Koutinas, and Komaitis (2007), ^f: Vichi et al. (2007), ^g: Lee, Umamo, Shibamoto, and Lee (2005), ^h: Goodner (2008)], MS = tentative identification by mass spectra obtained from NIST107, NIST21, and SZTERP libraries.

**Nd.: not detected

***Variation within treatments is not greater than 10% in all cases.

Figure captions

Figure 1. Electron micrographs of *Pistacia terebinthus* resin surface (A) and *L. casei* ATCC 393 cells in *Pistacia terebinthus* encapsulating matrix (B, C).

Figure 2. % Survival of *L. casei* in myzithra cheeses during refrigerated storage (4°C) for 30 days.

Figure 3. SPME GC–MS spectra (full scan mode chromatogram - A) presenting the detected terpenes (% total area of hydrocarbons - B) of myzithra cheese with *L. casei* encapsulated cells within *Pistacia terebinthus* resin (ME) from the 1st day of storage at 4°C.

Figure 4. Sensory evaluation of produced myzithra chesses from the 1st day of storage at 4°C.

Highlights

- Manufacture of a novel functional myzithra cheese with probiotic characteristics.
- *Pistacia terebinthus* viscus matrix as encapsulation support of probiotic *Lactobacillus* cells.
- Significant reduction of fungi/yeasts in cheeses with incorporated *Pistacia terebinthus* resin.
- SPME GC/MS analysis indicates the upgraded terpenoid profile of cheeses with *P. terebinthus* resin.
- Terpene exceptional aroma and possible antimicrobial/ antioxidant effects to cheese products.