The potential of siRNA based drug delivery in respiratory disorders: recent advances and progress

Tambuwalla, M. M. (2019). The potential of siRNA based drug delivery in respiratory disorders: recent advances and progress. *Drug Development Research*, 80(6), 714-730. <https://doi.org/10.1002/ddr.21571>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Drug Development Research

Publication Status:
Published (in print/issue): 30/09/2019

DOI:
[10.1002/ddr.21571](https://doi.org/10.1002/ddr.21571)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

The potential of siRNA based drug delivery in respiratory disorders: recent advances and progress

Kamal Dua^{*a,b,c}, Ridhima Wadhwa^d, Gautam Singhvi^e, VamshiKrishna Rapalli^e, Shakti Dhar Shukla^c, Madhur D. Shastri^f, Gaurav Gupta^g, Saurabh Satija^h, Meenu Mehta^h, Navneet Khurana^h, Rajendra Awasthiⁱ, Pawan Kumar Maurya^j, Lakshmi Thangavelu^k, S Rajesh Kumar^k, Murtaza M. Tambuwala^l, Trudi Collet^m, Philip M. Hansbro^{b,c,n}, Dinesh Kumar Chellappan^{o,*}

^aDiscipline of Pharmacy, Graduate School of Health, University of Technology Sydney, Ultimo NSW 2007, Australia.

^bCentenary Institute, Royal Prince Alfred Hospital, Camperdown, NSW 2050, Australia.

^cPriority Research Centre for Healthy Lungs, Hunter Medical Research Institute (HMRI) & School of Biomedical Sciences and Pharmacy, University of Newcastle, Callaghan, NSW 2308, Australia.

^dFaculty of Life Sciences and Biotechnology, South Asian University, Akbar Bhawan, Chanakyapuri, New Delhi 110021, India.

^eDepartment of Pharmacy, Birla Institute of Technology and Science (BITS), Pilani, 333031, India

^fSchool of Health Sciences, College of Health and Medicine, University of Tasmania, Launceston, Australia.

^gSchool of Pharmaceutical Sciences, Jaipur National University, Jagatpura, 302017, Jaipur, India

^hSchool of Pharmaceutical Sciences, Lovely Professional University, Jalandhar-Delhi G.T. Road (NH-1), Phagwara-144411, Punjab, India.

ⁱAmity Institute of Pharmacy, Amity University, Sec. 125, Noida 201303, Uttar Pradesh, India

^jDepartment of Biochemistry, Central University of Haryana, Jant-Pali, Mahendergarh District-123031, Haryana, India.

^kNanobiomedicine Lab, Department of Pharmacology, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences, Chennai-600077, Tamil Nadu, India.

^lSchool of Pharmacy and Pharmaceutical Sciences, Ulster University, Coleraine, County Londonderry, BT52 1SA, Northern Ireland, United Kingdom.

^mInovative Medicines Group, Institute of Health & Biomedical Innovation, Queensland University of Technology, Kelvin Grove, Brisbane, Queensland, Australia 4059.

ⁿSchool of Life Sciences, University of Technology Sydney, Sydney, NSW 2007, Australia.

^oDepartment of Life Sciences, School of Pharmacy, International Medical University, Bukit Jalil, Kuala Lumpur 57000, Malaysia.

***Corresponding authors**

Dr. Kamal Dua

Discipline of Pharmacy, Graduate School of Health,
University of Technology Sydney, P.O. Box: 123 Broadway,
NSW 2007, Australia.
Tel: +61 2 95147387; E-mail: kamalpharmacist02@gmail.com

Dr Dinesh Kumar Chellappan

Department of Life Sciences, International Medical University,
Bukit Jalil 57000, Kuala Lumpur, Malaysia
Tel: +60 126361308; E-mail: dineshkumarchellappan.imu@gmail.com, Dinesh_Kumar@imu.edu.my

Abstract

Lung diseases are the leading cause of mortality worldwide. The currently available therapies are not sufficient, leading to the urgent need for new therapies with sustained anti-inflammatory effects. Small/short or silencing interfering RNA (siRNA) has potential therapeutic implications through post-transcriptional downregulation of the target gene expression. siRNA is essential in gene regulation, so is more favorable over other gene therapies due to its small size, high specificity, potency and no or low immune response. In chronic respiratory diseases, local and targeted delivery of siRNA is achieved *via* inhalation. The effectual delivery can be attained by the generation of aerosols *via* inhalers and nebulizers which overcomes anatomical barriers, alveolar macrophage clearance and mucociliary clearance. In this review, we discuss the different siRNA nanocarrier systems for chronic respiratory diseases, for safe and effective delivery. siRNA mediated pro-inflammatory gene or miRNA targeting approach can be a useful approach in combating chronic respiratory inflammatory conditions and thus providing sustained drug delivery, reduced therapeutic dose, and improved patient compliance. This review will be of high relevance to the formulation, biological and translational scientists working in the area of respiratory diseases.

Keywords: siRNA; delivery systems; pulmonary; nanocarriers; RNA interference

1 Introduction

2 Chronic respiratory disease (CRD), in particular, asthma and chronic obstructive pulmonary disease
3 (COPD), are amongst the leading causes of mortality and morbidity that also exert huge health and
4 economic burden globally ("Global, regional, and national deaths, prevalence, disability-adjusted life
5 years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990-2015: a
6 systematic analysis for the Global Burden of Disease Study 2015," 2017). Asthma is a complex and
7 heterogeneous chronic inflammatory disorder, primarily affecting the airways. The disease is largely
8 'allergic' in nature and is characterized by the upregulation of genes that lead to multiple inflammatory
9 cascades implicated in asthma (Moheimani et al., 2016). Current asthma therapies focus on reduction of
10 symptoms and limit exacerbations during the course of the disease. However, the proportion of asthmatics
11 with the uncontrolled disease remain significantly high and utilizes the majority of healthcare expenses
12 globally (P. M. Hansbro *et al.*, 2017; Peters, Ferguson, Deniz, & Reisner, 2006). Despite the increasing
13 classification/categorization of asthmatics into various endotypes/phenotypes (based on a number of
14 molecular biomarkers, clinical presentation and responsiveness to common therapies)(Fajt & Wenzel,
15 2015), the key pathological feature of asthma is involvement of multiple inflammatory mediators that are
16 often regulated by 'key' genes/proteins, which are also referred as master transcription regulators(Sel,
17 Henke, Dietrich, Herz, & Renz, 2006). Thus, targeting these key pro-inflammatory genes would
18 potentially improve the overall disease outcomes(Sel et al., 2006).

19 COPD is another complex and multi-factorial respiratory disease that is caused primarily due to chronic
20 exposure to cigarette smoking(Rennard & Vestbo, 2006). In addition, exposures to biomass smoke, air
21 pollution and a variety of occupational exposures to chemical dust and fumes also play an important role
22 in the onset and progression of COPD(KC, Shukla, Gautam, Hansbro, & O'Toole, 2018). COPD
23 constitutes multiple inflammatory pathways, that are induced by chronic exposure to irritants and, in
24 addition, due to an imbalance between the oxidase/anti-oxidant ratio and/or proteases/anti-proteases ratio
25 (Peter J. Barnes, 2016). The course of the disease is further complicated by frequent acute exacerbations,
26 which are described as worsening of disease symptoms that require a change of daily medications and
27 often requiring hospitalizations(Rodriguez-Roisin, 2000). Acute exacerbations are primarily caused by
28 infections (bacterial and/or viral) which dramatically increase the risk of mortality and morbidity amongst
29 COPD patients(Sapey & Stockley, 2006). In addition, the lack of effective therapies to limit the onset
30 and/or progression of COPD further aids in increased global morbidity and mortality, especially in aging
31 populations.

1 Idiopathic pulmonary fibrosis (IPF) is a life-threatening lung disease that involves the progressive loss of
2 lung function along with clinically significant thickness/stiffness of lung tissues, generally accompanied
3 by tissue scarring. The 3- and 5-year mortality rates in IPF patients are almost 50%(Meltzer & Noble,
4 2008). Integrative network analysis revealed a total of 27 genes (CHIT1, CXCL14, LPPR4, etc.) and 22
5 miRNAs (in particular, miR-409-5p and has-miR-376c) that are associated with the disease
6 development(L. Wang, Huang, Zhang, Chen, & Zhao, 2018). Targeting these and similar nucleic acids
7 that are crucial in the onset/progression of IPF would certainly lead to the development of novel therapies
8 for treating IPF and enhancing the quality of life in IPF patients.

9 Lung cancer is one of the leading causes of cancer-related mortality globally(Bray et al., 2018). A number
10 of genetic mutations have been linked to the pathogenesis of two major types of lung cancer, including
11 non-small-cell lung cancer (e.g., KRAS, EGFR, ALK, MET exon 14, BRAF, PIK3CA, ROS1, HER2, and
12 RET)("Comprehensive molecular profiling of lung adenocarcinoma," 2014) and squamous cell carcinoma
13 (PIK3CA, PTEN and amplification of FGFR1)(Rosell & Karachaliou, 2016). Importantly, circulating
14 cell-free nucleic acids are now being investigated/utilized in both diagnosis and personalised treatments
15 for lung cancers, which have been reviewed in detail by Sorber *et al.*(Sorber *et al.*, 2017). Given the
16 immense burden and limited treatment options for lung cancer, nucleic acid-based therapies may present a
17 novel therapeutic front to develop personalised therapies.

18 Respiratory infections, especially with viruses and bacteria, further complicate the course of chronic
19 respiratory diseases. In particular, viral pathogens, such as rhinovirus, influenza virus and respiratory
20 syncytial virus, are often implicated in the exacerbations of respiratory diseases, including asthma, COPD
21 and IPF(Wark, Tooze, Powell, & Parsons, 2013). In addition, bacterial pathogens (e.g., *Haemophilus*
22 *influenzae*, *Streptococcus pneumoniae*, and *Moraxella catarrhalis*), either on its own or as a secondary
23 pathogen following prior viral infections seem to be more detrimental to the lung health and poorer
24 quality of life(Didierlaurent, Goulding, & Hussell, 2007; Wark *et al.*, 2013). A better understanding of the
25 interactions between respiratory infections and chronic respiratory diseases (asthma, COPD) is essential
26 for the design of more effective preventive and treatment strategies. For instance, tuberculosis seems to
27 increase the risk of COPD and vice versa (O'Toole, Shukla, & Walters, 2015). Moreover, tuberculosis
28 increases the oxidative burden in the lungs (K. Dua *et al.*, 2018; Shastri *et al.*, 2018), regulated by
29 multiple inflammatory mediators (K. Dua *et al.*, 2019), that could be targeted with the novel, nucleic acid-
30 based therapies for reducing the treatment regimens.

31 Despite the huge burden of major respiratory disease, there is a lack of effective treatments that could
32 limit the disease onset and/or disease progression(K Dua, DK Chellappan, *et al.*, 2018; K Dua, Gupta,

Chellappan, Shukla, & Hansbro, 2018; Kamal Dua, Vamshi Krishna Rapalli, *et al.*, 2018; Kamal Dua, Shukla, de Jesus Andreoli Pinto, & Hansbro, 2017; Rapalli *et al.*, 2018). Thus, extensive research into potentially novel therapeutics is urgently required. Currently, the potential of novel classes of therapeutics, which are nucleic acid-based, for treating lung diseases is under intense investigation due to their crucial roles in regulating gene expression(Kamal Dua, Terezinha de Jesus Andreoli Pinto, *et al.*, 2018; Kamal Dua, Hansbro, Foster, & Hansbro, 2017; Kamal Dua, Shukla, Tekade, & Hansbro, 2017). For instance, antisense oligonucleotides, which are single-strand DNAs or RNAs that selectively bind to complementary mRNAs and modulate their functions, could potentially result in up-/down-regulation of particular genes, thus aiding in limiting the disease progression (Bennett, Baker, Pham, Swayze, & Geary, 2017). Similarly, small interference RNAs (siRNAs) are double-strand RNA molecules (~21-23 base pairs in length) capable to ‘silence’ specific genes of a known sequence responsible for genome stability(J. K. Lam, Chow, Zhang, & Leung, 2015). siRNA works at two stages: post-transcriptional gene silencing (PTGS) resulting in direct sequence-specific cleavage causing repression of translation and degradation resulting in transcriptional gene silencing (TGS). siRNA is found associated with effector associations which are recognized by RNA-induced silencing complexes (RISCs)(Martinez, Patkaniowska, Urlaub, Lührmann, & Tuschl, 2002). Further, this siRNA uses its full sequence for the recognition of the target sequence and cleaves the target mRNA (Jie Wang, Lu, Wientjes, & Au, 2010). Micro RNAs (miRNAs) are single-stranded, endogenous non-coding RNA molecules (~18–24 base pairs long). The miRNAs act as important regulators for a variety of immunological and cellular pathways (Awasthi, Madan, Malipeddi, Dua, & Kulkarni, 2019; Hansbro & Dua, 2018; D. D. Nguyen & Chang, 2017).

Notably, siRNAs have been previously speculated to be an effective nucleic acid-based therapy in asthma(M. Choi, J. Gu, M. Lee, & T. Rhim, 2017; Luo *et al.*, 2012), COPD(Luo *et al.*, 2012), IPF(C. N. D'Alessandro-Gabazza *et al.*, 2012), lung cancer(Merkel, Rubinstein, & Kissel, 2014) and respiratory infections(Merkel *et al.*, 2014). We have attempted to primarily focus on the potential drug delivery strategies to enhance the efficacy of siRNAs in treating chronic respiratory diseases, because siRNA can result in multiple gene mutations (oncogenes and tumor suppressor genes) and acts as efficient and promising cure of disease in comparison to other therapies.

Drug delivery systems for siRNA in pulmonary diseases

siRNA has an enormous prospective for the treatment or avoidance of different lung infections. When the RNA particles effectively enter the target, they may obstruct the expression of specific gene series through RNA interference (RNAi) to produce remedial impacts(Amreddy *et al.*, 2018; Ayatollahi *et al.*,

2017). The greatest deterrent to translating siRNA treatment from the research facilities into the clinics is delivery. A perfect delivery operator ought to shield the siRNA from enzymatic debasement, encourage cell take-up and advance endosomal escape inside the cells, with irrelevant poisonous effect(V. Capel *et al.*, 2018; Jin *et al.*, 2018). Pulmonary targeting could be accomplished by fundamental delivery or lung delivery. The latter administrative route could possibly upgrade siRNA maintenance in the lungs and lessen fundamental poisonous impacts. The delivery design should be planned cautiously so as to boost the deposition of siRNA to the unhealthy region of the aviation routes. In majority of the lung siRNA treatment studies *in vivo*, siRNA was conveyed either intranasally or intratracheally (Figure: 1) (I. d'Angelo *et al.*, 2018; Ding *et al.*, 2018; He *et al.*, 2018).

Limited studies have been reported on siRNA formulations *via* inhalation, although, it is expected for potential future prospects. Following are the delivery systems of siRNA in pulmonary diseases:

1. Lipid-based delivery vectors

The lipid-based delivery system is generally used to deliver siRNA *in vitro* and *in vivo*. Ordinarily, cationic lipids or liposomes utilizes negatively charged siRNA through electrostatic forces and are known as lipoplexes. Several commercially available agents are lipid-based, some of which are for *in vivo* lung delivery, for example, DharmFECT, **lipofectamine**, and Oligofectamine. The main difficulties of utilizing lipid-based conveyance vectors in the clinical setup are their harmfulness and their non-specific activation of inflammatory cytokines and interferon reactions. Lipid-based delivery vectors can be classified into five types of molecules: Cationic lipoplexes and liposomes; PEGylated lipids; Neutral lipids; Lipids particles and Lipid-like molecules(Kaczmarek *et al.*, 2018; Liu *et al.*, 2019; Mokhtarieh, Lee, Kim, & Lee, 2018; O. S. Muddineti, A. Shah, S. V. K. Rompicharla, B. Ghosh, & S. Biswas, 2018).

2. Polymer-based delivery vectors

The polymer-based delivery vectors have an adaptable nature, which enables their physicochemical qualities to change effectively to accommodate their purpose. Also, it has been recommended that polymers usually do not induce a strong immune response (Ni *et al.*, 2018; Otsuka *et al.*, 2017). Polymer-based vectors are separated into two classes: polycations and polymeric nanoparticles. Engineered polycations, for example, polyamidoamine dendrimers and polyethyleneimine and natural polycations, for example, chitosan are utilized for conveying DNA for a long time(Y. Qiu *et al.*, 2017; Sasaki & Guo, 2018).

3. Peptide-based delivery vectors

1 Since the discovery of TAT protein from HIV-1, which assists in the uptake of the virus in the cell, a
2 range of cell-penetrating peptides (CPPs) have been identified. CPPs are most commonly utilized as
3 vehicles of restorative macromolecules. This technique can thus be used to deliver siRNA (Veilleux *et al.*,
4 2018; P. Y. Xu *et al.*, 2018). CPPs and subsidiaries have been explored for siRNA conveyance which
5 include MPG, TAT, transportan, penetratin, CADY and LAH4. The peptides are either covalently
6 connected to siRNA through disulfide bond development or electrostatically in a non-covalent way (Yuan
7 *et al.*, 2017; D. Zhang *et al.*, 2018).

8 **Immune response in siRNA-mediated drug delivery to lungs**

9 Airway inflammation is the major contributing factor in the pathogenesis of lung diseases, which
10 corresponds to the degree of symptoms, airways hyper-responsiveness, and obstruction. Therefore,
11 siRNA mediated delivery vehicle, as well as the products of secondary RNAi, stimulate the immune
12 system by enhancing the expression or suppressing pro-inflammatory cytokines, interferons, and toll-like
13 receptors *via* signaling pathways (Sioud, 2015). This is evident from an *in vivo* study for intratracheal
14 administration of penetratin (peptide) conjugated siRNA capable of penetrating cytosol, activating several
15 biochemical pathways and elevates the expression of TNF- α , IL-12, and p40. It also activates the innate
16 immune system to target p38 MAP kinase (Moschos *et al.*, 2007).

17 The approach of siRNA based therapeutics against tuberculosis infections directs the gene expression
18 modulation of the host rather than the bacilli (Man, Chow, Casettari, Gonzalez-Juarrero, & Lam, 2016).
19 The genes that regulate autophagy in the host can be targets for *M. tuberculosis* infection such as Rap22a,
20 Bfl-1/A1, Ras homologue enriched in brain (Rheb) and UV radiation resistance associated genes
21 (UVRAG) (Kathania, Raje, Raje, Dutta, & Majumdar, 2011; Kim *et al.*, 2015; Roberts, Chua, Kyei, &
22 Deretic, 2006; Jinli Wang *et al.*, 2013). The suppression of these genes can hinder bacterial growth and
23 proliferation. Another strategy is immunosuppression to inhibit bacterial growth. TNF- α and IFN γ are
24 essential for macrophage activation to initiate granuloma formation for the proliferation of *M.*
25 *tuberculosis* (Ramakrishnan, 2012; Silva Miranda, Breiman, Allain, Deknuydt, & Altare, 2012).
26 Chemokine and Lymphotoxin/XCL-1 activate CD8⁺T cells during severe tuberculosis infection. In
27 addition to this, XCL-1 regulates IFN- γ and CD4⁺T cells to maintain granuloma. When siRNA targeting
28 XCL-1 is delivered in TB infected mice, the expression of XCL-1 is greatly reduced, therefore,
29 decreasing T lymphocytes, IFN γ , and granuloma (Rosas-Taraco, Higgins, Sanchez-Campillo, *et al.*,
30 2009).

31 The intranasal siRNA mediated delivery of SOCS3 in chronic asthma mouse model had been reported to
32 improve eosinophil count and airways hyper-responsiveness. This has also been correlated to improved

mucus secretion, collagen reduction, and airway remodeling. siRNA mediated SOC3 reduces RhoA/Rho kinase signaling pathway, thereby, regulating inflammation, bronchial smooth muscle contractions and upregulating IL-13, IL-4 *via* RhoA protein and STAT6 activation. This reduces cytokine expression and airways hyper-responsiveness. It is known that IL-17 and IL-23 are responsible for neutrophilic and eosinophilic inflammation in mice (Figure.2)(Molet *et al.*, 2001; Wakashin *et al.*, 2008). Thus, siRNA mediated silencing of SOC3 regulates IL-17 expression(Staff, 2014).

siRNA drug delivery in asthma/allergic airway diseases

The combination therapy for asthma includes the use of inhaled corticosteroids, β_2 adrenergic receptor agonists, injected immunoglobulin E antibodies and quick-relief medications for effective control of asthma (Peter J Barnes, 2004). However, conventional inhaled corticosteroid therapies do not work in severe asthmatic patients (refractory asthma) and may cause adverse effects after long-term treatment (Kamal Dua, Hansbro, & Hansbro, 2017; Philip M Hansbro *et al.*, 2017; Mealey, Kenyon, Avdalovic, & Louie, 2007). Therefore, more attention is needed in asthma research focused on the development of target-specific therapeutics. It has been previously reported that proteins such as chemokines(Rosenwasser, Zimmermann, Hershey, Foster, & Rothenberg, 2003) and cytokines (Peter J Barnes, 2008) are involved in the asthmatic-inflammatory processes. Leukotrienes are also involved in the pathogenesis of asthma (Ogawa & Calhoun, 2006).

Omalizumab (Xolair® - a monoclonal antibody against IgE), reslizumab (Cinqair® in the US and Cinqaero® in Europe -a humanized antibody against human interleukin-5) and mepolizumab (Nucala® - blocks interleukin-5) are commercially available potential targeted therapeutics to treat asthma (Catley, Coote, Bari, & Tomlinson, 2011; Corren, 2012). Due to the complex nature of the disease, there is no single medication available for effective control of the asthmatic symptoms. Such targeted therapies may provide a valuable breakthrough in the pathophysiology of asthma and increase the possibility of decreasing asthma burden (M Choi, J Gu, M Lee, & T Rhim, 2017; Cook & Bochner, 2010).

siRNA delivery to the pulmonary system has ameliorated the therapeutic benefits of RNA interference (RNAi) for asthma (H.-Y. Huang & Chiang, 2009). RNAi has been reported to be effective in blocking functions of molecular targets of asthma. siRNAs (21-23 bp in length) are involved in the sequence-specific degradation of messenger RNA (mRNA) and decrease the expression of the corresponding proteins (Agrawal *et al.*, 2003; Xie & Merkel, 2015). Prolonged lung retention of siRNA administered *via* the pulmonary route can reduce systemic side effects and improve the therapeutic benefits in the

1 treatment of asthma (Deng *et al.*, 2014; Rettig & Behlke, 2012; Xie & Merkel, 2015). Intranasal
2 administration of siRNA formulation with or without a transfection agent has been reported to inhibit
3 replication of the respiratory syncytial virus in mice models (Whitehead, Langer, & Anderson, 2009).

4 Xie *et al.*, (2016) developed a siRNA based delivery system, transferring polyethylenimine (Tf-PEI), to
5 target specific delivery of siRNAs and activated T cells in the lung. The optimized polyplexes possess
6 ideal physical properties such as zeta-potential, size, distribution and siRNA condensation efficiency.
7 Formulated polyplexes showed significant enhancement in cellular uptake and gene knockdown.
8 Biodistribution studies in murine asthmatic model confirmed effective delivery of siRNA to the activated
9 T cells (Xie *et al.*, 2016). Based on the literature evidence, Xie and Merkel summarized that chemokines,
10 cytokines, tyrosine kinases, transcription factors, and co-stimulatory factors are the target of siRNA-
11 mediated asthma treatment. Further, the authors proposed potential applications of targeted siRNA
12 delivery to macrophages, T cells, and dendritic cells in asthma therapy (Xie & Merkel, 2015).

13 Wang *et al.*, 2008 reported imiquimod cream-chitosan nanoparticle system containing siRNA green
14 indicator (siGLO) or siRNA for natriuretic peptide receptor (siNPRA). The formulation was applied to
15 the mice skin. Fluorescence microscopy confirmed the delivery of SiGLO to the lungs *via* transdermal
16 route. OVA-sensitized asthmatic BALB/c mice model treated with imiquimodcream-siNPRA chitosan
17 nanoparticles showed significant (< 0.05) decrease in airway hyper-responsiveness, lung histopathology,
18 eosinophilia and pro-inflammatory cytokines IL-4 and IL-5 (X. Wang *et al.*, 2008).

19 Packaging RNAs (pRNAs), a component of the bacteriophage phi29-packaging motor, have been used to
20 deliver signal transducer and activator of transcription (STAT5b) siRNA to asthmatic spleen
21 lymphocytes. Reverse transcription-polymerase chain reaction (RT-PCR) study showed that the STAT5b
22 gene mRNA expression was effectively inhibited by the pRNA dimer. It is suggested that pRNA dimer
23 carrying aptamer (ligand to interact with receptors) and siRNA can deliver functional siRNA to cells (C.
24 Qiu, Peng, Shi, & Zhang, 2012). The first report on lung alveolar epithelial A549 cell targeting by
25 siRNA-generation four, amine-terminated poly (amidoamine) dendrimerdendriplexes and about 40% gene
26 silencing *via* siRNA exposed to the propellant used in oral inhalation devices was given by Conti *et al.*,
27 (Conti, Brewer, Grashik, Avasarala, & da Rocha, 2014).

28 Choi *et al.*, developed dexamethasone and vitamin D binding protein (VDBP) siRNA combination
29 therapeutic system for the treatment of asthma. The results showed that the dexamethasone-conjugated
30 polyethyleneimine/ vitamin D binding protein (DEXA-PEI/VDBP) siRNA, reduced goblet cell
31 hyperplasia, ovalbumin sensitization, challenge-induced enhancement of airway inflammation, and

expressions of interleukin-4 (IL-13), interleukin-4 (IL-4) and eosinophil mobilizing chemokine (CCL11) (M Choi *et al.*, 2017). Wu *et al.*, investigated the ability of c-kit silenced siRNA to decrease the inflammation caused by allergic asthma using asthmatic mouse model treated with intranasal anti-c-kit siRNA. The result showed that siRNA significantly inhibited c-kit gene expression and decreased airway mucus secretion. The production of stem cell factor, IL-4, and IL-5 declined significantly by c-kit siRNA. However, no effect was recorded on interferon- γ (IFN- γ) generation (Wu *et al.*, 2014).

Ambient particulate matters (PMs) are the major causative agents for asthma and chronic obstructive pulmonary disease, by increasing mucus hypersecretion and inflammation. Wang *et al.*, used ambient particulate matter (PM)-exposed human bronchial epithelial cells (HBEC) to determine the function of Amphiregulin (AREG), a ligand for epidermal growth factor receptor (EGFR), in PM-induced inflammation and mucus hypersecretion. The AREG-siRNA significantly suppressed the PM-induced inflammation and mucus hypersecretion. This also suppressed the activation of the EGFR-AKT/ERK pathway (Jian Wang, Zhu, Wang, Chen, & Song, 2019).

siRNA drug delivery in lung cancer

Lung cancer is the leading cause of death all over the world (Siegel, Naishadham, & Jemal, 2012). The widely available therapies include chemotherapy, radiotherapy, and surgery, while, NSCLC treatment is majorly dependent on chemotherapy. Conventionally, chemotherapy has been practiced by intravenous (IV) administration. However, IV administration has side effects like bioavailability of the drug throughout the body *via* the bloodstream, which affects both malignant as well as healthy cells, the death of healthy cells causes adverse side effects like hair loss, fatigue and infections (Gandhi *et al.*, 2018; Moding, Kastan, & Kirsch, 2013). Dobashi *et al.*, reported that the Akt/mTOR pathway is abnormally activated in NSCLC (Dobashi, Watanabe, Miwa, Suzuki, & Koyama, 2011). Several tumor suppressors have been found to be mutated in NSCLC, proposing the function of mTOR pathway. Furthermore, the use of mTOR inhibitors is responsible for adverse side effects on healthy cells. Several molecular targeting drugs such as gefitinib cause inhibition of phosphorylation and tyrosine kinase activity of intracellular ATP binding domain of EGFR via competitive inhibition, erlotinib also inhibits tyrosine kinase and bevacizumab inhibits angiogenesis (Y.K. Oh & Park, 2009; Tiseo, Bartolotti, Gelsomino, & Bordi, 2010). Due to numerous side effects, there is an emerging need for therapeutics for the treatment of lung cancer especially NSCLC (Sadowski, Kotulska, & Jóźwiak, 2016).

Taratula and his co-workers developed nanostructured lipid-based carriers (NLCs) for simultaneous delivery of anticancer drug and siRNA specifically for lung cancer. The drug encapsulated nanocarriers

1 induce cell death, whereas, siRNA suppresses multi-drug resistance. Two types of siRNA were delivered
2 to improve efficacy; the first siRNA was targeting the MRP1 (Multidrug resistance-associated protein)
3 mRNA which is responsible for the suppression of main drug efflux transporter. The second siRNA was
4 targeting Bcl2 (B-cell lymphoma) mRNA, which suppresses cellular anti-apoptotic defense. The drug was
5 delivered to lungs by inhalation after encapsulation in NLCs. Inhalation leads to high accumulation of
6 nanocarriers in lungs whereas, intravenous injection led to major accumulation in liver, spleen, and
7 kidney compared to lungs. The developed NLC formulation effectively delivered the drug and siRNA
8 into cancer cells. This induced cell death of lung tumor cells by targeted gene silencing (Taratula,
9 Kuzmov, Shah, Garbuzenko, & Minko, 2013).

10 Xu *et al.*, studied the combined effect of doxorubicin and Bcl2 siRNA using polyethyleneimine as a
11 carrier for pulmonary delivery. Confocal laser scanning microscopy and flow cytometry exhibited a high
12 cellular uptake of drug and siRNA in B16F10 cell lines. Real-time polymerase chain reaction (RT-PCR)
13 observations had high gene silencing, where, 70% of Bcl2 mRNA were bashed down. The combination
14 has increased cell apoptosis and cell proliferation inhibition in B1F10 cells. The *in-vivo* studies in mice
15 showed high accumulation of doxorubicin and siRNA in metastatic lung cancer upon pulmonary delivery
16 (C.-N. Xu *et al.*, 2017).

17 Hybrid lipid-polymer nanoparticles comprising of dipalmitoylphosphatidylcholine and poly(lactic-co-
18 glycolic) acid were developed to encapsulate siRNA. siGENOMESMART pool siRNA was used to check
19 their fate on the human epithelial airway barrier, which acts against α and β subunits of the sodium trans
20 epithelial channel. The developed nanoparticles exhibited ~150nm hydrodynamic diameter, with -25mV
21 zeta potential. *In-vitro* aerosolization studies were performed on a triple cell co-culture model which
22 mimic human epithelial airway barrier. There were no changes in nanoparticulate structure by
23 transmission electron microscopic imaging after nebulization. Nanoparticles were internalized in
24 epithelial cells and there were no cytotoxic effects or acute inflammation towards cell components. *In-*
25 *vitro* inhibition of sodium trans-epithelial channel protein expression was evaluated in A549 cell lines,
26 which confessed prolonged inhibition (Ivana d'Angelo *et al.*, 2018).

27 Capel *et al.*, reported delivery of siRNA using water-soluble piperazine substituted chitosan derivatives
28 for efficient delivery by inhalation. The piperazine derivative chitosan is water-soluble at physiological
29 pH, and forms nano-complexes with siRNA up to the size of 300nm at a relatively low polymer to siRNA
30 ratio (5:1). Glyceraldehyde-3-phosphate dehydrogenase (GADPH) targeting siRNA was complexed with
31 siRNA, *in-vitro* studies performed on lung epithelial cells revealed chitosan and siRNA complexes
32 exhibited silencing of a gene from 40-80%. There was no effect of the aerosolization by PenCentury™

1 microsprayer device on particle size and integrity. *In-vivo* studies performed to determine the potential of
2 piperazine chitosan complex in subcutaneous bioluminescent tumor A540-lucxenograft model. There was
3 a significant reduction of the tumor with absence of adverse effects. The modified chitosan siRNA
4 complexes were found to be safe and had the potential to deliver by inhalation therapy (Victoria Capel *et*
5 *al.*, 2018).

6 Ihara *et.al.*, developed dry powdered chitosan siRNA complexes, and its gene silencing efficiency was
7 quantified histologically after intratracheal administration into murine lungs. EGFP-siRNA chitosan
8 complex efficacy was studied in EGFP transgenic mice and mice carrying metastatic lung cancer of
9 Lewis lung carcinoma. Transgenic mice are divided into three groups, where, one group was treated with
10 targeting siRNA, one group treated with non-targeting siRNA and one was left without any treatment.
11 The fluorescence in bronchus, bronchioles, alveolar walls of the group treated with targeting siRNA is
12 reduced to a large extent in comparison to mice group treated with non-targeting siRNA and group
13 without treatment. Similarly, the same results were observed in the metastatic lung tumor consisting of
14 mice groups. The results conclude a high extent of gene silencing in proximal airways compared to
15 peripheral lung tissues. The study proves pulmonary delivery of siRNA is a predominant approach to
16 target gene expression in respiratory disorders involving airways, parenchyma and lung cancers (Ihara *et*
17 *al.*, 2015).

18 A dry powder siRNA based formulation was developed targeting vascular endothelial growth factors
19 (VEGF), which inhibits lung tumor growth in mice. *In-vivo* studies were performed on mice with
20 metastatic lung cancer, which were induced by B16F10 melanoma cells or Lewis lung carcinoma cells.
21 VEGF siRNA efficiency in gene suppression was evaluated by treating B16F10 and Lewis lung
22 carcinoma cell lines. There was reduced VEGF protein and mRNA pertaining to VEGF. *In-vivo* studies
23 were performed in tumor-bearing mice, where, chitosan was used to deliver siRNA. Prior to delivery of
24 siRNA, VEGF levels were measured in bronchoalveolar lavage fluid (BALF) collected from mice bearing
25 a tumor. The results showed reduced VEGF concentrations in BALF after single intratracheal
26 administration of dry siRNA powder. Repeated intratracheal administration reduced tumor growth in the
27 lungs. An *in-vitro* inhalation performance study was performed with jethaler single, which exhibited a
28 low-pressure drop. These results suggest chitosan dry powder of siRNA is a novel strategy for lung
29 cancer-specific and high gene silencing effect (Miwata *et al.*, 2018).

30 Bohr and his co-workers developed phosphorus-based dendrimers for delivery of siRNA. Pyrrolidinium
31 and morpholinium were selected as protonated amino groups for better compatibility. The dendriplexes
32 form strong complexes with siRNA targeting tumor necrosis factor- α (TNF- α). The *in-vitro* studies

revealed high cellular uptake and *in-vitro* silencing efficiency of TNF- α in RAW264.7 lipopolysaccharide activated macrophage cell line with pyrrolidinium dendriplexes. The improved efficiency of pyrrolidinium complexes is expected due to high pKa value which improves stronger siRNA complex formation. Nasal administration of complexed dendriplexes has higher efficacy towards lung injury in comparison to non-complexed siRNA (Table 1) (Bohr *et al.*, 2017).

The RNAi mechanism and siRNA provides the potential for designing therapeutics for the treatment of disease like cancer (Leung & Whittaker, 2005). siRNA has intrinsic efficacy as it utilizes the endogenous RNAi pathway, reduces the expression of disease-linked genes and can be used for any gene with its complementary sequence (Vogelstein & Kinzler, 2004). Several genes, their mutations, and pathways have been found to be associated in different cancers, thus it is evident that siRNA can provide therapeutic effect in cancers. siRNA mediated silencing of cancer associated proteins causes a remarkable apoptotic effect (Pai *et al.*, 2006). The major barriers for effective delivery of siRNA to the lungs are complex branching of lungs associated with biomechanical barriers including mucus over the airways and the airways cell membrane. Gene silencing will be achieved only when siRNA delivered is stable, is of good concentration, penetrates the cell and reaches the cytoplasm (Durcan, Murphy, & Cryan, 2008). Zhang *et al.*, reported siRNA counters K-RAS mutants and observed the anti-cancer effect by reducing K-RAS in lung cancer cell line. Additionally, adenovirus-mediated siRNA precisely targets RAS and acts as a drug for lung cancer treatment (Z. Zhang, Jiang, Yang, & Wang, 2006). Han *et al.*, proposed the use of p65 siRNA for anti-tumor effect by blocking PI3-kinase and NF κ B (Han & Roman, 2006).

Presently, in clinical trials, siRNA was locally administered to the target site to bypass the systemic delivery but the systemic route is required for the treatment of cancers and other diseases. For an *in vivo* delivery system, it ought to be biocompatible, non-immunogenic and biodegradable. The siRNA should be effectively delivered to the target site and must be protected from the action of serum nucleases. The system must evade from immediate hepatic or renal clearance and foster endosomal siRNA release into the cytoplasm for endogenous RISC interaction (Juliano, Alam, Dixit, & Kang, 2008). Development and validation of different strategies of siRNA delivery are underway. Zhang *et al.*, studies the *in vitro* delivery of encapsulated human double-minute gene 2-specific siRNA in arginine octamer surface-modified liposomes. The complex was reported to be stable for 24 hours in blood and had potentially good transfection in different lung cancer cell lines (C. Zhang *et al.*, 2006). Another study based on siRNA against human survivin was coated with cationic liposomes containing DOTAP and cholesterol (1:1 molar ratio) which resulted in LPD (liposome-polycation-DNA) nanoparticles. Further, LPD was PEGylated for ligand targeting and steric stabilization for selective delivery to the lungs. Further analysis suggested that PEGylated LPD has an anti-cancer effect via surviving downregulation (Li & Huang,

2006). In the *in vivo* mouse model, LPD nanoparticle encapsulating siRNA for epidermal growth factor showed anti-tumor activity in combination with cisplatin on intravenous injection (Li, Chen, Hackett, & Huang, 2008). Biodegradable cationic poly (amino-ether) (mPAE) were accessed as a carrier for mTOR siRNA for lower toxic effect, the prohibition against nuclease degradation and inhibition of cancer cell proliferation (Gandhi *et al.*, 2018). Chono *et al.*, reported on the immunotoxicity and organ defects of siRNA-LPD nanoparticle administration intravenously (Chono, Li, Conwell, & Huang, 2008). Cationic immune-liposomes conjugated with anti-transferrin receptor single chain antibody fragment with the fluorescent label have been studied for systemic delivery for lung cancer metastasis. It was observed that the labeled siRNA was distributed in lung metastasis rather than liver (Pirollo *et al.*, 2006). Cationic single-walled carbon nanotubes having siRNA for telomerase reverse transcriptase are being currently studied as *in vitro* lung cancer models. *In vitro* internalization of siRNA suppresses target gene expression. Furthermore, this model has also been reported for mice model for subcutaneous Lewis lung tumors (Zhuohan Zhang *et al.*, 2006). Xu and his colleagues developed a pH-sensitive nanoparticulate system for co-delivery of doxorubicin and survivin siRNA. Doxorubicin was conjugated with polyethylenimine by a pH-sensitive hydrazine bond using 3-maleimidopropionic acid hydrazide. The formed polyethyleneimine-doxorubicin -3- maleimidopropionic acid hydrazides are cationic in nature, and form complexes with anionic survivin siRNA with electrostatic interactions. On pulmonary delivery of these complexes in B16F10 tumor-bearing mice, resulted in the high accumulation of doxorubicin and siRNA in lungs. There was limited accumulation in case of normal lung tissues which indicates targeted drug delivery. The nanoparticulate system has improved anti-tumor efficacy in comparison to individual delivery of doxorubicin or surviving siRNA (C. Xu, Tian, Wang, Wang, & Chen, 2016). A similar study was performed using Bcl2 siRNA with pH-sensitive polyethyleneimine hydrazine doxorubicin complex. The *in-vitro* and *in-vivo* studies showed pH sensitive complex nanoparticles improved anti-tumor efficacy by pulmonary administration. There were high deposition and prolonged retention time in the lungs by pulmonary administration (C. Xu *et al.*, 2015).

Self-assembled cholesterol conjugated chitosan nanoparticles were used to deliver curcumin and siRNA concurrently to achieve a synergistic effect against the cancer cells. Curcumin and siRNA were internalized by clathrin-dependent endocytosis in a time-dependent manner. This method was successful for A549 human lung carcinoma cell line for the co-delivery of siRNA and hydrophobic drug (Omkara Swami Muddineti, Aashma Shah, Sri Vishnu Kiran Rompicharla, Balaram Ghosh, & Swati Biswas, 2018). Similarly, cationic polyethyleneimine-polylactic acid (PEI-PLA) was synthesized for systemic delivery of paclitaxel and siRNA for the knockdown of survivin gene for lung carcinoma. Upon nanoparticle uptake by the A549 cells, they turn electrically neutral due to lower endosomal pH. These

nanoparticles have pH-responsive property as pH 5.5 leads to drug release while pH 7.4 for cellular uptake. This study proved tumor growth inhibition along with surviving mRNA knockdown. Such co-delivery systems provide large surface area for high drug loading, time-dependent drug release which allows extended exposure to tumor treatment, passive targeting, lower cytotoxicity increased the proliferative effect and anti-cancer activity (Jin *et al.*, 2018) as depicted in the Figure:3.

Inhalation therapy can be potentially used for siRNA delivery due to high gene silencing effect and sequence specificity. *In vivo* study on mice with Lewis lung carcinoma reveals the effect of intratracheal vascular endothelial growth factor (VEGF) siRNA dry powder delivery downregulates the VEGF levels in both tumor tissue as well as broncho-alveolar lavage and reduces the metastatic loci in lungs (Miwata *et al.*, 2018). The aerosol composed of PEA and siRNA for Akt1 was administered in mice with urethane-induced inhaled lung cancer. The use of aerosol for 4 weeks reduces Akt1 levels and inhibits the tumor progression (C.-X. Xu *et al.*, 2008).

Clinical trial studies for siRNA drug delivery in pulmonary diseases

A drug called ExcellairTM was developed by ZaBeCor Pharmaceuticals for the treatment of asthma. The drug targets mRNA of spleen tyrosine kinase (Syk) which is responsible for activation of several pro-inflammatory transcription factors. In the Phase I of the study, patients received siRNA ExcellairTM via inhalation for 21 days (Watts & Corey, 2010). The drug did not cause any side effects to the asthma patients and almost 75% of the patients reported improved breathing and reduction in the use of inhalers while placebo patients showed no improvement. Further, in 2009, ExcellairTM entered Phase II of clinical trials but in 2015 it was discontinued as Syk can act as both suppressor and promoter of cell growth (Krisenko & Geahlen, 2015).

Alnylan Pharmaceuticals developed ALN-RSV01, a siRNA therapeutic to target mRNA of viral protein in respiratory syncytial virus (RSV) (DeVincenzo *et al.*, 2010). ALN-RSV01 targets nucleocapsid (N) protein of RSV which is essential for viral replication. In the Phase I of the clinical trial, 100 healthy males of 18 to 45 years age group were accounting to 65 having single and multiple dose of ALN-RSV01 while 36 were placebo. All the volunteers were given the therapeutic doses via nasal spray. No severe adverse effects were observed in different treated groups which lead ALN-RSV01 to enter Phase II (DeVincenzo *et al.*, 2008). The Phase II study comprised of 85 healthy males of 18-45 years of age. All subjects received RSV01 inoculation at day 0, and the ALN-RSV01 treated cohorts received the siRNA intranasal spray at days -1, 0,

+1, +2, and +3. A statistically significant reduction in detected RSV by quantitative culture and real-time PCR was reported for patients receiving 150mg of study drug, the highest dosage tested. Averaged for all treated patients vs. placebo, an acquisition over time effect was observed by either PCR or quantitative culture. Indeed, the strongest effects of treatment with ALN-RSV01 were observed by its prophylactic efficacy. The drug provided an antiviral effect over an 11-day time course, which resulted in reduced infection over time noticeable within 3–4 days after inoculation (DeVincenzo *et al.*, 2008). The safety trial for antiviral activity was conducted for lung transplant patients with RSV. In Phase II b it was showed the safety of ALN-RSV01 treated RSV infections across the broader groups for lung transplant (Alvarez *et al.*, 2009).

Atu-027 is composed of siRNA with lipoplex delivery system represents RNAi mediated suppression of protein kinase N3 (PKN3) in vascular endothelial cells and prevents lung metastasis. Phase Ib trials for the safety of Atu-027 with gemcitabine have been completed (Schultheis *et al.*, 2014).

Future prospects of drug delivery with siRNA in pulmonary diseases

RNA interference plays a key role in the treatment of several disorders. siRNA induces gene silencing by acting on sequence-specific cleavage of complementary mRNA (messenger RNA) and thereby inhibiting protein synthesis. siRNA based therapy was found to be a better strategy over the existing therapeutics, such as drug molecules, monoclonal antibodies and proteins (Fujita, Takeshita, Kuwano, & Ochiya, 2013). Apart from its advantages, administration and delivery of siRNA is the major challenge. siRNA undergoes degradation in the presence of serum nucleases when they are administered directly into the blood. Several strategies are utilized for delivery of siRNA to the target organ (T. Nguyen, Menocal, Harborth, & Fruehauf, 2008). siRNA demonstrated as potential therapeutic agents to treat pulmonary disorders including lung cancer, infectious diseases, airway inflammatory diseases, and cystic fibrosis. Delivery of siRNA directly by pulmonary route has added advantages such as reduced dose, reduced systemic side effects and reduced degradation due to a lower concentration of nuclease enzymes in airways. Pulmonary delivery of siRNA can also be helpful in systemic action, due to the large surface area, thin epithelium and high vascularization in alveoli which favors rapid absorption of siRNA (Fujita *et al.*, 2013). Inhalation is the most preferred and easy mode of non-invasive administration which can be applied for siRNA by liquid aerosol or dry aerosol formulations. There is a need for a high attention regarding stability and biological activity of siRNA at the time of formulation development and delivery. However, the pulmonary delivery of these siRNA is challenging due to mucociliary clearance by ciliated

epithelium cells, mucus, alveolar fluid and macrophages along the airways. Particles inhaled get deposited on ciliated cells and eventually get cleared by cough and swallowed. The mucus secreted, form a thin film thereby restricting diffusion and penetration of siRNA into the cell membrane. The alveolar fluids form a thin film as a pulmonary surfactant (phospholipids and surfactant proteins) obstruct permeation efficiency of lipid-based formulations to some extent and there was no effect in case of polymer-based systems. Macrophages engulf the inhaled particles as part of the defense mechanism. There will be altered conditions like increased mucus secretion, viscosity and ciliary clearance in diseased conditions. siRNA with a negative charge and high molecular weight (13kDa), has also contributed to the poor ability to cross cell membrane even it reaches the target surface area. Viral vectors and non-viral vectors are used for the delivery of siRNA. Viral vectors are found to be more efficient to transfer genetic material into host cells. Despite of its advantages, activation of immune responses after repeated administration may lead to organ failure and chances of serious concerns. Non-viral vectors which include lipids, polymers, inorganic materials and transfection agents (i.e., Lipofectamine, Oligofectamine, TransIT-TKO and DharmaFECT) are more vastly utilized for siRNA delivery. The ideal characteristic feature for a siRNA delivery system should include a) Protection from enzymatic degradation, b) Ability to penetrate cell membrane (facilitate cell uptake) c) It should able to protect from endosomal degradation and induce gene silencing d) Should not affect siRNA activity and specificity e) Non-toxic (Feldmann & Merkel, 2015; J. K.-W. Lam, Liang, & Chan, 2012; Yingshan Qiu, Lam, Leung, & Liang, 2016; Youngren-Ortiz, Gandhi, España-Serrano, & Chougule, 2016, 2017). Several strategies are utilized to overcome stated challenges in the delivery of siRNA some of them are discussed below.

Conclusion

The potential siRNA based therapeutics for lung diseases have to be explored further. The nanoparticle-based inhalable and aerosols have been studied extensively for improved delivery and clinical efficiency. Therefore, *ex vivo* models for inhaled particulate distribution and *in vivo* models for pharmacokinetics/pharmacodynamics are being taken under consideration to understand their distribution, safety, and efficiency of siRNA in the pulmonary system. Different organic and inorganic nanoparticle with varied size, charge, and chemistry have been used as carriers of siRNA or siRNA drug conjugate delivery. Also, upon internalization at the target site siRNA should be able to escape the endosomal mechanism. Furthermore, a nanoparticle with small size have been preferred for longevity at the target site, non-specific interactions and to prevent the off-site toxic effect.

Acknowledgements

PMH is funded by a Fellowship from the National Health Medical Research Council, NHMRC (1079187). We would like to acknowledge NHMRC and the Graduate School of Health (GSH), University of Technology Sydney (UTS), NSW, Australia for the funding.

References

- Agrawal, N., Dasaradhi, P., Mohammed, A., Malhotra, P., Bhatnagar, R. K., & Mukherjee, S. K. (2003). RNA interference: biology, mechanism, and applications. *Microbiology and molecular biology reviews*, 67(4), 657-685.
- Alvarez, R., Elbashir, S., Borland, T., Toudjarska, I., Hadwiger, P., John, M., . . . Kahn, J. (2009). RNA interference-mediated silencing of the respiratory syncytial virus nucleocapsid defines a potent antiviral strategy. *Antimicrobial agents and chemotherapy*, 53(9), 3952-3962.
- Amreddy, N., Babu, A., Panneerselvam, J., Srivastava, A., Muralidharan, R., Chen, A., . . . Ramesh, R. (2018). Chemo-biologic combinatorial drug delivery using folate receptor-targeted dendrimer nanoparticles for lung cancer treatment. *Nanomedicine*, 14(2), 373-384. doi:10.1016/j.nano.2017.11.010
- Asai-Tajiri, Y., Matsumoto, K., Fukuyama, S., Kan-o, K., Nakano, T., Tonai, K., . . . Nakanishi, Y. (2014). Small interfering RNA against CD86 during allergen challenge blocks experimental allergic asthma. *Respiratory research*, 15(1), 132.
- Awasthi, R., Madan, J. R., Malipeddi, H., Dua, K., & Kulkarni, G. T. (2019). Therapeutic strategies for targeting non-coding RNAs with special emphasis on novel delivery systems. *Non-coding RNA Investigation*, 3.
- Ayatollahi, S., Salmasi, Z., Hashemi, M., Askarian, S., Oskuee, R. K., Abnous, K., & Ramezani, M. (2017). Aptamer-targeted delivery of Bcl-xL shRNA using alkyl modified PAMAM dendrimers into lung cancer cells. *Int J Biochem Cell Biol*, 92, 210-217. doi:10.1016/j.biocel.2017.10.005
- Barnes, P. J. (2004). New drugs for asthma. *Nature reviews Drug discovery*, 3(10), 831.
- Barnes, P. J. (2008). The cytokine network in asthma and chronic obstructive pulmonary disease. *The Journal of clinical investigation*, 118(11), 3546-3556.
- Barnes, P. J. (2016). Inflammatory mechanisms in patients with chronic obstructive pulmonary disease. *Journal of Allergy and Clinical Immunology*, 138(1), 16-27. doi:10.1016/j.jaci.2016.05.011
- Bennett, C. F., Baker, B. F., Pham, N., Swayze, E., & Geary, R. S. (2017). Pharmacology of Antisense Drugs. *Annu Rev Pharmacol Toxicol*, 57, 81-105. doi:10.1146/annurev-pharmtox-010716-104846
- Bitko, V., Musiyenko, A., Shulyayeva, O., & Barik, S. (2005). Inhibition of respiratory viruses by nasally administered siRNA. *Nature medicine*, 11(1), 50.
- Bohr, A., Tsapis, N., Andreana, I., Chamarat, A., Foged, C., Delomenie, C., . . . Mignani, S. (2017). Anti-Inflammatory Effect of Anti-TNF- α SiRNA Cationic Phosphorus Dendrimer Nanocomplexes Administered Intranasally in a Murine Acute Lung Injury Model. *Biomacromolecules*, 18(8), 2379-2388.
- Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R. L., Torre, L. A., & Jemal, A. (2018). Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*, 68(6), 394-424. doi:10.3322/caac.21492
- Capel, V., Villasaliu, D., Watts, P., Clarke, P. A., Luxton, D., Grabowska, A. M., . . . Stolnik, S. (2018). Water-soluble substituted chitosan derivatives as technology platform for inhalation delivery of siRNA. *Drug Deliv*, 25(1), 644-653. doi:10.1080/10717544.2018.1440668

- 1 Capel, V., Vllasaliu, D., Watts, P., Clarke, P. A., Luxton, D., Grabowska, A. M., . . . Stolnik, S. (2018). Water-
2 soluble substituted chitosan derivatives as technology platform for inhalation delivery of siRNA.
3 *Drug delivery*, 25(1), 644-653.
- 4 Catley, M. C., Coote, J., Bari, M., & Tomlinson, K. L. (2011). Monoclonal antibodies for the treatment of
5 asthma. *Pharmacology & therapeutics*, 132(3), 333-351.
- 6 Choi, M., Gu, J., Lee, M., & Rhim, T. (2017). A new combination therapy for asthma using dual-function
7 dexamethasone-conjugated polyethylenimine and vitamin D binding protein siRNA. *Gene Ther*,
8 24(11), 727-734. doi:10.1038/gt.2017.83
- 9 Choi, M., Gu, J., Lee, M., & Rhim, T. (2017). A new combination therapy for asthma using dual-function
10 dexamethasone-conjugated polyethylenimine and vitamin D binding protein siRNA. *Gene*
11 *therapy*, 24(11), 727.
- 12 Chono, S., Li, S.-D., Conwell, C. C., & Huang, L. (2008). An efficient and low immunostimulatory
13 nanoparticle formulation for systemic siRNA delivery to the tumor. *Journal of Controlled*
14 *Release*, 131(1), 64-69.
- 15 Chow, M. Y., Qiu, Y., Lo, F. F., Lin, H. H., Chan, H.-K., Kwok, P. C., & Lam, J. K. (2017). Inhaled powder
16 formulation of naked siRNA using spray drying technology with L-leucine as dispersion enhancer.
17 *International journal of pharmaceutics*, 530(1-2), 40-52.
- 18 Comprehensive molecular profiling of lung adenocarcinoma. (2014). *Nature*, 511(7511), 543-550.
19 doi:10.1038/nature13385
- 20 Conde, J., Tian, F., Hernández, Y., Bao, C., Cui, D., Janssen, K.-P., . . . Jesús, M. (2013). In vivo tumor
21 targeting via nanoparticle-mediated therapeutic siRNA coupled to inflammatory response in
22 lung cancer mouse models. *Biomaterials*, 34(31), 7744-7753.
- 23 Conti, D. S., Brewer, D., Grashik, J., Avsarala, S., & da Rocha, S. R. (2014). Poly (amidoamine) dendrimer
24 nanocarriers and their aerosol formulations for siRNA delivery to the lung epithelium. *Molecular*
25 *pharmaceutics*, 11(6), 1808-1822.
- 26 Cook, M. L., & Bochner, B. S. (2010). Update on biological therapeutics for asthma. *World Allergy*
27 *Organization Journal*, 3(6), 188.
- 28 Corren, J. (2012). Inhibition of interleukin-5 for the treatment of eosinophilic diseases. *Discovery*
29 *medicine*, 13(71), 305-312.
- 30 D'Alessandro-Gabazza, C. N., Kobayashi, T., Boveda-Ruiz, D., Takagi, T., Toda, M., Gil-Bernabe, P., . . .
31 Suzuki, N. (2012). Development and preclinical efficacy of novel transforming growth factor- β 1
32 short interfering RNAs for pulmonary fibrosis. *American journal of respiratory cell and molecular*
33 *biology*, 46(3), 397-406.
- 34 D'Alessandro-Gabazza, C. N., Kobayashi, T., Boveda-Ruiz, D., Takagi, T., Toda, M., Gil-Bernabe, P., . . .
35 Gabazza, E. C. (2012). Development and preclinical efficacy of novel transforming growth factor-
36 beta1 short interfering RNAs for pulmonary fibrosis. *Am J Respir Cell Mol Biol*, 46(3), 397-406.
37 doi:10.1165/rcmb.2011-0158OC
- 38 d'Angelo, I., Costabile, G., Durantie, E., Brocca, P., Rondelli, V., Russo, A., . . . Petri-Fink, A. (2018). Hybrid
39 Lipid/Polymer Nanoparticles for Pulmonary Delivery of siRNA: Development and Fate Upon In
40 Vitro Deposition on the Human Epithelial Airway Barrier. *Journal of aerosol medicine and*
41 *pulmonary drug delivery*, 31(3), 170-181.
- 42 d'Angelo, I., Costabile, G., Durantie, E., Brocca, P., Rondelli, V., Russo, A., . . . Ungaro, F. (2018). Hybrid
43 Lipid/Polymer Nanoparticles for Pulmonary Delivery of siRNA: Development and Fate Upon In
44 Vitro Deposition on the Human Epithelial Airway Barrier. *J Aerosol Med Pulm Drug Deliv*, 31(3),
45 170-181. doi:10.1089/jamp.2017.1364
- 46 Darcan-Nicolaisen, Y., Meinicke, H., Fels, G., Hegend, O., Haberland, A., Köhl, A., . . . Henke, W. (2009).
47 Small interfering RNA against transcription factor STAT6 inhibits allergic airway inflammation
48 and hyperreactivity in mice. *The Journal of Immunology*, 182(12), 7501-7508.

- 1 Deng, Y., Wang, C. C., Choy, K. W., Du, Q., Chen, J., Wang, Q., . . . Tang, T. (2014). Therapeutic potentials
2 of gene silencing by RNA interference: principles, challenges, and new strategies. *Gene*, 538(2),
3 217-227.
- 4 DeVincenzo, J., Cehelsky, J. E., Alvarez, R., Elbashir, S., Harborth, J., Toudjarska, I., . . . Vaishnav, A. K.
5 (2008). Evaluation of the safety, tolerability and pharmacokinetics of ALN-RSV01, a novel RNAi
6 antiviral therapeutic directed against respiratory syncytial virus (RSV). *Antiviral research*, 77(3),
7 225-231.
- 8 DeVincenzo, J., Lambkin-Williams, R., Wilkinson, T., Cehelsky, J., Nochur, S., Walsh, E., . . . Vaishnav, A.
9 (2010). A randomized, double-blind, placebo-controlled study of an RNAi-based therapy directed
10 against respiratory syncytial virus. *Proceedings of the National Academy of Sciences*, 107(19),
11 8800-8805.
- 12 Didierlaurent, A., Goulding, J., & Hussell, T. (2007). The impact of successive infections on the lung
13 microenvironment. *Immunology*, 122(4), 457-465. doi:10.1111/j.1365-2567.2007.02729.x
- 14 Ding, L., Zhu, C., Yu, F., Wu, P., Chen, G., Ullah, A., . . . Oupicky, D. (2018). Pulmonary delivery of
15 polyplexes for combined PAI-1 gene silencing and CXCR4 inhibition to treat lung fibrosis.
16 *Nanomedicine*, 14(6), 1765-1776. doi:10.1016/j.nano.2018.05.005
- 17 Dobashi, Y., Watanabe, Y., Miwa, C., Suzuki, S., & Koyama, S. (2011). Mammalian target of rapamycin: a
18 central node of complex signaling cascades. *International journal of clinical and experimental*
19 *pathology*, 4(5), 476.
- 20 Dua, K., Chellappan, D., Singhvi, G., de Jesus, A. P. T., Gupta, G., & Hansbro, P. (2018). Targeting
21 microRNAs using nanotechnology in pulmonary diseases. *Panminerva medica*, 60(4), 230.
- 22 Dua, K., de Jesus Andreoli Pinto, T., Chellappan, D. K., Gupta, G., Bebawy, M., & Hansbro, P. M. (2018).
23 Advancements in nano drug delivery systems: a challenge for biofilms in respiratory diseases.
24 *Panminerva medica*.
- 25 Dua, K., Gupta, G., Chellappan, D., Shukla, S., & Hansbro, P. (2018). Targeting bacterial biofilms in
26 pulmonary diseases in pediatric population. *Minerva pediatrica*.
- 27 Dua, K., Hansbro, N. G., Foster, P. S., & Hansbro, P. M. (2017). MicroRNAs as therapeutics for future drug
28 delivery systems in treatment of lung diseases. *Drug delivery and translational research*, 7(1),
29 168-178.
- 30 Dua, K., Hansbro, N. G., & Hansbro, P. M. (2017). Steroid resistance and concomitant respiratory
31 infections: A challenging battle in pulmonary clinic. *EXCLI journal*, 16, 981.
- 32 Dua, K., Malya, V., Singhvi, G., Wadhwa, R., Krishna, R. V., Shukla, S. D., . . . Hansbro, P. M. (2019).
33 Increasing complexity and interactions of oxidative stress in chronic respiratory diseases: An
34 emerging need for novel drug delivery systems. *Chem Biol Interact*, 299, 168-178.
35 doi:10.1016/j.cbi.2018.12.009
- 36 Dua, K., Rapalli, V. K., Shukla, S. D., Singhvi, G., Shastri, M. D., Chellappan, D. K., . . . Pinto, T. D. J. A.
37 (2018). Multi-drug resistant Mycobacterium tuberculosis & oxidative stress complexity:
38 Emerging need for novel drug delivery approaches. *Biomedicine & Pharmacotherapy*, 107, 1218-
39 1229.
- 40 Dua, K., Rapalli, V. K., Shukla, S. D., Singhvi, G., Shastri, M. D., Chellappan, D. K., . . . Hansbro, P. M.
41 (2018). Multi-drug resistant Mycobacterium tuberculosis & oxidative stress complexity:
42 Emerging need for novel drug delivery approaches. *Biomed Pharmacother*, 107, 1218-1229.
43 doi:10.1016/j.biopha.2018.08.101
- 44 Dua, K., Shukla, S. D., de Jesus Andreoli Pinto, T., & Hansbro, P. M. (2017). Nanotechnology: Advancing
45 the translational respiratory research. *Interventional Medicine and Applied Science*, 9(1), 39-41.
- 46 Dua, K., Shukla, S. D., Tekade, R. K., & Hansbro, P. M. (2017). Whether a novel drug delivery system can
47 overcome the problem of biofilms in respiratory diseases? *Drug delivery and translational*
48 *research*, 7(1), 179-187.

- 1 Durcan, N., Murphy, C., & Cryan, S.-A. (2008). Inhalable siRNA: potential as a therapeutic agent in the
2 lungs. *Molecular pharmaceuticals*, 5(4), 559-566.
- 3 Fajt, M. L., & Wenzel, S. E. (2015). Asthma phenotypes and the use of biologic medications in asthma
4 and allergic disease: the next steps toward personalized care. *J Allergy Clin Immunol*, 135(2),
5 299-310; quiz 311. doi:10.1016/j.jaci.2014.12.1871
- 6 Feldmann, D. P., & Merkel, O. M. (2015). The advantages of pulmonary delivery of therapeutic siRNA.
7 *Therapeutic delivery*, 6(4), 407-409.
- 8 Fujita, Y., Takeshita, F., Kuwano, K., & Ochiya, T. (2013). RNAi therapeutic platforms for lung diseases.
9 *Pharmaceuticals*, 6(2), 223-250.
- 10 Gandhi, N. S., Godeshala, S., Koomoa-Lange, D.-L. T., Miryala, B., Rege, K., & Chougule, M. B. (2018).
11 Bio-reducible Poly (Amino Ethers) Based mTOR siRNA Delivery for Lung Cancer. *Pharmaceutical*
12 *research*, 35(10), 188.
- 13 Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with
14 disability for chronic obstructive pulmonary disease and asthma, 1990-2015: a systematic
15 analysis for the Global Burden of Disease Study 2015. (2017). *Lancet Respir Med*, 5(9), 691-706.
16 doi:10.1016/s2213-2600(17)30293-x
- 17 Goh, F. Y., Cook, K. L., Upton, N., Tao, L., Lah, L. C., Leung, B. P., & Wong, W. F. (2013). Receptor-
18 interacting protein 2 gene silencing attenuates allergic airway inflammation. *The Journal of*
19 *Immunology*, 1202416.
- 20 Han, S., & Roman, J. (2006). Fibronectin induces cell proliferation and inhibits apoptosis in human
21 bronchial epithelial cells: pro-oncogenic effects mediated by PI3-kinase and NF-κB. *Oncogene*,
22 25(31), 4341.
- 23 Hansbro, P. M., & Dua, K. (2018). Rajendra Awasthi, Anurag Kumar Singh, Gaurav Mishra, Anand
24 Maurya, Dinesh Kumar Chellappan, Gaurav Gupta. *Circular RNAs: Biogenesis and Functions*,
25 1087, 1.
- 26 Hansbro, P. M., Kim, R. Y., Starkey, M. R., Donovan, C., Dua, K., Mayall, J. R., . . . Wood, L. G. (2017).
27 Mechanisms and treatments for severe, steroid-resistant allergic airway disease and asthma.
28 *Immunological reviews*, 278(1), 41-62.
- 29 Hansbro, P. M., Kim, R. Y., Starkey, M. R., Donovan, C., Dua, K., Mayall, J. R., . . . Horvat, J. C. (2017).
30 Mechanisms and treatments for severe, steroid-resistant allergic airway disease and asthma.
31 *Immunol Rev*, 278(1), 41-62. doi:10.1111/imr.12543
- 32 He, M., Li, K., Yu, C., Lv, B., Zhao, N., Deng, J., . . . Wang, L. (2018). In vitro study of FUZ as a novel
33 potential therapeutic target in non-small-cell lung cancer. *Life Sci*, 197, 91-100.
34 doi:10.1016/j.lfs.2018.02.007
- 35 Hong, S.-H., Minai-Tehrani, A., Chang, S.-H., Jiang, H.-L., Lee, S., Lee, A.-Y., . . . Cho, M.-H. (2014).
36 Correction: Knockdown of the Sodium-Dependent Phosphate Co-Transporter 2b (NPT2b)
37 Suppresses Lung Tumorigenesis. *PloS one*, 9(1), 10.1371/annotation/08286cd08288-08527f-
38 08284f08214-08856f-57267107efa57267108.
- 39 Howard, K. A., Rahbek, U. L., Liu, X., Damgaard, C. K., Glud, S. Z., Andersen, M. Ø., . . . Besenbacher, F.
40 (2006). RNA interference in vitro and in vivo using a novel chitosan/siRNA nanoparticle system.
41 *Molecular Therapy*, 14(4), 476-484.
- 42 Huang, H.-Y., & Chiang, B.-L. (2009). siRNA as a therapy for asthma. *Current opinion in molecular*
43 *therapeutics*, 11(6), 652-663.
- 44 Huang, Z.-Y., Kim, M.-K., Kim-Han, T.-H., Indik, Z. K., & Schreiber, A. D. (2013). Effect of locally
45 administered Syk siRNA on allergen-induced arthritis and asthma. *Molecular immunology*, 53(1),
46 52-59.
- 47 Ihara, D., Hattori, N., Horimasu, Y., Masuda, T., Nakashima, T., Senoo, T., . . . Kohno, N. (2015).
48 Histological quantification of gene silencing by intratracheal administration of dry powdered

small-interfering RNA/chitosan complexes in the murine lung. *Pharmaceutical research*, 32(12), 3877-3885.

Jin, M., Jin, G., Kang, L., Chen, L., Gao, Z., & Huang, W. (2018). Smart polymeric nanoparticles with pH-responsive and PEG-detachable properties for co-delivering paclitaxel and survivin siRNA to enhance antitumor outcomes. *International journal of nanomedicine*, 13, 2405.

Juliano, R., Alam, M. R., Dixit, V., & Kang, H. (2008). Mechanisms and strategies for effective delivery of antisense and siRNA oligonucleotides. *Nucleic acids research*, 36(12), 4158-4171.

Kaczmarek, J. C., Kauffman, K. J., Fenton, O. S., Sadtler, K., Patel, A. K., Heartlein, M. W., . . . Anderson, D. G. (2018). Optimization of a Degradable Polymer-Lipid Nanoparticle for Potent Systemic Delivery of mRNA to the Lung Endothelium and Immune Cells. *Nano Lett*, 18(10), 6449-6454. doi:10.1021/acs.nanolett.8b02917

Kathania, M., Raje, C. I., Raje, M., Dutta, R. K., & Majumdar, S. (2011). Bfl-1/A1 acts as a negative regulator of autophagy in mycobacteria infected macrophages. *The international journal of biochemistry & cell biology*, 43(4), 573-585.

KC, R., Shukla, S. D., Gautam, S. S., Hansbro, P. M., & O'Toole, R. F. (2018). The role of environmental exposure to non-cigarette smoke in lung disease. *Clinical and Translational Medicine*, 7(1), 39. doi:10.1186/s40169-018-0217-2

Khaitov, M. R., Shilovskiy, I. P., Nikonova, A. A., Shershakova, N. N., Kamyshnikov, O. Y., Babakhin, A. A., . . . Khaitov, R. M. (2014). Small interfering RNAs targeted to interleukin-4 and respiratory syncytial virus reduce airway inflammation in a mouse model of virus-induced asthma exacerbation. *Human gene therapy*, 25(7), 642-650.

Kim, J. K., Yuk, J.-M., Kim, S. Y., Kim, T. S., Jin, H. S., Yang, C.-S., & Jo, E.-K. (2015). MicroRNA-125a inhibits autophagy activation and antimicrobial responses during mycobacterial infection. *The Journal of Immunology*, 194(11), 5355-5365.

Krisenko, M. O., & Geahlen, R. L. (2015). Calling in SYK: SYK's dual role as a tumor promoter and tumor suppressor in cancer. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1853(1), 254-263. doi:https://doi.org/10.1016/j.bbamcr.2014.10.022

Lam, J. K.-W., Liang, W., & Chan, H.-K. (2012). Pulmonary delivery of therapeutic siRNA. *Advanced drug delivery reviews*, 64(1), 1-15.

Lam, J. K., Chow, M. Y., Zhang, Y., & Leung, S. W. (2015). siRNA Versus miRNA as Therapeutics for Gene Silencing. *Mol Ther Nucleic Acids*, 4, e252. doi:10.1038/mtna.2015.23

Leung, R. K., & Whittaker, P. A. (2005). RNA interference: from gene silencing to gene-specific therapeutics. *Pharmacology & therapeutics*, 107(2), 222-239.

Li, S.-D., Chen, Y.-C., Hackett, M. J., & Huang, L. (2008). Tumor-targeted delivery of siRNA by self-assembled nanoparticles. *Molecular Therapy*, 16(1), 163-169.

Li, S.-D., & Huang, L. (2006). Targeted delivery of antisense oligodeoxynucleotide and small interference RNA into lung cancer cells. *Molecular pharmaceutics*, 3(5), 579-588.

Liu, Q., Wang, D., Xu, Z., Huang, C., Zhang, C., He, B., . . . Qian, H. (2019). Targeted delivery of Rab26 siRNA with precisely tailored DNA prism for lung cancer therapy. *ChemBiochem*. doi:10.1002/cbic.201800761

Lomas-Neira, J., Perl, M., Venet, F., Chung, C.-S., & Ayala, A. (2012). The role and source of TNF- α in hemorrhage induced priming for septic lung injury. *Shock (Augusta, Ga.)*, 37(6), 611.

Luo, Y., Zhai, X., Ma, C., Sun, P., Fu, Z., Liu, W., & Xu, J. (2012). An inhalable beta(2)-adrenoceptor ligand-directed guanidylated chitosan carrier for targeted delivery of siRNA to lung. *J Control Release*, 162(1), 28-36. doi:10.1016/j.jconrel.2012.06.005

Man, D. K., Chow, M. Y., Casattari, L., Gonzalez-Juarrero, M., & Lam, J. K. (2016). Potential and development of inhaled RNAi therapeutics for the treatment of pulmonary tuberculosis. *Advanced drug delivery reviews*, 102, 21-32.

- 1 Martinez, J., Patkaniowska, A., Urlaub, H., Lührmann, R., & Tuschl, T. (2002). Single-stranded antisense
2 siRNAs guide target RNA cleavage in RNAi. *Cell*, 110(5), 563-574.
- 3 Mealey, F. H., Kenyon, N. J., Avdalovic, M. V., & Louie, S. (2007). Difficult-to-control asthma in adults.
4 *The American journal of medicine*, 120(9), 760-763.
- 5 Meltzer, E. B., & Noble, P. W. (2008). Idiopathic pulmonary fibrosis. *Orphanet J Rare Dis*, 3, 8.
6 doi:10.1186/1750-1172-3-8
- 7 Merkel, O. M., Rubinstein, I., & Kissel, T. (2014). siRNA delivery to the lung: what's new? *Adv Drug Deliv*
8 *Rev*, 75, 112-128. doi:10.1016/j.addr.2014.05.018
- 9 Miwata, K., Okamoto, H., Nakashima, T., Ihara, D., Horimasu, Y., Masuda, T., . . . Hamada, H. (2018).
10 Intratracheal Administration of siRNA Dry Powder Targeting Vascular Endothelial Growth Factor
11 Inhibits Lung Tumor Growth in Mice. *Molecular Therapy-Nucleic Acids*, 12, 698-706.
- 12 Moding, E. J., Kastan, M. B., & Kirsch, D. G. (2013). Strategies for optimizing the response of cancer and
13 normal tissues to radiation. *Nature reviews Drug discovery*, 12(7), 526.
- 14 Moheimani, F., Hsu, A. C., Reid, A. T., Williams, T., Kicic, A., Stick, S. M., . . . Knight, D. A. (2016). The
15 genetic and epigenetic landscapes of the epithelium in asthma. *Respir Res*, 17(1), 119.
16 doi:10.1186/s12931-016-0434-4
- 17 Mokhtarieh, A. A., Lee, J., Kim, S., & Lee, M. K. (2018). Preparation of siRNA encapsulated
18 nanoliposomes suitable for siRNA delivery by simply discontinuous mixing. *Biochim Biophys Acta*
19 *Biomembr*, 1860(6), 1318-1325. doi:10.1016/j.bbmem.2018.02.027
- 20 Molet, S., Hamid, Q., Davoineb, F., Nutku, E., Tahaa, R., Pagé, N., . . . Chakir, J. (2001). IL-17 is increased
21 in asthmatic airways and induces human bronchial fibroblasts to produce cytokines. *Journal of*
22 *Allergy and Clinical Immunology*, 108(3), 430-438.
- 23 Moschos, S. A., Jones, S. W., Perry, M. M., Williams, A. E., Erjefalt, J. S., Turner, J. J., . . . Lindsay, M. A.
24 (2007). Lung delivery studies using siRNA conjugated to TAT (48– 60) and penetratin reveal
25 peptide induced reduction in gene expression and induction of innate immunity. *Bioconjugate*
26 *chemistry*, 18(5), 1450-1459.
- 27 Muddineti, O. S., Shah, A., Rompicharla, S. V. K., Ghosh, B., & Biswas, S. (2018). Cholesterol-grafted
28 chitosan micelles as a nanocarrier system for drug-siRNA co-delivery to the lung cancer cells.
29 *International journal of biological macromolecules*, 118, 857-863.
- 30 Muddineti, O. S., Shah, A., Rompicharla, S. V. K., Ghosh, B., & Biswas, S. (2018). Cholesterol-grafted
31 chitosan micelles as a nanocarrier system for drug-siRNA co-delivery to the lung cancer cells. *Int*
32 *J Biol Macromol*, 118(Pt A), 857-863. doi:10.1016/j.ijbiomac.2018.06.114
- 33 Nguyen, D. D., & Chang, S. (2017). Development of Novel Therapeutic Agents by Inhibition of Oncogenic
34 MicroRNAs. *Int J Mol Sci*, 19(1). doi:10.3390/ijms19010065
- 35 Nguyen, T., Menocal, E. M., Harborth, J., & Fruehauf, J. H. (2008). RNAi therapeutics: an update on
36 delivery. *Current opinion in molecular therapeutics*, 10(2), 158-167.
- 37 Ni, S., Liu, Y., Tang, Y., Chen, J., Li, S., Pu, J., & Han, L. (2018). GABAB receptor ligand-directed trimethyl
38 chitosan/tripolyphosphate nanoparticles and their pMDI formulation for survivin siRNA
39 pulmonary delivery. *Carbohydr Polym*, 179, 135-144. doi:10.1016/j.carbpol.2017.09.075
- 40 O'Toole, R. F., Shukla, S. D., & Walters, E. H. (2015). TB meets COPD: An emerging global co-morbidity in
41 human lung disease. *Tuberculosis (Edinb)*, 95(6), 659-663. doi:10.1016/j.tube.2015.08.005
- 42 Ogawa, Y., & Calhoun, W. J. (2006). The role of leukotrienes in airway inflammation. *Journal of Allergy*
43 *and Clinical Immunology*, 118(4), 789-798.
- 44 Oh, B., & Lee, M. (2014). Combined delivery of HMGB-1 box A peptide and S1PLyase siRNA in animal
45 models of acute lung injury. *Journal of Controlled Release*, 175, 25-35.
- 46 Oh, Y.-K., & Park, T. G. (2009). siRNA delivery systems for cancer treatment. *Advanced drug delivery*
47 *reviews*, 61(10), 850-862.

- 1 Okuda, T., Kito, D., Oiwa, A., Fukushima, M., Hira, D., & Okamoto, H. (2013). Gene silencing in a mouse
2 lung metastasis model by an inhalable dry small interfering RNA powder prepared using the
3 supercritical carbon dioxide technique. *Biological and Pharmaceutical Bulletin*, 36(7), 1183-1191.
- 4 Otsuka, M., Shiratori, M., Chiba, H., Kuronuma, K., Sato, Y., Niitsu, Y., & Takahashi, H. (2017). Treatment
5 of pulmonary fibrosis with siRNA against a collagen-specific chaperone HSP47 in vitamin A-
6 coupled liposomes. *Exp Lung Res*, 43(6-7), 271-282. doi:10.1080/01902148.2017.1354946
- 7 Pai, S., Lin, Y., Macaes, B., Meneshian, A., Hung, C., & Wu, T. (2006). Prospects of RNA interference
8 therapy for cancer. *Gene therapy*, 13(6), 464.
- 9 Peters, S. P., Ferguson, G., Deniz, Y., & Reisner, C. (2006). Uncontrolled asthma: a review of the
10 prevalence, disease burden and options for treatment. *Respir Med*, 100(7), 1139-1151.
11 doi:10.1016/j.rmed.2006.03.031
- 12 Pirollo, K. F., Zon, G., Rait, A., Zhou, Q., Yu, W., Hogrefe, R., & Chang, E. H. (2006). Tumor-targeting
13 nanoimmunoliposome complex for short interfering RNA delivery. *Human gene therapy*, 17(1),
14 117-124.
- 15 Qiu, C., Peng, W., Shi, F., & Zhang, T. (2012). Bottom-up assembly of RNA nanoparticles containing phi29
16 motor pRNA to silence the asthma STAT5b gene. *Genet Mol Res*, 11(3), 3236-3245.
- 17 Qiu, Y., Chow, M. Y. T., Liang, W., Chung, W. W. Y., Mak, J. C. W., & Lam, J. K. W. (2017). From Pulmonary
18 Surfactant, Synthetic KL4 Peptide as Effective siRNA Delivery Vector for Pulmonary Delivery. *Mol*
19 *Pharm*, 14(12), 4606-4617. doi:10.1021/acs.molpharmaceut.7b00725
- 20 Qiu, Y., Lam, J., Leung, S., & Liang, W. (2016). Delivery of RNAi therapeutics to the airways—from bench
21 to bedside. *Molecules*, 21(9), 1249.
- 22 Ramakrishnan, L. (2012). Revisiting the role of the granuloma in tuberculosis. *Nature Reviews*
23 *Immunology*, 12(5), 352.
- 24 Rapalli, V. K., Singhvi, G., Dubey, S. K., Gupta, G., Chellappan, D. K., & Dua, K. (2018). Emerging landscape
25 in psoriasis management: From topical application to targeting biomolecules. *Biomedicine &*
26 *Pharmacotherapy*, 106, 707-713.
- 27 Rennard, S. I., & Vestbo, J. (2006). COPD: the dangerous underestimate of 15%. *Lancet*, 367(9518), 1216-
28 1219. doi:10.1016/s0140-6736(06)68516-4
- 29 Rettig, G. R., & Behlke, M. A. (2012). Progress toward in vivo use of siRNAs-II. *Molecular therapy*, 20(3),
30 483-512.
- 31 Roberts, E. A., Chua, J., Kyei, G. B., & Deretic, V. (2006). Higher order Rab programming in
32 phagolysosome biogenesis. *The Journal of cell biology*, 174(7), 923-929.
- 33 Rodriguez-Roisin, R. (2000). Toward a Consensus Definition for COPD Exacerbations. *CHEST*, 117(5),
34 398S-401S. doi:10.1378/chest.117.5_suppl_2.398S
- 35 Rosas-Taraco, A. G., Higgins, D. M., Sanchez-Campillo, J., Lee, E. J., Orme, I. M., & González-Juarrero, M.
36 (2009). Intrapulmonary delivery of XCL1-targeting small interfering RNA in mice chronically
37 infected with Mycobacterium tuberculosis. *American journal of respiratory cell and molecular*
38 *biology*, 41(2), 136-145.
- 39 Rosas-Taraco, A. G., Higgins, D. M., Sánchez-Campillo, J., Lee, E. J., Orme, I. M., & González-Juarrero, M.
40 (2009). Intrapulmonary delivery of XCL1-targeting small interfering RNA in mice chronically
41 infected with Mycobacterium tuberculosis. *American journal of respiratory cell and molecular*
42 *biology*, 41(2), 136-145.
- 43 Rosas-Taraco, A. G., Higgins, D. M., Sánchez-Campillo, J., Lee, E. J., Orme, I. M., & González-Juarrero, M.
44 (2011). Local pulmonary immunotherapy with siRNA targeting TGFβ1 enhances antimicrobial
45 capacity in Mycobacterium tuberculosis infected mice. *Tuberculosis*, 91(1), 98-106.
- 46 Rosell, R., & Karachaliou, N. (2016). Large-scale screening for somatic mutations in lung cancer. *Lancet*,
47 387(10026), 1354-1356. doi:10.1016/s0140-6736(15)01125-3

- 1 Rosenwasser, L. J., Zimmermann, N., Hershey, G. K., Foster, P. S., & Rothenberg, M. E. (2003).
2 Chemokines in asthma: cooperative interaction between chemokines and IL-13. *Journal of*
3 *Allergy and Clinical Immunology*, 111(2), 227-242.
- 4 Sadowski, K., Kotulska, K., & Jóźwiak, S. (2016). Management of side effects of mTOR inhibitors in
5 tuberous sclerosis patients. *Pharmacological Reports*, 68(3), 536-542.
- 6 Sapey, E., & Stockley, R. A. (2006). COPD exacerbations · 2: Aetiology. *Thorax*, 61(3), 250-258.
7 doi:10.1136/thx.2005.041822
- 8 Sasaki, S., & Guo, S. (2018). Nucleic Acid Therapies for Cystic Fibrosis. *Nucleic Acid Ther*, 28(1), 1-9.
9 doi:10.1089/nat.2017.0696
- 10 Schultheis, B., Strumberg, D., Santel, A., Vank, C., Gebhardt, F., Keil, O., . . . Khan, M. (2014). First-in-
11 human phase I study of the liposomal RNA interference therapeutic Atu027 in patients with
12 advanced solid tumors. *Journal of Clinical Oncology*, 32(36), 4141-4148.
- 13 Sel, S., Henke, W., Dietrich, A., Herz, U., & Renz, H. (2006). Treatment of allergic asthma by targeting
14 transcription factors using nucleic-acid based technologies. *Curr Pharm Des*, 12(25), 3293-3304.
- 15 Shastri, M. D., Shukla, S. D., Chong, W. C., Dua, K., Peterson, G. M., Patel, R. P., . . . O'Toole, R. F. (2018).
16 Role of Oxidative Stress in the Pathology and Management of Human Tuberculosis. *Oxid Med*
17 *Cell Longev*, 2018, 7695364. doi:10.1155/2018/7695364
- 18 Siegel, R., Naishadham, D., & Jemal, A. (2012). Cancer statistics for hispanics/latinos, 2012. *CA: a cancer*
19 *journal for clinicians*, 62(5), 283-298.
- 20 Silva Miranda, M., Breiman, A., Allain, S., Deknuydt, F., & Altare, F. (2012). The tuberculous granuloma:
21 an unsuccessful host defence mechanism providing a safety shelter for the bacteria? *Clinical and*
22 *Developmental Immunology*, 2012.
- 23 Sioud, M. (2015). Overcoming the challenges of siRNA activation of innate immunity: design better
24 therapeutic siRNAs. In *RNA Interference* (pp. 301-319): Springer.
- 25 Sorber, L., Zwaenepoel, K., Deschoolmeester, V., Van Schil, P. E., Van Meerbeeck, J., Lardon, F., . . .
26 Pauwels, P. (2017). Circulating cell-free nucleic acids and platelets as a liquid biopsy in the
27 provision of personalized therapy for lung cancer patients. *Lung Cancer*, 107, 100-107.
28 doi:10.1016/j.lungcan.2016.04.026
- 29 Staff, P. O. (2014). Correction: Gene Silencing of SOCS3 by siRNA Intranasal Delivery Inhibits Asthma
30 Phenotype in Mice. *PloS one*, 9(8), e105924.
- 31 Sung, D. K., Kong, W. H., Park, K., Kim, J. H., Kim, M. Y., Kim, H., & Hahn, S. K. (2013). Noncovalently
32 PEGylated CTGF siRNA/PDMAEMA complex for pulmonary treatment of bleomycin-induced lung
33 fibrosis. *Biomaterials*, 34(4), 1261-1269.
- 34 Taratula, O., Kuzmov, A., Shah, M., Garbuzenko, O. B., & Minko, T. (2013). Nanostructured lipid carriers
35 as multifunctional nanomedicine platform for pulmonary co-delivery of anticancer drugs and
36 siRNA. *Journal of Controlled Release*, 171(3), 349-357.
- 37 Tiseo, M., Bartolotti, M., Gelsomino, F., & Bordi, P. (2010). Emerging role of gefitinib in the treatment of
38 non-small-cell lung cancer (NSCLC). *Drug design, development and therapy*, 4, 81.
- 39 Tompkins, S. M., Lo, C.-Y., Tumpey, T. M., & Epstein, S. L. (2004). Protection against lethal influenza virus
40 challenge by RNA interference in vivo. *Proceedings of the National Academy of Sciences*,
41 101(23), 8682-8686.
- 42 Veilleux, D., Gopalakrishna Panicker, R. K., Chevrier, A., Biniecki, K., Lavertu, M., & Buschmann, M. D.
43 (2018). Lyophilisation and concentration of chitosan/siRNA polyplexes: Influence of buffer
44 composition, oligonucleotide sequence, and hyaluronic acid coating. *J Colloid Interface Sci*, 512,
45 335-345. doi:10.1016/j.jcis.2017.09.084
- 46 Vogelstein, B., & Kinzler, K. W. (2004). Cancer genes and the pathways they control. *Nature medicine*,
47 10(8), 789.

- 1 Wakashin, H., Hirose, K., Maezawa, Y., Kagami, S.-i., Suto, A., Watanabe, N., . . . Iwakura, Y. (2008). IL-23
2 and Th17 cells enhance Th2-cell-mediated eosinophilic airway inflammation in mice. *American*
3 *journal of respiratory and critical care medicine*, 178(10), 1023-1032.
- 4 Wang, J., Lu, Z., Wientjes, M. G., & Au, J. L.-S. (2010). Delivery of siRNA therapeutics: barriers and
5 carriers. *The AAPS journal*, 12(4), 492-503.
- 6 Wang, J., Yang, K., Zhou, L., Wu, Y., Zhu, M., Lai, X., . . . Huang, C. (2013). MicroRNA-155 promotes
7 autophagy to eliminate intracellular mycobacteria by targeting Rheb. *PLoS pathogens*, 9(10),
8 e1003697.
- 9 Wang, J., Zhu, M., Wang, L., Chen, C., & Song, Y. (2019). Amphiregulin potentiates airway inflammation
10 and mucus hypersecretion induced by urban particulate matter via the EGFR-PI3K α -AKT/ERK
11 pathway. *Cellular signalling*, 53, 122-131.
- 12 Wang, L., Huang, W., Zhang, L., Chen, Q., & Zhao, H. (2018). Molecular pathogenesis involved in human
13 idiopathic pulmonary fibrosis based on an integrated microRNA-mRNA interaction network. *Mol*
14 *Med Rep*, 18(5), 4365-4373. doi:10.3892/mmr.2018.9456
- 15 Wang, X., Xu, W., Mohapatra, S., Kong, X., Li, X., Lockey, R. F., & Mohapatra, S. S. (2008). Prevention of
16 airway inflammation with topical cream containing imiquimod and small interfering RNA for
17 natriuretic peptide receptor. *Genetic vaccines and therapy*, 6(1), 7.
- 18 Wark, P. A., Tooze, M., Powell, H., & Parsons, K. (2013). Viral and bacterial infection in acute asthma and
19 chronic obstructive pulmonary disease increases the risk of readmission. *Respirology*, 18(6), 996-
20 1002. doi:10.1111/resp.12099
- 21 Watts, J. K., & Corey, D. R. (2010). Clinical status of duplex RNA. *Bioorganic & medicinal chemistry*
22 *letters*, 20(11), 3203-3207.
- 23 Whitehead, K. A., Langer, R., & Anderson, D. G. (2009). Knocking down barriers: advances in siRNA
24 delivery. *Nature reviews Drug discovery*, 8(2), 129.
- 25 Wu, W., Chen, H., Li, Y.-M., Wang, S.-Y., Diao, X., & Liu, K.-G. (2014). Intranasal sirna targeting c-kit
26 reduces airway inflammation in experimental allergic asthma. *International journal of clinical*
27 *and experimental pathology*, 7(9), 5505.
- 28 Xie, Y., Kim, N. H., Nadithe, V., Schalk, D., Thakur, A., Kılıç, A., . . . Merkel, O. M. (2016). Targeted delivery
29 of siRNA to activated T cells via transferrin-polyethylenimine (Tf-PEI) as a potential therapy of
30 asthma. *Journal of Controlled Release*, 229, 120-129.
- 31 Xie, Y., & Merkel, O. M. (2015). Pulmonary delivery of siRNA via polymeric vectors as therapies of
32 asthma. *Archiv der Pharmazie*, 348(10), 681-688.
- 33 Xu, C.-N., Tian, H.-Y., Wang, Y.-B., Du, Y., Chen, J., Lin, L., . . . Chen, X.-S. (2017). Anti-tumor effects of
34 combined doxorubicin and siRNA for pulmonary delivery. *Chinese Chemical Letters*, 28(4), 807-
35 812.
- 36 Xu, C.-X., Jere, D., Jin, H., Chang, S.-H., Chung, Y.-S., Shin, J.-Y., . . . Chae, C.-H. (2008). Poly (ester amine)-
37 mediated, aerosol-delivered Akt1 small interfering RNA suppresses lung tumorigenesis.
38 *American journal of respiratory and critical care medicine*, 178(1), 60-73.
- 39 Xu, C., Tian, H., Sun, H., Jiao, Z., Zhang, Y., & Chen, X. (2015). A pH sensitive co-delivery system of siRNA
40 and doxorubicin for pulmonary administration to B16F10 metastatic lung cancer. *RSC Advances*,
41 5(125), 103380-103385.
- 42 Xu, C., Tian, H., Wang, P., Wang, Y., & Chen, X. (2016). The suppression of metastatic lung cancer by
43 pulmonary administration of polymer nanoparticles for co-delivery of doxorubicin and Survivin
44 siRNA. *Biomaterials science*, 4(11), 1646-1654.
- 45 Xu, P. Y., Kankala, R. K., Pan, Y. J., Yuan, H., Wang, S. B., & Chen, A. Z. (2018). Overcoming multidrug
46 resistance through inhalable siRNA nanoparticles-decorated porous microparticles based on
47 supercritical fluid technology. *Int J Nanomedicine*, 13, 4685-4698. doi:10.2147/ijn.s169399

- Yoon, P. O., Park, J. W., Lee, C.-M., Kim, S. H., Kim, H.-N., Ko, Y., . . . Kwon, T. (2016). Self-assembled micelle interfering RNA for effective and safe targeting of dysregulated genes in pulmonary fibrosis. *Journal of Biological Chemistry*, jbc. M115. 693671.
- Youngren-Ortiz, S. R., Gandhi, N. S., España-Serrano, L., & Chougule, M. B. (2016). Aerosol delivery of siRNA to the lungs. Part 1: Rationale for gene delivery systems. *KONA Powder and Particle Journal*, 33, 63-85.
- Youngren-Ortiz, S. R., Gandhi, N. S., España-Serrano, L., & Chougule, M. B. (2017). Aerosol Delivery of siRNA to the Lungs. Part 2: Nanocarrier-based Delivery Systems. *KONA Powder and Particle Journal*, 34, 44-69.
- Yuan, Z. Q., Chen, W. L., You, B. G., Liu, Y., Yang, S. D., Li, J. Z., . . . Zhang, X. N. (2017). Multifunctional nanoparticles co-delivering EZH2 siRNA and etoposide for synergistic therapy of orthotopic non-small-cell lung tumor. *J Control Release*, 268, 198-211. doi:10.1016/j.jconrel.2017.10.025
- Zhang, C., Tang, N., Liu, X., Liang, W., Xu, W., & Torchilin, V. P. (2006). siRNA-containing liposomes modified with polyarginine effectively silence the targeted gene. *Journal of controlled release*, 112(2), 229-239.
- Zhang, D., Lee, H., Wang, X., Rai, A., Groot, M., & Jin, Y. (2018). Exosome-Mediated Small RNA Delivery: A Novel Therapeutic Approach for Inflammatory Lung Responses. *Mol Ther*, 26(9), 2119-2130. doi:10.1016/j.ymthe.2018.06.007
- Zhang, Z., Jiang, G., Yang, F., & Wang, J. (2006). Knockdown of mutant K-ras expression by adenovirus-mediated siRNA inhibits the in vitro and in vivo growth of lung cancer cells. *Cancer biology & therapy*, 5(11), 1481-1486.
- Zhang, Z., Yang, X., Zhang, Y., Zeng, B., Wang, S., Zhu, T., . . . Yang, R. (2006). Delivery of telomerase reverse transcriptase small interfering RNA in complex with positively charged single-walled carbon nanotubes suppresses tumor growth. *Clinical Cancer Research*, 12(16), 4933-4939.

Table 1: siRNA delivery systems for targeted pulmonary therapy

Delivery system	siRNA delivered	Target	Route of administration	Outcome	Reference
Polyethylenimine-cis-aconitic anhydride-doxorubicin and (B cell) Bcl2 siRNA complex nanoparticles	Bcl-2 siRNA	Lung cancer	Intratracheal	The polyethylenimine-cis-aconitic anhydride-doxorubicin/Bcl2 siRNA complex nanoparticles are for treating metastatic lung cancer by pulmonary delivery with low side effects on the normal tissues.	(C. Xu et al., 2015)
Arginine-glycine-aspartic acid peptide (RGD) gold nanoparticles	c-mycsiRNA	Lung cancer	Intratracheal	c-myc-RGD gold nanoparticles are capable of targeting tumor cells, significant tumor growth inhibition, as well as extended survival of mice bearing tumors.	(Conde et al., 2013)

Glycerol propoxylate triacrylate and spermine	Sodium-dependent phosphate co-transporter 2b(NPT2b) siRNA	Lung cancer	Intratracheal	siNPT2b successfully suppressed lung cancer growth and decreased cancer cell proliferation and angiogenesis, facilitating apoptosis.	(Hong et al., 2014)
Chitosan dry powder	Luciferase siRNA	Lung cancer	Intratracheal	siRNA/chitosan powder prepared using supercritical carbon dioxide has an effective and specific gene silencing against the tumor cells metastasized in the lungs of mice	(Okuda et al., 2013)
Naked siRNA or TransIT-TKO	Phosphoproteins siRNA	Parainfluenza virus (PIV), Respiratory syncytial virus (RSV)	Intranasal	Animals were successfully protected from RSV and PIV infections specifically	(Bitko, Musiyenko, Shulyayeva, & Barik, 2005)
Naked siRNA	Nucleocapsid gene-specific siRNA	Viral infections	Intranasal	Significant reductions of viral load were achieved in both prophylactic and therapeutic regimens	(Alvarez et al., 2009)
Oligofectamine	Nucleocapsid protein and Polymerase acidic protein	Influenza type A	Intranasal and hydrodynamic injection	Treated animals lung virus titres were reduced and protected from lethal challenge with highly pathogenic viruses	(Tompkins, Lo, Tumpey, & Epstein, 2004)
Naked siRNA	Lymphotactin (XCL1) siRNA	Tuberculosis	Intratracheal	XCL1 expression in the lungs was significantly suppressed; decreased T lymphocytes, IFN- response and disorganized granulomatous lesions and high fibrosis	(Rosas-Taraco, Higgins, Sánchez-Campillo, et al., 2009)
Naked siRNA	Transforming growth factor- β 1 siRNA	Tuberculosis	Intratracheal	Increased expression of antimicrobial mediators, with the reduced bacterial load in the lungs of treated mice	(Rosas-Taraco et al., 2011)
Naked siRNA	Suppressor	Asthma	Intranasal	Decrease in lung	(Staff, 2014)

	s of cytokine signaling protein 3 (SOCS) siRNA			eosinophilia, normalization of hyperresponsiveness, increase in mucus secretion and reduction in collagen deposition in the lungs.	
Naked siRNA	Interleukin -4 siRNA and Phosphopr oteins siRNA	Asthma	Intranasal	Eosinophilia in bronchoalveolar lavage fluid, hyperresponsiveness and airway inflammation were significantly reduced	(Khaitov et al., 2014)
Naked siRNA	Signal transducer and activator of transcription factor 6 (STAT6) siRNA	Asthma	Intratracheal and intranasal	Allergen-induced lung inflammation was significantly reduced and Expression of key cytokines (IL-4, IL-13) and allergen-induced inflammation in lung tissues were significantly reduced	(Darcen- Nicolaisen et al., 2009)
Naked siRNA	Receptor- interacting protein 2 (Rip2) siRNA	Asthma	Intratracheal	Ovalbumin-induced cytokine release, inflammatory cell infiltration and mucus hypersecretion was inhibited. elevation of serum Ovalbumin- specific IgE level was markedly suppressed	(Goh et al., 2013)
Naked siRNA	cluster of differentiation 86 (CD86) siRNA	Asthma	Intratracheal	Ovalbumin-induced airway eosinophilia, airway hyperresponsiveness, and cytokines production was reduced	(Asai-Tajiri et al., 2014)
Naked siRNA	Spleen tyrosine kinase (syk) siRNA	Asthma	Intranasal	siRNA administration by intranasal route inhibited inflammatory cells in the bronchoalveolar lavage fluid (BALF) of allergen sensitized mice.	(Z.-Y. Huang, Kim, Kim- Han, Indik, & Schreiber, 2013)
Naked siRNA	c-kit siRNA	Asthma	Intranasal	Airway mucus secretion and eosinophil infiltration in BALF was effectively reduced. C-kit further	(Wu et al., 2014)

				reduced the production of stem cell factor, IL-4, and IL-5, but had no effect on interferon- γ (IFN- γ) generation	
Transferrin polyethylenimine (Tf-PEI)	Fluorescently labeled siRNA	Asthma	Intratracheal	Tf-PEI polyplexes selectively delivered siRNA to activated T cells	(Xie et al., 2016)
R3V6 peptides were used as a carrier. (Ternary complex of siS1PLYase, HMGB1A, and R3V6 was produced by charge interaction)	siS1PLYase/HMGB1A/R3V6 ternary complex	Acute lung injury	Intratracheal	siS1PLYase/HMGB1A/R3V6 complex reduced the levels of IL-6 and TNF- α more efficiently compared to HMGB1A alone and siS1PLYase/R3V6 complex in lipopolysaccharides activated macrophages and reduced the inflammatory response and apoptosis in acute lung injury	(B. Oh & Lee, 2014)
Naked siRNA, i.v, liposomes	Tumor necrosis factor- α	Acute lung injury	Intratracheal	Systemic injection but not intratracheal delivery of TNF- α siRNA significantly reduced the incidence of acute lung injury. Results suggest pulmonary endothelial and/or other possible vascular resident cells, not epithelial cells, play a greater role in mediating the TNF- α priming response in hemorrhage/sepsis-induced acute lung injury.	(Lomas-Neira, Perl, Venet, Chung, & Ayala, 2012)
Naked siRNA	Transforming growth factor- β 1 (TGF- β 1) siRNA	Pulmonary fibrosis	Intratracheal	Levels of inflammatory cytokines, including IFN- α and IFN- β , were not significantly affected, whereas TGF- β 1 was significantly inhibited.	(Corina N D'Alessandro-Gabazza et al., 2012)
PEGylated poly(dimethylamino)ethylmethacrylate (PDMAEMA)	Connective tissue growth factor (CTGF) siRNA	Pulmonary fibrosis	Intratracheal	There was a reduction in collagen deposition, inflammatory cytokines production and drastic attenuation of pulmonary fibrosis	(Sung et al., 2013)

Self-assembled micelle interfering RNA (siRNA) nanoparticles	Amphiregulin (AR) and connective tissue growth factor (CTGF) targeting siRNA	Pulmonary fibrosis	Intratracheal/intravenous	Collagen accumulation was significantly reduced and lung function was substantially restored in TGF- β transgenic mice.	(Yoon et al., 2016)
Chitosan-based siRNA nanoparticle	siRNA specific to the BCR/ABL-1 junction sequence	Potential of chitosan nanocarriers	Nasal	In bronchiole epithelial cells of transgenic EGFP (endogenous enhanced green fluorescent protein) mice.	(Howard et al., 2006)
Spray dried naked siRNA using L-leucine dispersion enhancer	siRNA targeting interleukin 10	2% W/W siRNA developed into an inhalable dry powder	Not applicable	The integrity of siRNA was successfully retained after spray drying. Spray dried powder were crystal in nature with low moisture levels traits stable formulation.	(Chow et al., 2017)