Radiographically Occult Medial Cuneiform Impaction Fracture

Mabry, L., Patti, T. N., & Bleakley, C. M. (2019). Radiographically Occult Medial Cuneiform Impaction Fracture. *Journal of Orthopaedic and Sports Physical Therapy*, 49(9), 675-675. <https://doi.org/10.2519/jospt.2019.8778>

[Link to publication record in Ulster University Research Portal](#)

Published in:

Journal of Orthopaedic and Sports Physical Therapy

Publication Status:

Published (in print/issue): 30/09/2019

DOI:

[10.2519/jospt.2019.8778](https://doi.org/10.2519/jospt.2019.8778)

Document Version

Author Accepted version

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

[Heading] MUSCULOSKELETAL IMAGING

Radiographically Occult Medial Cuneiform Impaction Fracture

Lance M. Mabry, PT, DPT, Department of Physical Therapy, High Point University, High Point, NC.

Taylor N. Patti, Physical Therapy Doctoral Program, High Point University, High Point, NC.

Chris M. Bleakley, PT, PhD, Department of Physical Therapy, High Point University, High Point, NC.

A 25-year-old woman reported to the emergency department with right medial midfoot pain after kicking a wall, inducing a first-ray axial/plantar flexion compressive force. The patient was unable to weight bear immediately after the injury. Erythema and edema ~~was~~were present at the medial midfoot. Radiographs were noncontributory (**FIGURE 1**, available at www.jospt.org). The patient was diagnosed with a metatarsal contusion and was instructed to use ice and, ibuprofen_; and to weight bear as tolerated on crutches.

The patient returned to full weight bearing 3 days later_; however, persistent pain 3.5 weeks post injury led her to seek a direct-access physical therapy evaluation. Her pain was exacerbated with the first steps when getting out of bed and by walking barefoot. The examination revealed mild ecchymosis at the medial and plantar midfoot. During gait, the patient ~~was noted to bore~~weight bear along the lateral foot to avoid pronation in stance and experienced 3/10 to 4/10 pain. The medial cuneiform was tender to palpation. Due to concern for a medial cuneiform fracture or a medial midfoot sprain, the physical therapist requested magnetic resonance imaging, which confirmed a medial cuneiform fracture (**FIGURE 2; FIGURE 3**, available at www.jospt.org).

The patient utilized a cam boot for 2 weeks; transitioned to a flat, hard-soled shoe at 5 weeks post injury_; and footwear restrictions were removed 6 weeks post

injury. The patient returned to running and lunges with only mild discomfort 12 weeks post injury.

Isolated medial cuneiform fractures are commonly missed at baseline evaluation and are typically occult in initial radiographs.² Definitive diagnosis usually occurs after continued symptoms prompt advanced diagnostic imaging. Conservative management is commonly sufficient to restore prior function.¹ *J Orthop Sports Phys Ther* 2019;49(9):xxx. doi:10.2519/jospt.2019.8778

REFERENCES

1. Babu NS, Gambardella GV, Bowlby MA. Isolated fracture of the medial cuneiform: a case report. *J Am Podiatr Med Assoc.* 2017;107:436-439. <https://doi.org/10.7547/16-013>
2. Eraslan A, Ozyurek S, Erol B, Ercan E. Isolated medial cuneiform fracture: a commonly missed fracture. *BMJ Case Rep.* 2013;2013:bcr2013010093. <https://doi.org/10.1136/bcr-2013-010093>

[online]

FIGURE 1. (A) Anterior-to-posterior, (B) oblique, and (C) lateral radiographs of the right foot, taken on the day of the injury, demonstrating no signs of fracture.

FIGURE 2. (A) Long-axis, axial, T2-weighted right-foot magnetic resonance image demonstrating cortical irregularity (arrow). (B) Sagittal, T1-weighted right-foot magnetic resonance image demonstrating signal hypointensity (arrow). (C) Sagittal, short-tau inversion recovery, weighted right-foot magnetic resonance image demonstrating signal hyperintensity (arrow) of the medial cuneiform without involvement of the plantar metatarsal ligaments, consistent with an isolated medial cuneiform impaction fracture.

[online]

FIGURE 3. [scrollable MRI]