



Anti-asthmatic prescriptions in children with and without congenital anomalies: a European data linkage study

Divin, N., Garne, E., Morris, J. K., & Loane, M. (2022). Anti-asthmatic prescriptions in children with and without congenital anomalies: a European data linkage study. *International Journal of Population Data Science*, 7(3), 117-117. <https://doi.org/10.23889/ijpds.v7i3.1889>

[Link to publication record in Ulster University Research Portal](#)

Published in:

International Journal of Population Data Science

Publication Status:

Published (in print/issue): 25/08/2022

DOI:

[10.23889/ijpds.v7i3.1889](https://doi.org/10.23889/ijpds.v7i3.1889)

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Anti-asthmatic prescriptions in children with and without congenital anomalies: a European data linkage study.

Natalie Divin¹, Ester Garne², Joan K Morris³, and Maria Loane¹¹Ulster University²Pediatric Department, Hospital Lillebaelt Kolding³St George's University of London

Objectives

Asthma is the most common chronic disease in childhood, yet little is known about rates of asthma and wheezing in children with congenital anomalies. This study explored the prevalence and risk of receiving anti-asthmatic prescriptions in children with congenital anomalies compared to children without anomalies in six European regions/countries.

Approach

This was a EUROlinkCAT population-based linkage cohort study involving children from 0-9 years of age born between 2000-2014. Congenital anomaly data from six EUROCAT registries were linked to births data in national/vital statistics and to electronic prescription databases. Prescription/pharmacy dispensing records across regions were standardised to a Common Data Model. Anatomical Therapeutic Chemical classification codes beginning with R03 were used to identify anti-asthmatic prescriptions. Random-effects meta-analyses were performed to identify both the relative risk (RR) of receiving >1 anti-asthmatic prescription in a year relative to the reference group, and the heterogeneity of prevalence rates across registries and age group.

Results

A total of 5.1% of children with congenital anomalies and 4.9% of reference children were dropped from the study as they were not linked. Children with congenital anomalies (n=60,662) had a higher prevalence of >1 anti-asthmatic prescription and a significantly higher risk of being prescribed anti-asthmatics (RR=1.41, 95% CI 1.35-1.48) compared to reference children (n=1,722,912). The increased risk was consistent across all age groups. Children with congenital anomalies were more likely to be prescribed beta-2 agonists (RR=1.71, 95% CI 1.60-1.83) and inhaled corticosteroids (RR=1.74, 95% CI 1.61-1.87). Children with oesophageal atresia, diaphragmatic hernia, genetic syndromes and chromosomal anomalies had over twice the risk of being prescribed anti-asthmatics compared to reference children. Regional differences in prevalence and risk of anti-asthmatic prescriptions were identified.

Conclusion

Children aged <10 years with congenital anomalies consistently had higher prevalence and risk of receiving >1 anti-asthmatic prescription across age group and across European regions. This study demonstrates that information on the prevalence of anti-asthmatic prescriptions issued/dispensed can be obtained through data linkage to monitor changes in prevalence over time.

