



Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach

Shimamoto, K., McElroy, E., & Ibuka, Y. (2022). Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach. *SSM - Population Health*, 20, [101269]. <https://doi.org/10.1016/j.ssmph.2022.101269>

[Link to publication record in Ulster University Research Portal](#)

Published in:
SSM - Population Health

Publication Status:
Published (in print/issue): 31/12/2022

DOI:
[10.1016/j.ssmph.2022.101269](https://doi.org/10.1016/j.ssmph.2022.101269)

Document Version
Author Accepted version

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Journal Pre-proof



Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach

Kyoko Shimamoto, Eoin McElroy, Yoko Ibuka

PII: S2352-8273(22)00248-8

DOI: <https://doi.org/10.1016/j.ssmph.2022.101269>

Reference: SSMPH 101269

To appear in: *SSM - Population Health*

Received Date: 18 May 2022

Revised Date: 30 September 2022

Accepted Date: 11 October 2022

Please cite this article as: Shimamoto K., McElroy E. & Ibuka Y., Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach, *SSM - Population Health* (2022), doi: <https://doi.org/10.1016/j.ssmph.2022.101269>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2022 Published by Elsevier Ltd.

Title: Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach

Authors: Kyoko Shimamoto^{1,2}, Eoin McElroy³ and Yoko Ibuka⁴

Affiliations:

1. Keio Global Research Institute, Keio University, 2-15-45 Mita, Minato-ku Tokyo 108-8345, Japan
2. Graduate School of Health Management, Keio University, 35 Shinanomachi, Shinjyuku-ku Tokyo 160-8582, Japan
3. School of Psychology, Ulster University, Cromore Road, Coleraine, Co. Londonderry BT52 1SA, United Kingdom
4. Department of Economics, Keio University, 2-15-45 Mita, Minato-ku Tokyo 108-8345, Japan

Corresponding author: Kyoko Shimamoto, kyoko.shimamoto@keio.jp

Declarations:

The authors declared that they have no competing interests.

Acknowledgement:

Authors acknowledge funding from the National Institute of Public Health of Japan; valuable contributions from Dr Yumiko Hosaka for conceptualization, literature search and review; and management support from the fund managers, Prof Hiroyuki Ishida, Prof Toru Takebayashi, and Prof Rei Goto.

1 **Title:** Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic
2 consequences in Japan: A structural equation modeling approach

3
4 **Abstract:**

5 **Background.** Health inequity in relation to COVID-19 infection and socioeconomic
6 consequences is a major global concern. Mental health issues in vulnerable populations have
7 received special attention in research and practice during the COVID-19 pandemic. However,
8 there is limited evidence on the nature of the anxieties experienced as a result of COVID-19,
9 and how such concerns vary across demographic groups.

10 **Aim.** This study examines anxiety among the working population of Japan (aged 18-59), in
11 terms of both COVID-19 infection and socioeconomic consequences, using an internationally
12 validated tool, the Pandemic Anxiety Scale (PAS).

13 **Methods.** Data were collected using an online survey (n=2,764). The analyses included an
14 exploratory factor analysis (EFA), a confirmatory factor analysis (CFA), and structural
15 equation modeling (SEM), followed by validation of the Japanese version of the PAS.

16 **Results.** A two-factor latent variable model shows the multidimensionality of anxiety in
17 regard to the COVID-19 pandemic and the disparity across population groups in predicting
18 the two defined anxiety dimensions. Several path coefficients showed somewhat unexpected
19 and/or unique results from Japan compared with previous European studies. Specifically,
20 self-reported health status was not significantly related to disease anxiety, and those who
21 were not in paid employment reported lower consequence anxiety. The SEM results showed
22 a greater number of significant exogenous variables for consequence anxiety compared to
23 disease anxiety, highlighting disparities in pandemic anxiety by socioeconomic status in
24 regard to socioeconomic consequences of the pandemic.

25 **Conclusion.** In contrast to existing European studies, evidence from the current study
26 suggests contextual patterns of health inequity. Due to the prolonged socioeconomic

- 1 consequences of the pandemic, multidisciplinary research on mental health issues and the
- 2 quality of life remains an important research agenda in exploring socioeconomic measures in
- 3 context, towards addressing inequity concerns.

Journal Pre-proof

1 **1. Background**

2
3 During the COVID-19 pandemic, adverse socioeconomic impacts and
4 disproportionate effects among vulnerable population groups have been studied globally.
5 Besides macro-economic repercussions of the pandemic (1, 2), disproportionate
6 socioeconomic impacts on low socioeconomic status groups have been reported in high-,
7 middle-, and low-income countries. For example, evidence suggests that younger
8 generations, low-income groups, women and children are particularly vulnerable to adverse
9 socioeconomic consequences of the pandemic through increased risk of unemployment,
10 decreased income, and domestic violence (3-6). The circumstances of the vulnerable
11 population groups and their trends reportedly differ in each setting, in terms of the pattern of
12 unemployment rate by gender, and the extent of the impact on poverty, household income
13 decline, and food insecurity by country. Thus, a contextual analysis and intervention across
14 different population groups would remain critical by setting.

15 Health equity has been receiving increasing attention in the last decade and especially
16 in the Sustainable Development Goals (SDGs) era. Generally, health equity is referred to and
17 defined as “the absence of unfair and avoidable or remediable differences in health among
18 population groups defined socially, economically, demographically or geographically”(7-9).
19 Since the onset of the COVID-19 pandemic, health equity perspectives have been further
20 underscored in the relevant health issues, including but not limited to access and delivery of
21 preventive and curative health care against COVID-19 infection (e.g., vaccination), infection
22 prevention and social determinants of health (e.g., socioeconomic position), and relevant
23 health outcomes (e.g., infection and mortality rates) (10-15). Such health equity issues have
24 been increasingly recognized in Japan by sub-population group, including the mental health
25 issues following the pandemic (e.g., psychological distress, suicide, health-related quality of

1 life) (16-20), whilst the evidence base is still limited in Japan with variations surrounding the
2 socioeconomic perspectives.

3 Multidisciplinary research investigating the mental health impact of the COVID-19
4 pandemic has been also recognized as a research priority since the early stages of the
5 pandemic. In particular, the need for research and interventions that address the
6 psychological, social, and neuro-scientific aspects of the pandemic has been underscored
7 (21). Studies caution against the adverse mental health outcomes during the COVID-19
8 pandemic and the variation of such deterioration by sociodemographic factor, suggesting the
9 negative influence of the COVID-19 pandemic on mental health issues and subsequent health
10 inequity concerns (22-24). Evidence of the adverse mental health impact of the COVID-19
11 pandemic on vulnerable population groups exists globally, and its disproportionate impact
12 among children and adolescents, for instance, is of major concern (22, 25-27). Concerns
13 about COVID-19 itself have been posited as a key factor in the increase in general mental
14 health problems during the pandemic (28, 29). Furthermore, evidence of COVID-19-related
15 anxiety among the lower health status population or the higher risk-taking population group
16 is mixed, suggesting some variations across the specific aspects of anxiety (30-32). Most
17 studies on anxiety about COVID-19, however, employ measures that focus on the disease
18 aspect only (33) and do not distinguish the multiple dimensions that comprise anxiety.

19 As such, there has been increasing recognition of the multidimensionality of COVID-
20 related anxiety during the pandemic, and several new COVID-specific anxiety measures have
21 been developed, tested and validated (33-36). Although the majority of the COVID-specific
22 anxiety measures address a single dimension of anxiety related to COVID-19 as a disease
23 itself, a relevant multidimensional measure was also developed, called the “COVID-19 Stress
24 Scales” that comprise five dimensions and thirty-three specific indicators (36). Following
25 this, the Pandemic Anxiety Scale (PAS) was developed and validated in the United Kingdom

1 (UK) as a feasible and practical scale among surveys, underscoring the multidimensionality
2 of pandemic anxiety using seven indicators. In particular, the PAS differentiates anxiety
3 about COVID-19 infection (i.e., “disease anxiety”) and negative socioeconomic
4 consequences of the pandemic (i.e., “consequence anxiety”) (35), both of which are critical
5 factors in the health and wellbeing of the population, showing differential associations with
6 demographics, social and health factors (e.g., gender, age, and chronic physical health
7 conditions). In addition, a study in Austria validated and employed the PAS, finding a
8 different pattern of pandemic anxiety from the original UK study (37).

9 However, studies that identify and compare distinct pandemic anxiety dimensions are
10 still limited among the general population across generations. The evidence base of anxiety
11 due to the COVID-19 pandemic has mainly focused on the so-called “unidimensional”
12 anxiety that spotlights fear and anxiety about COVID-19 infection and has been statistically
13 validated (35, 36, 38). Another study used a general mental health screening scale (i.e., the
14 Psychological Distress Scale K6) (39). Most of these studies employed a measure that is
15 calculated based on multiple questions or indicators, of which response scores are added to
16 provide a summative score (33, 40, 41). However, relevant measures that assess pandemic
17 anxiety as a latent construct considering reflective indicators are scarce. Furthermore, despite
18 its methodological advantage, evidence is still limited from studies using structural equation
19 modeling (SEM) in the assessment of mental health by measures of depression, anxiety, fear,
20 risk perceptions, and negative emotions during the COVID-19 pandemic (35, 42-47).

21 Therefore, this study aims to examine the associations between socioeconomic
22 position, health-related status and the multiple dimensions of anxiety during the pandemic,
23 namely, disease-related anxiety (i.e., “disease anxiety”) and socioeconomic consequence-
24 related anxiety (i.e., “consequence anxiety”). Using SEM, this study assessed multiple

1 dimensions of pandemic anxiety as a latent outcome measure of mental health issues, which
2 were predicted by a series of interrelated socioeconomic and health-related measures.

3
4

5 **2. Methods**

6
7

8 **(1) Study setting and data**

9

10 This study was conducted in Japan, particularly in the six prefectures where the
11 central government's emergency declaration was in effect and the new COVID-19 infection
12 cases were marked as the highest in the country at the time of study preparation in early
13 2021. The locations were Aichi, Chiba, Kanagawa, Osaka, Saitama, and Tokyo. Ethical
14 approval was obtained from the Research Ethics Committee of the School of Health
15 Management, Keio University, in February 2021.

16 Data were collected in March 2021 from the general working-age populations, aged
17 18 to 59 years. It was approximately a year after the onset of the pandemic in February 2020
18 and the government's direct payment program in the mid-2020. The government's COVID-
19 19 public vaccination program, which started around April 2021, had not yet been launched
20 at the time of the survey. The participants were registered as a survey panel for an
21 international online survey company, Cint Japan, one of the largest online survey companies
22 in Japan. Quota sampling methods were employed according to the national population
23 statistics by age and gender. For data collection, study sample distributions were weighted by
24 the sub-national population statistics of the target prefecture, approximating the distribution
25 of the sub-national population (48). The final study sample comprised 2,764 observations.

26
27

28 **(2) Analytic strategy and measures**

29

30 This study employed latent variable SEM, including both a structural portion (i.e.,
with measured variables) and a measurement portion (i.e., with latent constructs) (49). The
latent variable SEM comprises two latent constructs representing the distinct dimensions of

1 pandemic anxiety related to the COVID-19 pandemic, as described in the subsequent section.
2 The model includes two “endogenous variables”, which appear as dependent variables in one
3 of the equations; and multiple “exogenous variables”, which are never dependent variables
4 and are related to the socioeconomic position and health-related status of the study
5 participant. In the SEM approach in theory, Kline explains that the relationship between
6 variables is examined in terms of “path coefficients” that are indicated as an arrow assuming
7 a potential causal relationship. Thus, “X is a cause of Y” by the conceptual definition of
8 SEM (49).

9

10 ***Endogenous variables – “disease anxiety” and “consequence anxiety”***

11 Endogenous variables represent the anxiety about COVID-19 and the socioeconomic
12 consequences of the COVID-19 pandemic. These anxieties are defined and measured as
13 “disease anxiety” and “consequence anxiety” according to the PAS, which has been validated
14 in the UK (35). The scale comprises seven indicators, including four that are related to
15 COVID-19 infection and three that are related to the socioeconomic consequences of the
16 pandemic. Relevant indicators of “disease anxiety” are reflected in the question that asks
17 about the respondent’s anxiety about the disease itself, including anxiety about the infection
18 of the respondent, infection of family and friends, going out, and transmission of infection to
19 others. Those of “consequence anxiety” inquire about the adverse socioeconomic
20 consequences of the pandemic, including anxiety about missing school/work, reduction of
21 income, and the impact of COVID-19 on the labor market and economy. Respondents
22 reported their level of anxiety about each question on a five-point Likert scale (i.e., strongly
23 disagree, disagree, neither agree nor disagree, agree, strongly agree). A two latent variable
24 structure was confirmed based on the preliminary analysis results, as described in the
25 subsequent section.

1

2 ***Exogenous variables - socioeconomic position and health-related status***

3 Exogenous variables included socioeconomic position and health-related status of the survey
4 participants. Respondents' age was assessed as a continuous and categorical variable in
5 consideration of potential non-linear relationships between age and pandemic anxiety:
6 teenagers aged 18-19, those aged 20-29, aged 30-39, aged 40-49, and aged 50-59 (reference
7 group: ages 30-39). Gender was categorized as male, female, and other for those who
8 reported as "other" or "do not answer" (reference group: female). Education was categorized
9 as "high school or less" or "higher education" (i.e., technical college, 2-year college
10 education or higher) (reference: high school or less). Household income was measured in
11 quintiles, regarding the recent national household annual income data (lowest 20% quintile
12 for Japanese Yen - JPY two million or less; lower 20-40% quantile - JPY 3.42 million or less;
13 middle 40-60% quintile - JPY 5.23 million or less; higher 60-80% quintile - JPY 8.13 million
14 or less; and highest 20% quintile - above JPY 8.13 million) (reference: the highest 20%
15 income quintile) (48). Employment, measured as a binary variable, was asked if they were in
16 paid employment in the last four weeks preceding the survey (reference: not in paid
17 employment). Current schooling was also measured as a binary variable if the respondent
18 was a student or not at the time of the survey (reference: currently not in school). Marital or
19 partnership status was measured as a binary variable if the respondent had a partner
20 regardless of legal status at the time of the survey (reference: not married or having a
21 partner).

22 In addition, the health-related status of a respondent was measured using the
23 international tool developed by EuroQoL and employed internationally as a health outcome
24 measure in public health and health economics research. The five-level EQ-5D version (EQ-
25 5D-5L) consists of two measures: the EQ-5D descriptive system and the EQ-visual analog

1 scale (EQ-VAS). The EQ-5D descriptive system is a health-related quality of life (HRQoL)
2 measure comprising the following five dimensions. Respondents were asked about mobility,
3 self-care, usual activities, pain/discomfort, and anxiety/depression. These questions ask the
4 respondent to select the statement that best describes one's health on the date of the survey
5 for each dimension, and the answer options have five levels (e.g., no problems, slight
6 problems, moderate problems, severe problems, or unable) (50). The HRQoL score is a single
7 cardinal value assigning 0.0 for death and 1.0 for perfect health, and the score was calculated
8 according to the Japanese version's valuation study (range: - 0.025 to 1.000) (51). The EQ-
9 VAS is a measure of self-reported health, and respondents were asked to rate their health
10 status on the date of the survey, indicating 100 for the best health and 0 for the worst health
11 that the respondent could imagine (50).

12 (3) Analytic steps

13 Data analysis was conducted in four steps. First, a descriptive analysis was conducted
14 using STATA 17. Second, the psychometric properties of the Japanese version of the PAS
15 were tested with STATA, using skewness and kurtosis scores for normality and Cronbach's
16 alpha values for internal consistency. Third, factor analyses were conducted using Mplus
17 version 8.7. An exploratory factor analysis (EFA) was utilized to assess the underlying factor
18 structure of the PAS using geomin rotation. The decision on the number of factors to retain
19 was based on an inspection of the eigenvalues and scree plot. A confirmatory factor analysis
20 (CFA) was employed to examine the appropriateness and generalizability of the identified
21 multi-factor structure, which represents the multiple dimensions of pandemic anxiety, as a
22 measurement portion of the SEM. Fourth, SEM was conducted with Mplus version 8.7 to
23 examine the mechanism by which socioeconomic positions and health-related status
24 predicted multidimensional anxiety about the COVID-19 pandemic.
25
26

1 The SEM analyzed two equations simultaneously for the two defined endogenous
2 variables in the model and estimated standardized coefficients, such that the model enabled
3 an examination of the multidimensionality of pandemic anxiety and a comparison of path
4 coefficients in terms of the effect size across exogenous variables of different metrics. These
5 equations separately and simultaneously regressed the two dimensions of pandemic anxiety
6 using polychoric correlations and probit regressions with weighted least squares estimation
7 (in particular, WLSMV weighted least square mean and variance adjusted). In the model, all
8 exogenous variables were designated as covarying because of the potential relatedness among
9 exogenous variables. In addition, the errors/disturbances of the two latent dimensions of
10 anxiety were covarying, as the unobserved aspects of these constructs were likely to be
11 associated with each other (49, 52).

12 Model fit was assessed using the following recommended indices. Root Mean
13 Square Error of Approximation (RMSEA) was examined to be less than 0.06 as a close fit
14 and 0.08 as an acceptable fit. A Comparative Fit Index (CFI) and Tucker-Lewis Index (TLI)
15 were assessed to be 0.95 or higher as a recommended close fit (53).

16 Diagnostic procedures prior to the main analysis included bivariate analyses of
17 selected variables and testing for multicollinearity and normality using STATA 17. These
18 diagnoses have suggested safely ignoring multicollinearity among the exogenous variables,
19 according to the Variance Inflation Factor (VIF) < 10, and assuming normal distributions of
20 the endogenous variable indicator score (according to skewness and kurtosis scores)(54).

21

22 **3. Results**

23 **(1) Descriptive results**

24 Table 1 shows the descriptive statistics for pandemic anxiety, socioeconomic position, and
25 health-related status of the study population. Mean scores of each pandemic anxiety indicator

1 showed some variation across different types of anxiety. The respondents' mean age was
2 approximately 40 and ranged from 18 to 59 years. Around two-thirds had a college-level
3 education or higher and paid employment in the last four weeks preceding the survey. Their
4 household income almost approximated the national income quintile level, with somewhat
5 lower proportions in the second lowest income group. Approximately half were married or
6 had a partner, regardless of legal status. Self-reported health status was 74 points out of 100
7 points on average. The mean HRQoL score is 0.897, against the highest possible score of
8 1.000.

9 [INSERT TABLE 1 HERE]

10

11 **(2) Psychometric properties of the Pandemic Anxiety Scale**

12 Skewness and kurtosis scores for each item of the PAS indicated normality, using a
13 conventional guideline of values ± 3 (54). Cronbach's alpha values suggested relatively high
14 internal consistency among the seven items of the PAS (scale reliability coefficient=0.861).

15

16 **(3) Factor analysis results**

17 The EFA and CFA results suggested a two-factor structure of pandemic anxiety – “disease
18 anxiety” and “consequence anxiety”. Table 2 reports the factor loadings of each indicator on
19 the respective factors from CFA, with the loading of the first indicator being set free and the
20 variance of the latent factor set to one. The EFA indicated that the first factor and an
21 eigenvalue of 4.164, and the second had an eigenvalue 0.973, which is narrowly below the
22 conventional cut-off of 1.0. The third factor had an eigenvalue of 0.649. Based on these
23 results and the patterns of factor loadings, we chose to retain the two-factor structure in
24 subsequent analyses. The two-factor structure had better model fit indices than the one-factor
25 structure in terms of CFI/TLI and RMSEA (55). The CFA results confirmed the two-factor

1 structure with model fit indices that were close or acceptable according to the aforementioned
2 thresholds (e.g., CFI/TLI) (Table 2), in support of the appropriateness and generalizability of
3 the measurement portion of the SEM. The two defined factors had a standardized correlation
4 of 0.71 with statistical significance.

5 [INSERT TABLE 2 HERE]

6

7 (4) SEM results

8 The results of the final adjusted SEM are shown in Table 3 and Figure 1, which report the
9 standardized regression coefficients. The coefficients are indicated as a single-headed arrow
10 pointing from X (cause) to Y (effect) in the figure, assuming a potential causal relationship
11 by the traditional conceptual definition of the SEM approach in this analysis (49). The model
12 fit indices show that the model fit the data well (CFI=0.959; TLI=0.942; RMSEA=0.071).
13 Sensitivity analysis results are reported in a supplement file (Supplement 1).

14

15 [INSERT TABLE 3 AND FIGURE 1 HERE]

16

17 The regression coefficients of the exogenous variables predicting **disease anxiety** are
18 reported in the column 1 of Table 3. Gender differences were statistically significant, and
19 males reported the lower level of disease anxiety than females ($b = -0.126$). The oldest age
20 group age 50-59 reported the lower disease anxiety than those aged 30-39, whilst those age
21 40-49 reported the lower disease anxiety at borderline significance. Unmarried respondents
22 and those with higher quality of life scores also reported lower disease anxiety. The rest of
23 the exogenous variables, however, did show a statistically significant association with disease
24 anxiety, including education, current schooling, household income, paid employment, and
25 self-reported health. Comparisons of the standardized coefficients suggest that gender,

1 marital relationship and age differences reflect the largest effect sizes in predicting disease
2 anxiety among the selected exogenous variables.

3 Further, the standardized coefficients of exogenous variables predicting **consequence**
4 **anxiety** are reported in the column 2 of Table 3. Compared to the results for disease anxiety,
5 there are a greater number of exogenous variables with statistically significant coefficients
6 for consequence anxiety. Specifically, gender differences were significant, with males
7 reporting the lower level of consequence anxiety than females ($b = -0.079$). Respondents aged
8 50-59 reported lower consequence anxiety compared to those aged 30-39 ($b = 0.085$).
9 Relative to the highest 20% income quintiles, the rest of the income quintile groups reported
10 higher consequence anxiety, although no clear gradient patterns were found across groups.
11 Paid employment and marital relationship also had a positive association. In addition, the
12 health-related Quality of Life (HRQoL) showed a negative association, indicating that the
13 better the HRQoL, the lower the consequence anxiety. However, the remaining exogenous
14 variables, including education, current schooling and the self-reported health status, did not
15 show a statistically significant relationship with consequence anxiety. Among the selected
16 exogenous variables, differences in paid employment, income, and marital relationship
17 reflected the larger effect sizes in predicting consequence anxiety.

18

19 **4. Discussion**

20 This study examined the associations linking socioeconomic positions and health-related
21 status with the multidimensions of anxiety related to COVID-19 infection and adverse
22 socioeconomic consequences of the pandemic, among the Japanese working population aged
23 18-59. Using a SEM approach, this analysis provides evidence of the two interrelated yet
24 distinct dimensions of pandemic anxiety, as they are related to and likely to be influenced by
25 a series of interrelated individual-level socioeconomic and health-related factors.

1 Key findings from this study emphasize the concern about the disproportionate
2 socioeconomic impact of the pandemic and consequently, the widening health inequity
3 following the pandemic, of which the trend and transition are likely to differ by setting. First,
4 substantial gender differences exist in terms of the two pandemic anxiety dimensions. This
5 gender pattern is also consistent with findings from the UK, showing higher anxiety among
6 females than males in the both anxiety dimensions (35). This pandemic anxiety pattern may
7 reflect the general concern in Japanese society, together with several other societal concerns
8 that demonstrate gender differences, such as females being disproportionately affected by the
9 adverse socioeconomic consequences of the pandemic compared to males (3, 16, 56). In
10 addition, the lower disease anxiety among males than females may have an influence on, at
11 least in part, the relatively higher proportion of COVID-19 infection among males, although
12 the gender pattern undergoes transitions and variations by context (11, 57-59).

13 Second, this study shows unexpected results that are inconsistent with existing
14 evidence. Self-reported health status showed no significant relationship with disease anxiety
15 or consequence anxiety, whilst the HRQoL was negatively associated with both dimensions;
16 that is, the higher the QoL, the lower the pandemic anxiety. This association should be
17 further investigated in future studies, given this inconsistency with the existing literature on
18 mental health issues among less healthy groups during the pandemic (e.g., fear, anxiety, and
19 depression) (30-32). In addition, while the UK study demonstrated negative associations
20 between household income and the both dimensions of pandemic anxiety (35), unique
21 evidence is also shown from this Japanese study. Specifically, although significant negative
22 associations between consequence anxiety and economically disadvantaged groups were
23 somewhat foreseen in Japan – as concerned globally that the lower socioeconomic groups are
24 disproportionately affected during the pandemic (14, 15, 56, 60) – this finding suggests that
25 those who are not in paid employment report lower consequence anxiety. Emergency

1 financial support programs by both central and regional governments, as well as non-profit
2 organizations, might have possibly mitigated, at least in part, the high-level consequence
3 anxiety among the financially-vulnerable population in Japan. The potential of the effect of
4 these emergency financial schemes, targeting both individuals and enterprises, on quality of
5 life, mental health, and suicide prevention was also discussed in Japanese studies, as a
6 possible important mitigating factor (17, 20). In consideration of mid- and long-term
7 socioeconomic consequences of the pandemic, such financial and social support schemes
8 beyond short-term emergency schemes should be strengthened widely, as these measures are
9 recommended and shown to be effective (61-63).

10 Third, this study shows unexpected patterns of generational differences in pandemic
11 anxiety. In this Japanese model, older generations reported lower anxiety in both dimensions,
12 disease anxiety and consequence anxiety. Higher consequence anxiety among younger
13 generations was somewhat foreseen, in accordance with the descriptive evidence and general
14 societal concerns that younger generations have been more severely and negatively affected
15 in the labor market compared to older generations who are relatively more stable in
16 employment status (16). This Japanese finding is not consistent with the UK study finding no
17 significant age differences regarding pandemic anxiety among the working-age adult
18 population. Significant age differences were found, however, among UK adolescents (35).
19 Considering the prolonged adverse socioeconomic consequences in Japan to date, a
20 subsequent longitudinal analysis should further investigate generational disparities in anxiety
21 related to the COVID-19 pandemic.

22 There are some limitations to this study. First, this was a cross-sectional survey; thus,
23 causal inference on the hypothesized pathway is tentative, according to the controversy
24 surrounding SEM (64). Second, the study sample was drawn from selected geographic areas
25 and was not nationally representative, while the sample was randomly drawn from the survey

1 panel approximating sub-national demographic patterns. Third, there may have been
2 unobserved variables in the study that influenced mental health issues in general and the
3 pandemic anxiety in particular, while the significant correlation of disturbances between the
4 two pandemic anxiety dimensions suggest that their unobserved aspects are interrelated.
5 Fourth, given the shifting of COVID-19 infection patterns and government policy, relevant
6 pandemic anxiety indicators and questions must be reviewed as appropriate (37). Fifth, future
7 investigations should consider further in-depth analyses including mediation and multigroup
8 analyses (Supplement 2). Finally, it should be noted that the timing of relevant studies and
9 analytic model structures differ across countries (e.g., Austria, Japan, and the UK). Owing to
10 the differential pattern of COVID-19 infection and transition globally, and the
11 methodological features and differences across the settings (e.g., differences in data
12 collection strategies and measures), the comparative inference of pandemic anxiety across
13 contexts requires careful attention.

14 Despite these limitations, this study examined health inequity related to mental health
15 issues and pandemic anxiety, in consideration of the multidimensionality of anxiety during
16 the COVID-19 pandemic. This evidence from a latent variable SEM underscores the two
17 distinct and interrelated anxiety dimensions, suggesting a unique pattern and predictors of
18 each dimension, as well as substantial concerns about mental health issues related to the
19 socioeconomic consequences of the pandemic. This distinction between the two anxiety
20 dimensions highlights the more substantial disparities with mental health issues, which are
21 likely to be due to the socioeconomic consequences of the pandemic as a key driver of
22 widening health and social gaps in Japan. Compared to European studies, this unique finding
23 from Japan suggests critical and possibly high potential mitigating measures to buffer the
24 serious socioeconomic impact among the most vulnerable populations. Relevant social and
25 economic support policies and programs need to be warranted for vulnerable populations,

- 1 beyond the short-term emergency funding scheme in the process of transition and recovery
- 2 from the COVID-19 pandemic. Due to the persistent disproportionate socioeconomic impact
- 3 of the pandemic on vulnerable populations globally, multidisciplinary research on mental
- 4 health issues and quality of life remains an important research agenda in exploring
- 5 socioeconomic measures in context, towards addressing inequity concerns.

Journal Pre-proof

1 Reference

- 2 1. McKibbin W, Fernando R. The global macroeconomic impacts of COVID-19: Seven
3 scenarios. *Asian Economic Papers*. 2021;20(2):1-30.
- 4 2. Ceylan RF, Ozkan B, Mulazimogullari E. Historical evidence for economic effects of
5 COVID-19. Springer; 2020. p. 817-23.
- 6 3. Churchill B. COVID - 19 and the immediate impact on young people and
7 employment in Australia: A gendered analysis. *Gender, Work & Organization*.
8 2021;28(2):783-94.
- 9 4. Egger D, Miguel E, Warren SS, Shenoy A, Collins E, Karlan D, et al. Falling living
10 standards during the COVID-19 crisis: Quantitative evidence from nine developing countries.
11 *Science advances*. 2021;7(6):eabe0997.
- 12 5. Ahmed F, Ahmed Ne, Pissarides C, Stiglitz J. Why inequality could spread COVID-19.
13 *The Lancet Public Health*. 2020;5(5):e240.
- 14 6. Loayza N. Costs and trade-offs in the fight against the COVID-19 pandemic: A
15 developing country perspective. *World Bank Research and Policy Briefs*. 2020(148535).
- 16 7. World Health Organization (WHO). Social Determinants of Health [Available from:
17 https://www.who.int/health-topics/social-determinants-of-health#tab=tab_3 (Accessed on
18 20 July 2022).
- 19 8. Patel V, Saxena S, Lund C, Thornicroft G, Baingana F, Bolton P, et al. The Lancet
20 Commission on global mental health and sustainable development. *The lancet*.
21 2018;392(10157):1553-98.
- 22 9. Marmot M, Friel S, Bell R, Houweling TA, Taylor S, Health CoSDo. Closing the gap in a
23 generation: health equity through action on the social determinants of health. *The lancet*.
24 2008;372(9650):1661-9.
- 25 10. World Health Organization (WHO). Vaccine equity [Available from:
26 <https://www.who.int/campaigns/vaccine-equity> (Accessed on 21 July 2022).
- 27 11. Khalatbari-Soltani S, Cumming RC, Delpierre C, Kelly-Irving M. Importance of
28 collecting data on socioeconomic determinants from the early stage of the COVID-19
29 outbreak onwards. *J Epidemiol Community Health*. 2020;74(8):620-3.
- 30 12. Forbes H, Morton CE, Bacon S, McDonald HI, Minassian C, Brown JP, et al.
31 Association between living with children and outcomes from covid-19: OpenSAFELY cohort
32 study of 12 million adults in England. *Bmj*. 2021;372:n628.
- 33 13. Mathur R, Rentsch CT, Morton CE, Hulme WJ, Schultze A, MacKenna B, et al. Ethnic
34 differences in SARS-CoV-2 infection and COVID-19-related hospitalisation, intensive care
35 unit admission, and death in 17 million adults in England: an observational cohort study
36 using the OpenSAFELY platform. *Lancet*. 2021;397(10286):1711-24.
- 37 14. Williamson EJ, Walker AJ, Bhaskaran K, Bacon S, Bates C, Morton CE, et al. Factors
38 associated with COVID-19-related death using OpenSAFELY. *Nature*. 2020;584(7821):430-6.
- 39 15. Baena-Díez JM, Barroso M, Cordeiro-Coelho SI, Díaz JL, Grau M. Impact of COVID-19
40 outbreak by income: hitting hardest the most deprived. *J Public Health (Oxf)*.
41 2020;42(4):698-703.
- 42 16. Ueda M, Nordström R, Matsubayashi T. Suicide and mental health during the COVID-
43 19 pandemic in Japan. *Journal of public health (Oxford, England)*. 2020.
- 44 17. Tanaka T, Okamoto S. Increase in suicide following an initial decline during the
45 COVID-19 pandemic in Japan. *Nature human behaviour*. 2021;5(2):229-38.

- 1 18. Yoshioka T, Okubo R, Tabuchi T, Odani S, Shinozaki T, Tsugawa Y. Factors associated
2 with serious psychological distress during the COVID-19 pandemic in Japan: a nationwide
3 cross-sectional internet-based study. *BMJ open*. 2021;11(7):e051115-e.
- 4 19. Yoshikawa Y, Kawachi I. Association of Socioeconomic Characteristics With
5 Disparities in COVID-19 Outcomes in Japan. *JAMA Netw Open*. 2021;4(7):e2117060.
- 6 20. Ikeda T, Igarashi A, Odani S, Murakami M, Tabuchi T. Health-related quality of life
7 during COVID-19 pandemic: assessing impacts of job loss and financial support programs in
8 Japan. *Applied research in quality of life*. 2021:1-17.
- 9 21. Holmes EA, O'Connor RC, Perry VH, Tracey I, Wessely S, Arseneault L, et al.
10 Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental
11 health science. *The Lancet Psychiatry*. 2020;7(6):547-60.
- 12 22. Patel K, Robertson E, Kwong ASF, Griffith GJ, Willan K, Green MJ, et al. Psychological
13 Distress Before and During the COVID-19 Pandemic Among Adults in the United Kingdom:
14 Coordinated Analyses of 11 Longitudinal Studies. *medRxiv*. 2022:2021.10.22.21265368.
- 15 23. Ettman CK, Cohen GH, Abdalla SM, Sampson L, Trinquart L, Castrucci BC, et al.
16 Persistent depressive symptoms during COVID-19: a national, population-representative,
17 longitudinal study of US adults. *The Lancet Regional Health-Americas*. 2022;5:100091.
- 18 24. Pierce M, Hope H, Ford T, Hatch S, Hotopf M, John A, et al. Mental health before and
19 during the COVID-19 pandemic: a longitudinal probability sample survey of the UK
20 population. *The Lancet Psychiatry*. 2020;7(10):883-92.
- 21 25. Meherali S, Punjani N, Louie-Poon S, Abdul Rahim K, Das JK, Salam RA, et al. Mental
22 Health of Children and Adolescents Amidst COVID-19 and Past Pandemics: A Rapid
23 Systematic Review. *Int J Environ Res Public Health*. 2021;18(7).
- 24 26. Panchal U, Salazar de Pablo G, Franco M, Moreno C, Parellada M, Arango C, et al. The
25 impact of COVID-19 lockdown on child and adolescent mental health: systematic review. *Eur*
26 *Child Adolesc Psychiatry*. 2021:1-27.
- 27 27. Samji H, Wu J, Ladak A, Vossen C, Stewart E, Dove N, et al. Review: Mental health
28 impacts of the COVID-19 pandemic on children and youth - a systematic review. *Child*
29 *Adolesc Ment Health*. 2021.
- 30 28. Labrague LJ, de Los Santos JAA. Fear of Covid - 19, psychological distress, work
31 satisfaction and turnover intention among frontline nurses. *Journal of nursing management*.
32 2021;29(3):395-403.
- 33 29. Mailliez M, Griffiths MD, Carre A. Validation of the french version of the fear of
34 covid-19 scale and its associations with depression, anxiety, and differential emotions.
35 *International journal of mental health and addiction*. 2021:1-15.
- 36 30. Musche V, Kohler H, Bäuerle A, Schweda A, Weismüller B, Fink M, et al. COVID-19-
37 Related Fear, Risk Perception, and Safety Behavior in Individuals with Diabetes. *Healthcare*
38 *(Basel)*. 2021;9(4).
- 39 31. Kohler H, Bäuerle A, Schweda A, Weismüller B, Fink M, Musche V, et al. Increased
40 COVID-19-related fear and subjective risk perception regarding COVID-19 affects behavior in
41 individuals with internal high-risk diseases. *J Prim Care Community Health*.
42 2021;12:2150132721996898.
- 43 32. Akgor U, Fadiloglu E, Soyak B, Unal C, Cagan M, Temiz BE, et al. Anxiety, depression
44 and concerns of pregnant women during the COVID-19 pandemic. *Arch Gynecol Obstet*.
45 2021;304(1):125-30.

- 1 33. Ahorsu DK, Lin C-Y, Imani V, Saffari M, Griffiths MD, Pakpour AH. The fear of COVID-
2 19 scale: development and initial validation. *International journal of mental health and*
3 *addiction*. 2020;1-9.
- 4 34. Bernardo ABI, Mendoza NB, Simon PD, Cunanan ALP, Dizon JIWT, Tarroja MCH, et al.
5 Coronavirus Pandemic Anxiety Scale (CPAS-11): development and initial validation. *Current*
6 *Psychology*. 2020.
- 7 35. McElroy E, Patalay P, Moltrecht B, Shevlin M, Shum A, Creswell C, et al. Demographic
8 and health factors associated with pandemic anxiety in the context of COVID - 19. *British*
9 *Journal of Health Psychology*. 2020;25(4):934-44.
- 10 36. Taylor S, Landry CA, Paluszek MM, Fergus TA, McKay D, Asmundson GJ.
11 Development and initial validation of the COVID Stress Scales. *Journal of Anxiety Disorders*.
12 2020;72:102232.
- 13 37. Kubb C, Foran HM. Measuring COVID-19 related anxiety in parents: Psychometric
14 comparison of four different inventories. *JMIR mental health*. 2020;7(12):e24507.
- 15 38. Lin CY, Hou WL, Mamun MA, Aparecido da Silva J, Broche - Pérez Y, Ullah I, et al.
16 Fear of COVID - 19 Scale (FCV - 19S) across countries: Measurement invariance issues.
17 *Nursing open*. 2021;8(4):1892-908.
- 18 39. Kessler RC, Andrews G, Colpe LJ, Hiripi E, Mroczek DK, Normand S-L, et al. Short
19 screening scales to monitor population prevalences and trends in non-specific psychological
20 distress. *Psychological medicine*. 2002;32(6):959-76.
- 21 40. Masuyama A, Shinkawa H, Kubo T. Validation and psychometric properties of the
22 Japanese version of the fear of COVID-19 scale among adolescents. *International Journal of*
23 *Mental Health and Addiction*. 2020:1-11.
- 24 41. Harper CA, Satchell LP, Fido D, Latzman RD. Functional fear predicts public health
25 compliance in the COVID-19 pandemic. *International journal of mental health and addiction*.
26 2021;19(5):1875-88.
- 27 42. Bechard LE, Bergelt M, Neudorf B, DeSouza TC, Middleton LE. Using the Health Belief
28 Model to Understand Age Differences in Perceptions and Responses to the COVID-19
29 Pandemic. *Frontiers in Psychology*. 2021;12(1216).
- 30 43. Li JB, Yang A, Dou K, Wang LX, Zhang MC, Lin XQ. Chinese public's knowledge,
31 perceived severity, and perceived controllability of COVID-19 and their associations with
32 emotional and behavioural reactions, social participation, and precautionary behaviour: a
33 national survey. *BMC Public Health*. 2020;20(1):1589.
- 34 44. Ding Y, Xu J, Huang S, Li P, Lu C, Xie S. Risk Perception and Depression in Public
35 Health Crises: Evidence from the COVID-19 Crisis in China. *Int J Environ Res Public Health*.
36 2020;17(16).
- 37 45. Lu P, Kong D, Shelley M. Risk Perception, Preventive Behavior, and Medical Care
38 Avoidance among American Older Adults During the COVID-19 Pandemic. *J Aging Health*.
39 2021;33(7-8):577-84.
- 40 46. Mahamid FA, Veronese G, Bdier D. Fear of coronavirus (COVID-19) and mental health
41 outcomes in Palestine: The mediating role of social support. *Curr Psychol*. 2021:1-10.
- 42 47. Mamun MA, Sakib N, Gozal D, Bhuiyan AI, Hossain S, Bodrud-Doza M, et al. The
43 COVID-19 pandemic and serious psychological consequences in Bangladesh: A population-
44 based nationwide study. *J Affect Disord*. 2021;279:462-72.

- 1 48. Japanese Ministry of Internal Affairs and Communications - Bureau of Statistics.
2 Statistics of Japan 2022 [Available from: <https://www.e-stat.go.jp/en> (Accessed on 1 March
3 2022)].
- 4 49. Kline RB. Principles and practice of structural equation modeling (Fourth edition):
5 Guilford publications; 2016.
- 6 50. The EuroQoL Group. EQ-5D-5L | About 2021 [Available from: <https://euroqol.org/eq-5d-instruments/eq-5d-5l-about/> (Accessed on 1 March 2022).
7
- 8 51. IKEDA S, SHIROIWA, T., IGARASHI, A., NOTO, S., FUKUDA, T., SAITO, S.,
9 SHIMOZUMA, K. Developing a Japanese version of the EQ-5D-5L value set. Journal of the
10 National Institute of Public Health (in Japanese). 2015;64(1):47-55.
- 11 52. Muthen L, Muthen B. Mplus Statistical Analysis With Latent Variables User's Guide
12 (Version 8). 2017.
- 13 53. Hu L-t, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis:
14 Conventional criteria versus new alternatives. Structural Equation Modeling: A
15 Multidisciplinary Journal. 1999;6(1):1-55.
- 16 54. Hair JF, Black WC, Babin BJ, Anderson RE, Tatham RL. Multivariate data analysis (Vol.
17 5. 3). Prentice hall, Upper Saddle River, NJ; 1998.
- 18 55. Pett M, Lackey N, Sullivan J. The use of factor analysis for instrument development in
19 health care research. Thousand Oaks, CA: Sage; 2003.
- 20 56. Han J, Meyer BD, Sullivan JX. Income and Poverty in the COVID-19 Pandemic.
21 Brookings Papers on Economic Activity. 2020:85-118.
- 22 57. Jin J-M, Bai P, He W, Wu F, Liu X-F, Han D-M, et al. Gender Differences in Patients
23 With COVID-19: Focus on Severity and Mortality. Frontiers in Public Health. 2020;8.
- 24 58. UN Women. COVID-19 and gender monitor 2020 [Available from:
25 <https://data.unwomen.org/resources/covid-19-and-gender-monitor> (Accessed on 1 March
26 2022)].
- 27 59. Ministry of health labour and welfare in Japan. Visualizing the data: information on
28 COVID-19 infections 2022 [Available from: <https://covid19.mhlw.go.jp/en/> (Accessd on
29 December 31, 2021)].
- 30 60. Almeida V, Barrios S, Christl M, De Poli S, Tumino A, van der Wielen W. The impact of
31 covid-19 on households income in the EU. The Journal of Economic Inequality.
32 2021;19(3):413-31.
- 33 61. Pinto AD, Perri M, Pedersen CL, Aratangy T, Hapsari AP, Hwang SW. Exploring
34 different methods to evaluate the impact of basic income interventions: a systematic
35 review. International journal for equity in health. 2021;20(1):1-20.
- 36 62. Tsai J, Huang M, Montgomery AE, Elbogen EB. Receipt, Spending, and Clinical
37 Correlates of the Economic Impact Payment Among Middle- and Low-Income U.S. Adults.
38 Psychiatr Serv. 2021;72(12):1377-84.
- 39 63. Hensher M. Covid-19, unemployment, and health: time for deeper solutions? BMJ.
40 2020;371:m3687.
- 41 64. Bollen KA, Pearl J. Eight myths about causality and structural equation models.
42 Handbook of causal analysis for social research: Springer; 2013. p. 301-28.
- 43

Table 1: Descriptive statistics of the study population in Japan (n=2,764)

Variables	Frequency	Mean (standard deviation) or Proportion/Percentage
Pandemic Anxiety Scale (PAS) with Likert scale (0=strongly disagree; 4=strongly agree)		
I'm worried that I will catch COVID-19 ^[c]	2764	2.55 (1.10)
I'm worried that family and friends will catch COVID-19 ^[c]	2764	2.74 (1.05)
I'm afraid to leave the house right now ^[c]	2764	1.81 (1.11)
I'm worried I might transmit the infection to someone else ^[c]	2764	2.30 (1.13)
I'm worried about missing school/work ^[c]	2764	1.97 (1.22)
I'm worried about the amount of money we have coming in ^[c]	2764	2.44 (1.17)
I'm worried about the long-term impact this will have on my job prospects and the economy ^[c]	2764	2.60 (1.10)
Socioeconomic position/demographics		
Age ^[c]	2764	38.80 (12.25)
Age 18-19	236	8.54%
Age 20-29	521	18.85%
Age 30-39	590	21.35%
Age 40-49	774	28.00%
Age 50-59	643	23.26%
Gender		
Male	1371	49.60%
Female	1372	49.64%
Other/do not answer	21	0.76%
Education		
Lower education (high school or lower)	970	35.09%
Higher education (technical college, 2-year college or higher)	1794	64.91%
Current schooling		
Currently in schooling	329	11.90%
Not in schooling	2435	88.10%
Paid employment (in the last 28 days)		
In paid employment	1914	69.25%
Not in paid employment	850	30.75%
Household income quintile		
Lowest 20% income	541	19.57%
Lower 20-40% income	372	13.46%
Middle 40-60% income	577	20.88%
Higher 60-80% income	661	23.91%
Highest 20% income	613	22.18%
Marital status/partner		
Currently married or have a partner	1279	46.27%
Not married or do not have a partner	1485	53.73%
Health-related status		
Self-reported health status ^[c] (100=best health; 0=worst health)	2764	74.29 (21.64)
Health-related quality of life (HRQoL) ^[c] (1=perfect health; 0=death)	2764	0.897 (0.167)

Note: [c]=continuous variables. Among those who are not in paid employment (850 observations), current students are 161 observations; females are 574 observations.

Table 2: Factor loadings and model fit statistics of the Pandemic Anxiety Scale among the study population in Japan (n=2,764)

	[Factor 1] Disease anxiety Factor loading	[Factor 2] Consequence anxiety Factor loading
Q1 I'm worried that I will catch COVID-19	0.901	N.A.
Q2 I'm worried that family and friends will catch COVID-19	0.909	N.A.
Q3 I'm afraid to leave the house right now	0.693	N.A.
Q4 I'm worried I might transmit the infection to someone else	0.788	N.A.
Q5 I'm worried about missing school/work	N.A.	0.879
Q6 I'm worried about the amount of money we have coming in	N.A.	0.997
Q7 I'm worried about the long-term impact this will have on my job prospects and the economy	N.A.	0.950
Model fit statistics		
CFI	0.957	
TLI	0.931	
RMSEA (90% Confidence Interval)	0.182 (0.173 - 0.191)	


Note: Factor loadings and model fit statistics are reported from a two-factor Confirmatory Factor Analysis (CFA), with the loading of the first indicator being set free and the variance of latent factor set one.

Table 3. Standardized path coefficients of the latent variable SEM on Pandemic Anxiety Scale (Japanese n=2,764)

Exogenous variables (X: independent variables in the equation):	Endogenous variables (Y: dependent variables in the equation):			
	[Column 1] Disease anxiety		[Column 2] Consequence anxiety	
	coefficient	p-value	coefficient	p-value
Male	-0.126	0.000	-0.079	0.000
Other gender	0.019	0.261	-0.005	0.771
Age 18-19	0.033	0.211	0.012	0.671
Age 20-29	0.029	0.247	0.012	0.653
Age 40-49	-0.048	0.063	-0.018	0.498
Age 50-59	-0.091	0.000	-0.085	0.002
Higher education	0.007	0.729	-0.012	0.596
Household income: the lowest 20%	-0.010	0.693	0.070	0.010
Household income: the lower 20-40%	-0.033	0.171	0.067	0.009
Household income: the middle 40-60%	0.007	0.784	0.101	0.000
Household income: the higher 60-80%	0.019	0.448	0.102	0.000
Paid employment	0.003	0.908	0.204	0.000
Current schooling	0.032	0.190	0.042	0.111
Marital relationship	0.100	0.000	0.102	0.000
Self-reported health (EQ-VAS)	0.022	0.326	-0.012	0.622
Health-related quality of life (HRQoL)	-0.073	0.002	-0.064	0.011
Model fit statistics				
CFI	0.959			
TLI	0.942			
RMSEA (90% Confidence Interval)	0.071 (0.067-0.074)			

Reference groups: gender=female; age=age 30-39; education=high-school or lower; household income=the highest 20% income quintile; employment=not in paid employment; schooling=not currently schooling; marital relationship=not married or do not have a partner.

Figure 1: Diagram of the latent variable SEM on Pandemic Anxiety Scale (Japanese n=2,764)



Note:

- 1) Variable labels in the SEM represent the following: male=male; genoth=other gender; hiedu=higher education; iq1=income quintile the lowest 20%; iq2=income quintile the second lowest 20-40%; iq3=income quintile the middle 40-60%; iq4=income quintile the higher 60-80%; eqvas=EQ-VAS; qol=HRQoL; pemp=paid employment; sch=schooling; married=married; g10=age 18-19; g20=age 20-29; g40=age 40-49; g50=age 50-59; F1=factor 1 on disease anxiety; F2=factor 2 on consequence anxiety; pas1-7=indicators per each PAS question.
- 2) Arrows (from left to right) indicate the relationship between the concerned variables in the tested SEM. An arrow between the two factors indicates factor correlations. All exogenous variables are covarying each other.

Supplement 1. Sensitivity Analysis: Standardized path coefficients of the latent variable SEM on Pandemic Anxiety Scale (Japanese sub-samples who are not currently schooling n=2,435)

Exogenous variables (X: independent variables in the equation):	Endogenous variables (Y: dependent variables in the equation):			
	[Column 1]		[Column 2]	
	Disease anxiety		Consequence anxiety	
	coefficient	p-value	coefficient	p-value
Male	-0.135	0.000	-0.081	0.000
Other gender	0.018	0.324	0.004	0.812
Age 18-19	0.020	0.343	0.013	0.557
Age 20-29	0.025	0.299	0.005	0.852
Age 40-49	-0.045	0.091	-0.012	0.664
Age 50-59	-0.092	0.001	-0.083	0.003
Higher education	-0.006	0.797	-0.013	0.565
Household income: the lowest 20%	-0.021	0.445	0.083	0.004
Household income: the lower 20-40%	-0.047	0.068	0.077	0.005
Household income: the middle 40-60%	0.007	0.800	0.113	0.000
Household income: the higher 60-80%	0.020	0.450	0.125	0.000
Paid employment	0.006	0.811	0.227	0.000
Marital relationship	0.096	0.000	0.105	0.000
Self-reported health (EQ-VAS)	-0.006	0.791	-0.039	0.124
Health-related quality of life (HRQoL)	-0.067	0.006	-0.067	0.012
Model fit statistics				
CFI	0.958			
TLI	0.940			
RMSEA (90% Confidence Interval)	0.076 (0.072-0.080)			

Reference groups: gender=female; age=age 30-39; education=high-school or lower; household income=the highest 20% income quintile; employment=not in paid employment; schooling=not currently schooling; marital relationship=not married or do not have a partner.

Supplement 2: Moderation effect results from the latent variable SEM on Pandemic Anxiety Scale (Japanese n=2,764)

	Endogenous variables			
	(Y: dependent variables in the equation):			
	[Column 1]		[Column 2]	
	Disease anxiety		Consequence anxiety	
Interaction terms	coefficient	p-value	coefficient	p-value
(X: independent variables in the equation):				
gender*education	0.018	0.650	0.064	0.331
employment*marital status	0.026	0.544	0.065	0.134
gender*employment	0.025	0.579	-0.029	0.539

Note: Standardized coefficients are reported. Each interaction term was included and tested respectively in the base model (Table 3 and Figure 1).

Table 1: Descriptive statistics of the study population in Japan (n=2,764)

Variables	Frequency	Mean (standard deviation) or Proportion/Percentage
Pandemic Anxiety Scale (PAS) with Likert scale (0=strongly disagree; 4=strongly agree)		
I'm worried that I will catch COVID-19 ^[c]	2764	2.55 (1.10)
I'm worried that family and friends will catch COVID-19 ^[c]	2764	2.74 (1.05)
I'm afraid to leave the house right now ^[c]	2764	1.81 (1.11)
I'm worried I might transmit the infection to someone else ^[c]	2764	2.30 (1.13)
I'm worried about missing school/work ^[c]	2764	1.97 (1.22)
I'm worried about the amount of money we have coming in ^[c]	2764	2.44 (1.17)
I'm worried about the long-term impact this will have on my job prospects and the economy ^[c]	2764	2.60 (1.10)
Socioeconomic position/demographics		
Age ^[c]	2764	38.80 (12.25)
Age 18-19	236	8.54%
Age 20-29	521	18.85%
Age 30-39	590	21.35%
Age 40-49	774	28.00%
Age 50-59	643	23.26%
Gender		
Male	1371	49.60%
Female	1372	49.64%
Other/do not answer	21	0.76%
Education		
Lower education (high school or lower)	970	35.09%
Higher education (technical college, 2-year college or higher)	1794	64.91%
Current schooling		
Currently in schooling	329	11.90%
Not in schooling	2435	88.10%
Paid employment (in the last 28 days)		
In paid employment	1914	69.25%
Not in paid employment	850	30.75%
Household income quintile		
Lowest 20% income	541	19.57%
Lower 20-40% income	372	13.46%
Middle 40-60% income	577	20.88%
Higher 60-80% income	661	23.91%
Highest 20% income	613	22.18%
Marital status/partner		
Currently married or have a partner	1279	46.27%
Not married or do not have a partner	1485	53.73%
Health-related status		
Self-reported health status ^[c] (100=best health; 0=worst health)	2764	74.29 (21.64)
Health-related quality of life (HRQoL) ^[c] (1=perfect health; 0=death)	2764	0.897 (0.167)

Note: [c]=continuous variables. Among those who are not in paid employment (850 observations), current students are 161 observations; females are 574 observations.

Table 2: Factor loadings and model fit statistics of the Pandemic Anxiety Scale among the study population in Japan (n=2,764)

	[Factor 1] Disease anxiety	[Factor 2] Consequence anxiety
	Factor loading	Factor loading
Q1 I'm worried that I will catch COVID-19	0.901	N.A.
Q2 I'm worried that family and friends will catch COVID-19	0.909	N.A.
Q3 I'm afraid to leave the house right now	0.693	N.A.
Q4 I'm worried I might transmit the infection to someone else	0.788	N.A.
Q5 I'm worried about missing school/work	N.A.	0.879
Q6 I'm worried about the amount of money we have coming in	N.A.	0.997
Q7 I'm worried about the long-term impact this will have on my job prospects and the economy	N.A.	0.950
Model fit statistics		
CFI	0.957	
TLI	0.931	
RMSEA (90% Confidence Interval)	0.182 (0.173 - 0.191)	


Note: Factor loadings and model fit statistics are reported from a two-factor Confirmatory Factor Analysis (CFA), with the loading of the first indicator being set free and the variance of latent factor set one.

Table 3. Standardized path coefficients of the latent variable SEM on Pandemic Anxiety Scale (Japanese n=2,764)

Exogenous variables (X: independent variables in the equation):	Endogenous variables (Y: dependent variables in the equation):			
	[Column 1] Disease anxiety		[Column 2] Consequence anxiety	
	coefficient	p-value	coefficient	p-value
Male	-0.126	0.000	-0.079	0.000
Other gender	0.019	0.261	-0.005	0.771
Age 18-19	0.033	0.211	0.012	0.671
Age 20-29	0.029	0.247	0.012	0.653
Age 40-49	-0.048	0.063	-0.018	0.498
Age 50-59	-0.091	0.000	-0.085	0.002
Higher education	0.007	0.729	-0.012	0.596
Household income: the lowest 20%	-0.010	0.693	0.070	0.010
Household income: the lower 20-40%	-0.033	0.171	0.067	0.009
Household income: the middle 40-60%	0.007	0.784	0.101	0.000
Household income: the higher 60-80%	0.019	0.448	0.102	0.000
Paid employment	0.003	0.908	0.204	0.000
Current schooling	0.032	0.190	0.042	0.111
Marital relationship	0.100	0.000	0.102	0.000
Self-reported health (EQ-VAS)	0.022	0.326	-0.012	0.622
Health-related quality of life (HRQoL)	-0.073	0.002	-0.064	0.011
Model fit statistics				
CFI	0.959			
TLI	0.942			
RMSEA (90% Confidence Interval)	0.071 (0.067-0.074)			

Reference groups: gender=female; age=age 30-39; education=high-school or lower; household income=the highest 20% income quintile; employment=not in paid employment; schooling=not currently schooling; marital relationship=not married or do not have a partner.

Figure 1: Diagram of the latent variable SEM on Pandemic Anxiety Scale (Japanese n=2,764)



Note:

- 1) Variable labels in the SEM represent the following: male=male; genoth=other gender; hiedu=higher education; iq1=income quintile the lowest 20%; iq2=income quintile the second lowest 20-40%; iq3=income quintile the middle 40-60%; iq4=income quintile the higher 60-80%; eqvas=EQ-VAS; qol=HRQoL; pemp=paid employment; sch=schooling; married=married; g10=age 18-19; g20=age 20-29; g40=age 40-49; g50=age 50-59; F1=factor 1 on disease anxiety; F2=factor 2 on consequence anxiety; pas1-7=indicators per each PAS question.
- 2) Arrows (from left to right) indicate the relationship between the concerned variables in the tested SEM. An arrow between the two factors indicates factor correlations. All exogenous variables are covarying each other.

Journal Pre-proof

Title: Health inequity in pandemic anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach

Highlights:

- Health inequity in pandemic anxiety is assessed by SES (socioeconomic status).
- A latent variable SEM shows the multidimensionality of pandemic anxiety.
- We find unique evidence from Japan compared to European studies.
- Lower consequence anxiety under no paid work may be related to buffering measures.
- Unexpected associations between anxiety and general health to be further examined.

Journal Pre-proof

Ethical statement:

For the manuscript titled as “Social disparities in mental health related to anxiety about COVID-19 infection and socioeconomic consequences in Japan: A structural equation modeling approach” by Kyoko Shimamoto, Eoin McElroy and Yoko Ibuka.

The authors declare that this manuscript is the authors' own original work, which has not been previously published elsewhere. The paper is not currently being considered for publication elsewhere. The paper reflects the authors' own research and analysis in a truthful and complete manner.

Journal Pre-proof

Author statement:

KS - Conceptualization; Methodology; Data curation; Formal analysis; Project administration; Writing - original draft, review & editing.

EM – Conceptualization; Methodology; Writing- review & editing.

YI – Conceptualization; Writing- review & editing.

Journal Pre-proof