



Can Nutritional Epigenomics Explain Persistent Effects of Periconceptual Folic Acid in the Methylome?

Lees-Murdock, D., & Ward, M. (2022). Can Nutritional Epigenomics Explain Persistent Effects of Periconceptual Folic Acid in the Methylome? *The Journal of nutrition*, 152(12), 2636-2637.
<https://doi.org/10.1093/jn/nxac224>

[Link to publication record in Ulster University Research Portal](#)

Published in:
The Journal of nutrition

Publication Status:
Published (in print/issue): 31/12/2022

DOI:
[10.1093/jn/nxac224](https://doi.org/10.1093/jn/nxac224)

Document Version
Author Accepted version

General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

Can nutritional epigenomics explain persistent effects of periconceptual folic acid in the methylome?

¹ Diane Lees-Murdock and ² Mary Ward

Author Affiliations: ¹Genomic Medicine and ²Nutrition Innovation Centre for Food and Health (NICHE), School of Biomedical Sciences, Ulster University, Coleraine, Northern Ireland, United Kingdom.

Corresponding Author: Professor Mary Ward, Nutrition Innovation Centre for Food and Health (NICHE), School of Biomedical Sciences, Ulster University, Coleraine, Northern Ireland, United Kingdom, BT52 1SA. Tel: +442870123076, E-mail mw.ward@ulster.ac.uk

Sources of Support: Supported by funding from Northern Ireland Chest Heart & Stroke Association (NICH2021_09: DLM & MW); Health Education Authority, North-South scheme (EpiHyper Project 429436: DLM & MW); BBSRC, JPI ERA-HDHL scheme (EpiBrain project, Grant Ref: BB/S020330/1: MW). The funders had no role in the subject or preparation of this manuscript.

Conflict of interest: No conflicts for any author

Nutritional epigenomics is a burgeoning field of interest given the potential of dietary factors to influence the epigenetic landscape, with potential for long-term health impacts (1,2). DNA methylation, essential for cell division and tissue specification, is a dynamic epigenetic mechanism that can be modified by certain nutritional exposures including one-carbon nutrients, (3,4) which act by altering the availability of substrates required for methylation reactions or by inhibiting the enzymes that catalyse these reactions. Folate and the metabolically related B-vitamins supply methyl groups for DNA methylation via formation of S-adenosylmethionine (SAM) within one-carbon

metabolism. Pregnancy and adolescence, periods of intense development and growth with increased demands for folate, coincide with dynamic changes in DNA methylation and thus represents a window of opportunity for gene-nutrient interactions to impact health.

The evidence for protection of the foetus against neural tube defects by maternal periconceptual folic acid (FA) supplementation from randomized controlled trials (RCTs) is conclusive, with women worldwide recommended to take supplements of 400mg FA/day in the periconception period (5). Beyond the first trimester, analysis of cord blood samples from the Folic Acid Supplementation in the Second and Third Trimesters (FASSTT) trial in pregnancy, indicated that continued supplementation affected DNA methylation of specific genes in the offspring, including those related to brain development (6,7,8). These findings offer a potential biological mechanism linking maternal folate status with offspring neurodevelopment (6,7,8) and lend support to the fact that the DNA methylome undergoes widespread change during in-utero development, however evidence of a dynamic methylation landscape beyond birth through to adolescence has not been widely investigated. Of note, one recent landmark investigation, involving a series of epigenome-wide association studies in 5019 blood samples, collected from 2348 participants of two large independent cohorts at multiple time-points from birth to late adolescence, reported that changes in methylation over time were spread across the epigenome at over half of the methylation (CpG) sites analysed (9).

Given the lack of available evidence, the study by Crider and colleagues published in this issue of the *Journal of Nutrition* (10), exploring the long-term effect of maternal FA supplementation on DNA methylation in adolescent offspring is timely. The study contributes novel data, in a population not exposed to FA fortification, on the impact of maternal exposure to FA on the DNA methylome of offspring during adolescence. The authors were unable to detect any significant differences in DNA methylation between offspring of mothers who had taken FA in the periconceptual period and those who had not. A secondary analysis was conducted to examine

potential differences in allelic frequencies at SNPs implicated in folate metabolism. The authors also show that there are no significant differences in genotype frequencies of genes in the folate metabolising pathway between mothers and adolescents. Standard linkage disequilibrium was not possible due to lack of paternal genotype information; therefore, the authors note that caution should be taken in interpretation of these results. This study of Chinese mother-adolescent pairs highlights the complexities of studying the effect of nutrients on the dynamics of DNA methylation during periods of plasticity, given that the changes in methylation reported were not significant. The timing of FA exposure is also an important factor and varied to some extent in this study, with the majority, but not all mothers commencing FA prior to becoming pregnant. In addition, even though offspring from mothers who were supplemented with FA may have had altered DNA methylation profiles at birth, the low habitual dietary folate intake of the children and adolescents (both urban and rural) investigated may have prevented retention of methylation patterns induced by high levels of FA during pregnancy. Furthermore, maternal exposures to environmental factors, e.g. smoking (11) and BMI (12) are known to modulate DNA methylation with persistent effects in the offspring.

A number of limitations of the work were highlighted by the authors. The Infinium Human Methylation 450k array was used to compare patterns of methylation from adolescents of mothers who took FA supplements versus those who did not. The 450k microarray is enriched for CpG-dense genomic regions (CpG islands) however there is growing evidence that methylation in CpG-poor regions, such as enhancers and gene bodies, also have significant functional consequences through altering gene expression (13). Thus, to advance the study of environmental influences on the methylome, future studies should consider methods which offer greater genomic coverage such as whole genome bisulfite sequencing (WGBS) or the Infinium EPIC array which assesses over 800k sites simultaneously, thus achieving a good balance between genomic coverage and cost. The authors also discussed the use of saliva samples vs the more commonly used blood for collection of DNA as a limitation however, recent evidence suggests that saliva may, in some cases, be a better proxy for certain tissues, including brain (14). Saliva also represents a much less invasive means of accessing

DNA thus may offer a superior, alternative for future large-scale studies in targeted populations, including children and adolescents. The observational nature of this study is also a limiting factor particularly pertinent to the field of nutrigenomics, with longitudinal studies enabling the differentiation of intra-individual differences from inter-individual differences (15). Furthermore, the relatively small number of adolescents included in this study may have impacted the ability to detect changes in methylation. Future studies in large longitudinal cohorts are needed to definitively determine if methylation changes can persist through the life course, including during adolescence and early adulthood.

The current study provided a unique opportunity to investigate the impact of FA exposure *in utero* on long term epigenome wide DNA methylation patterns, in a population not exposed to FA fortification. While the study found no differences in those exposed vs not exposed to FA supplementation, the research highlights the need for further well-designed studies, including RCTs, to determine whether epigenetic changes can persist beyond early life. Explicit data regarding the timing of exposure and prevailing diet in populations being investigated are necessary to understand the pathological methylation changes associated with B-vitamin supplementation in promoting healthy development and growth. Such studies will enable persistent changes in the epigenome of offspring, throughout the life course following maternal exposures, to be tracked to help unravel disease mechanisms.

ORIGINAL UNL

REFERENCES

1. Amenyah SD, Ward M, Strain JJ, McNulty H, Hughes CF, Dollin C, Walsh CP, Lees-Murdock DJ. Nutritional epigenomics and age-related disease. *Curr Dev Nutr*. 2020; 4(7): nzaa097.
2. Stover PJ, James WPT, Krook A, Garza C. Emerging Concepts on the Role of Epigenetics in the Relationships between Nutrition and Health. *J. Intern. Med*. 2018;284:37–49.
3. Amenyah SD, McMahon A, Ward M, Deane J, McNulty H, Hughes CF, Strain JJ, Horigan G, Purvis J, Walsh CP, Lees-Murdock DJ. Riboflavin supplementation alters global and gene-specific DNA methylation in adults with the MTHFR 677 TT genotype. *Biochimie*. 2020;173:17-26.
4. Amenyah SD, Ward M, McMahon A, Deane J, McNulty H, Hughes C, Strain JJ, Horigan G, Purvis J, Walsh CP, Lees-Murdock DJ. DNA methylation of hypertension-related genes and effect of riboflavin supplementation in adults stratified by genotype for the MTHFR C677T polymorphism. *Int J of Cardiol*. 2021;322: 233–239.
5. Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Eng J Med*. 1992;327:1832–5.
6. Caffrey A, Irwin RE, McNulty H, Strain JJ, Lees-Murdock DJ, McNulty BA, Ward M, Walsh CP, Pentieva K. Gene-specific DNA methylation in newborns in response to folic acid supplementation during the second and third trimesters of pregnancy: epigenetic analysis from a randomized controlled trial. *Am J Clin Nutr*. 2018;107:566-75.
7. Irwin RE, Thursby SJ, Ondicova M, Pentieva K, McNulty H, Richmond RC, Caffrey A, Lees-Murdock DJ, McLaughlin M, et al. A randomized controlled trial of folic acid intervention in pregnancy highlights a putative methylation-regulated control element at ZFP57. *Clin Epigenet*. 2019;11:31.
8. Ondičová M, Irwin RE, Thursby SJ, Hillman L, Caffrey C, Cassidy T, McLaughlin M, Lees-Murdock DJ, Ward M, Murphy M, et al. Folic acid intervention during pregnancy alters DNA methylation, affecting neural target genes through two distinct mechanisms. *Clin Epigenet*. 2022;14:63.

9. Mulder RH, Neumann A, Cecil CAM, Walton E, Houtepen LC, Simpkin AJ, Rijlaarsdam J, Heijmans BT, Gaunt TR, Felix JF, Jaddoe VVW, Bakermans-Kranenburg MJ, Tiemeier H, Relton CL, van IJzendoorn MH, Suderman M. Epigenome-wide change and variation in DNA methylation in childhood: trajectories from birth to late adolescence. *Hum Mol Genet.* 2021;30(1):119-134.
10. Crider et al., JN-2022-0525R
11. Rebecca C. Richmond, Andrew J. Simpkin, Geoff Woodward, Tom R. Gaunt, Oliver Lyttleton, Wendy L. McArdle, Susan M. Ring, Andrew D.A.C. Smith, Nicholas J. Timpson, Kate Tilling, et al., Prenatal exposure to maternal smoking and offspring DNA methylation across the lifecourse: findings from the Avon Longitudinal Study of Parents and Children (ALSPAC), *Hum Mol Genet.* 2015;24(8):2201–2217
12. Sharp GC, Salas LA, Monnereau C, Allard C, Yousefi P, Everson TM, Bohlin J, Xu Z, Huang RC, Reese SE et al., Maternal BMI at the start of pregnancy and offspring epigenome-wide DNA methylation: findings from the pregnancy and childhood epigenetics (PACE) consortium. *Hum Mol Genet.* 2017 Oct 15;26(20):4067-4085.
13. Gutierrez-Arcelus M, Ongen H, Lappalainen T, Montgomery SB, Buil A, Yurovsky A, Bryois J, Padioleau I, Romano L, Planchon A et al., Tissue-Specific Effects of Genetic and Epigenetic Variation on Gene Regulation and Splicing. *PLOS Genetics* 2015 11(1): e1004958.
14. Thomas M, Knoblich N, Wallisch A, Glowacz K, J Becker-Sadzio, Gundel F, Brückmann , Nieratschker V, Increased BDNF methylation in saliva, but not blood, of patients with borderline personality disorder. *Clin Epigenet* 2018;10:109.
15. Fitzmaurice et al., 2012 Applied Longitudinal Analysis, John Wiley & Sons, Hoboken, New Jersey.