



The impact of urinary incontinence on multiple health outcomes:

Soysal, P., Veronese, N., Ippoliti, S., Pizzol, D., Carrie, A. M., Stefanescu, S., López-Sánchez, G. F., Barnett, Y., Butler, L., Koyanagi, A., Jacob, L., Ghaydya, R. A., Sheyn, D., Hijaz, A. K., Oliva-Lozano, J. M., Muyor, J. M., Trott, M., Kronbichler, A., Grabovac, I., ... Smith, L. (2023). The impact of urinary incontinence on multiple health outcomes: an umbrella review of meta-analysis of observational studies. *Aging Clinical and Experimental Research*, 35(3), 479-495. Advance online publication. <https://doi.org/10.1007/s40520-022-02336-0>

[Link to publication record in Ulster University Research Portal](#)

Published in:

Aging Clinical and Experimental Research

Publication Status:

Published online: 13/01/2023

DOI:

[10.1007/s40520-022-02336-0](https://doi.org/10.1007/s40520-022-02336-0)

Document Version

Author Accepted version

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

1 **Review**

2 **The impact of urinary incontinence on multiple health outcomes: an umbrella review of**
 3 **meta-analysis of observational studies**

4 *Running Title: Urinary Incontinence and Health Outcome*

5 Pinar Soysal^{1*}, Nicola Veronese^{2*}, Simona Ippoliti^{3*}, Damiano Pizzol^{4*}, Anne Marie
 6 Carrie³, Simina Stefanescu³, Guillermo F. López-Sánchez⁵, Yvonne Barnett⁶, Laurie Butler⁶,
 7 Ai Koyanagi⁷, Louis Jacob⁸, Ramy Abou Ghaydya^{9,10}, David Sheyn^{9,10}, Adonis K Hijaz^{9,10},
 8 Jose M Oliva-Lozano¹¹, Jose M Muyor¹¹, Mike Trott¹², Andreas Kronbichler¹³, Igor
 9 Grabovac¹⁴, Mark Tully¹⁵, Lin Yang¹⁶, Jimin Hwang¹⁷, Jong Yeob Kim¹⁸, Seoyeon Park¹⁸,
 10 Junmin Song¹⁹, Jae Il Shin²⁰, Petre-Cristian Ilie^{3±}, Lee Smith^{11±}

11 *On behalf of the European Society of Geriatric Medicine Special Interest Group in*

12 *Systematic Reviews and Meta-Analyses*

13 *Authors contributed equally

14 ±Senior authors

- 15 1. Department of Geriatric Medicine, Faculty of Medicine, Bezmialem Vakif
 16 University, Adnan Menderes Bulvarı (Vatan Street), 34093 Fatih, İstanbul, Turkey.
- 17 2. Department of Internal Medicine, University of Palermo, Geriatrics Section, Palermo,
 18 Italy.
- 19 3. The Queen Elizabeth Hospital Foundation Trust, King's Lynn, United Kingdom.
- 20 4. Italian Agency for Development Cooperation, 33 Street, Amarat, Khartoum 79371,
 21 Sudan
- 22 5. Vision and Eye Research Institute, School of Medicine, Faculty of Health, Education,
 23 Medicine and Social Care, Anglia Ruskin University, Cambridge, United Kingdom.
- 24 6. Faculty of Science and Engineering, Anglia Ruskin University, Cambridge, UK, CB1
 25 1PT
- 26 7. Research and Development Unit, Parc Sanitari Sant Joan de Déu, CIBERSAM,
 27 08830, Barcelona, Spain; ICREA, Pg. Lluís Companys 23, 08010, Barcelona, Spain.

- 28 8. Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines, 78180
29 Montigny-le-Bretonneux, France.
- 30 9. Department of Urology, University Hospitals Cleveland Medical Center, 11000
31 Euclid Avenue, Cleveland, OH, USA
- 32 10. Case Western Reserve University School of Medicine, Cleveland, OH, USA
- 33 11. Health Research Centre, University of Almeria, Almeria, Spain.
- 34 12. The Cambridge, Centre for Sport and Exercise Sciences, Anglia Ruskin University,
35 Cambridge, UK, CB1 1PT
- 36 13. Department of Internal Medicine IV, Medical University Innsbruck, Innsbruck,
37 Austria.
- 38 14. Department of Social and Preventive Medicine, Centre for Public Health, Medical
39 University of Vienna, 1090 Wien, Austria
- 40 15. Institute of Mental Health Sciences, School of Health Sciences, Ulster University,
41 Newtown Abbey BT37 0QB, UK
- 42 16. Department of Cancer Epidemiology and Prevention Research, Cancer Control
43 Alberta, Alberta Health Services, Calgary, Canada
- 44 17. Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health,
45 Baltimore, MD, USA
- 46 18. Yonsei University College of Medicine, Seoul, Republic of Korea
- 47 19. Keimyung University School of Medicine, Daegu, Republic of Korea
- 48 20. Department of Pediatrics, Yonsei University College of Medicine, Seoul, Republic of
49 Korea

50

51 Corresponding author:

52 Professor Jae Il Shin, Department of Pediatrics, Yonsei University College of Medicine, Seoul
53 03722, Korea. (telephone: 82-2-2228-2050; FAX: 82-2-393-9118; email: SHINJI@yuhs.ac)

54

55 **Word count (abstract)** – 247 words56 **Word count (main text)** – 3,841 words

57 **Funding:** No financial support was provided for research conduct and/or preparation of the
58 article.

59 **Abstract**

60 **Background & Aim:** We aimed to capture the breadth of health outcomes that have been
61 associated with the presence of Urinary Incontinence (UI) and systematically assess the quality,
62 strength, and credibility of these associations through an umbrella review and integrated meta-
63 analyses.

64 **Methods:** We assessed meta-analyses of observational studies based on random-effect
65 summary effect sizes and their p-values, 95% prediction intervals, heterogeneity, small-study
66 effects, and excess significance. We graded the evidence from convincing (Class I) to weak
67 (Class IV).

68 **Results & Discussion:** From 3172 articles returned in search of the literature, 9 systematic
69 reviews were included with a total of 41 outcomes. Overall, 37 out of the 41 outcomes reported
70 nominally significant summary results ($p < 0.05$), with 22 associations surviving the application
71 of a more stringent p-value ($p < 10^{-6}$). UI was associated with worse scores than controls in
72 female sexual function (Class II), while it was also associated with a higher prevalence of
73 depression (odds ratio [OR]=1.815; 95% confidence interval [CI]: 1.551-2.124), and anxiety
74 (OR=1.498; 95%CI: 1.273-1.762) (Class IV). UI was associated with poorer quality of life
75 (QoL), higher rate of mortality (hazard ratio=2.392; 95%CI: 2.053-2.787) an increase in falls,
76 frailty, pressure ulcers, diabetes, arthritis, and fecal incontinence (Class IV).

77 **Conclusions:** UI is associated with female sexual dysfunction, with highly suggestive
78 evidence. However, the evidence of other adverse outcomes including depression, anxiety,
79 poorer QoL, higher mortality, falls, pressure ulcers, diabetes, arthritis, fecal incontinence, and
80 frailty is only weak. A multidimensional approach should be taken in managing UI in the
81 clinical setting.

82

83 **Keywords:** Urinary Incontinence, Health Outcomes, Umbrella Review, Meta-analysis

84 **Statements and Declarations:** None

85 **Competing Interests:** All authors confirm to have no actual or potential conflict of interests.

86

87 INTRODUCTION

88 The International Association of Urinary Incontinence (ICS) defines urinary incontinence (UI)
89 as complaints of involuntary leakage of urine, which is recognized by the World Health
90 Organization as an important health problem with significant social and economic burden[1,
91 2]. UI can be observed and may cause negative clinical consequences at any age[3, 4].
92 Regardless of the UI type, the prevalence among females is 17.6% between the ages of 20-39
93 years, 27.9% among females aged 20-49 years, while it is present in almost one out of every
94 two females aged over 60 years[3]. Among males, UI is extremely rare before the age of 65
95 years; however, its prevalence is 15% among those aged 65 years and over and 25% among
96 those aged 80 years and over[5]. Therefore, UI affects millions of people around the world,
97 and it is important to determine what complications it can cause.

98 For many years, there have been an increasing number of studies investigating the mental and
99 physical effects of UI on people of both sexes, different age groups, and in different physiologic
100 states, such as pregnant women or institutionalized geriatric patients[6, 7]. According to their
101 results, patients with UI have a lower quality of life (QoL), lower labor productivity, poorer
102 sexual function, and higher risk of major depressive disorder and social isolation[8]. Especially
103 in older adults, the risks of major adverse events are high, including falls and fractures, skin
104 infections, functional impairment, caregiver burden, institutionalization, and mortality[9-15].
105 Considering both the indirect costs associated with these complications and the health
106 expenditure required for the diagnosis and treatment of UI, the annual cost-of-illness for UI in
107 European populations was calculated as €7 billion; in the United States, UI was estimated to
108 cost \$82.6 billion in 2020[16, 17]. These studies, published with increasing frequency, resulted
109 in the publication of systematic reviews with meta-analyses. However, to date, most systematic
110 reviews have focused on a single disease endpoint, and no systematic evaluation of the
111 relationships between UI and various physical and mental health outcomes has been carried

112 out. A better understanding of the full spectrum of health risks associated with UI is important
113 for clinical practice.

114 Therefore, this study was designed to capture the breadth of results shown in observational
115 studies regarding UI and to systematically evaluate the quality, strength, and reliability of these
116 relationships. We used an umbrella review with integrated meta-analyses to combine evidence
117 from a wide range of outcomes and populations.

118

119 **METHODS**

120 This umbrella review followed a pre-planned but unpublished protocol available upon request
121 to the corresponding author. This study follows the Preferred Reporting Items for Systematic
122 Reviews and Meta-Analyses (PRISMA) indications for reporting findings of systematic
123 reviews and meta-analyses[18].

124

125 *Data sources and searches*

126 We conducted an umbrella review[19], searching the MEDLINE, Scopus, Embase databases,
127 Cochrane library, CINAHL and PSYCINFO from inception until 20th July 2022 with: “(Meta-
128 Analysis[ptyp] OR metaanaly*[tiab] OR meta-analy*[tiab] OR Systematic review [ptyp] OR
129 “systematic review” [tiab]) AND ("urin* incontinence"[tiab] OR "bladder incontinence"[tiab]
130 OR "urin* leakage"[tiab] OR "urine bladder incontinence"[tiab] OR "urine leakage"[tiab] OR
131 wetting [tiab] OR incontinence)[tiab]. In addition, we hand-searched the reference lists of
132 eligible5 articles.

133

134 *Study selection*

135 In this umbrella review, we included: (1) systematic reviews with meta-analyses that included
136 people with UI; (2) meta-analyses of observational studies (longitudinal or case-control) that
137 investigated the association of UI with any health-related outcome (e.g., QoL, falls, depression,
138 mortality). Two authors (PS, LS) independently performed title and abstract screening.
139 Disagreements were resolved through consensus with another independent author (DP). Full
140 texts of all potentially eligible articles were then retrieved by the same three authors and any
141 disagreement was resolved with another independent author (NV).

142

143 *Data extraction*

144 Two independent investigators (SI, MT) extracted the following information for each meta-
145 analysis independently: first author name; publication year; the number of studies; study
146 population; type of effect size; study design; the number of participants with (cases) and
147 without (controls) events for each study. We also extracted the study-specific estimated relative
148 risk for health outcomes (risk ratio, RR; odds ratio OR; hazard ratio, HR; mean difference,
149 WMD; standardized mean difference, SMD) and 95% confidence intervals (CIs). We finally
150 extracted the data for the Assessment of Multiple Systematic Reviews (AMSTAR)-2 tool[20].
151 When more than one meta-analysis on the same research question using the same study design
152 was identified, the one with the largest number of participants was selected.

153

154 *Data synthesis and analysis*

155 For each meta-analysis, we estimated the summary effect size and its 95% CI by using a
156 random-effects model based on the DerSimonian-Leird method[21]. We also estimated the
157 prediction intervals (PIs) and its 95% CI, which further accounts for between-study effects and
158 estimates the certainty of the association if a new study may address the same association[22-

159 24]. Between-study inconsistency was estimated with the I^2 metric, with values $\geq 50\%$
160 indicative of heterogeneity and values between 50-75% of moderate heterogeneity and $\geq 75\%$
161 of high heterogeneity[25]. Then, we calculated the evidence of small-study effects (i.e. whether
162 small studies inflated effect sizes) using the regression asymmetry test with a p-value < 0.10
163 being indicative of this potential bias[26].

164 Finally, we applied the excess of significance test (Ioannidis' test)[27]. Because of the limited
165 statistical power of this test, a lenient significance threshold ($p < 0.10$) was adopted[28]. We
166 considered the effect size of the largest study for each outcome, and based on this, we estimated
167 the power of each constituent study with an algorithm using a non-central t distribution. Excess
168 significance for each meta-analysis was considered whenever $p < 0.10$. All statistical analyses
169 were conducted using Stata, version 14.0 (StataCorp).

170

171 *Grading the evidence*

172 For observational studies, using the criteria mentioned above, significant associations (i.e.
173 $p < 0.05$) were categorized into convincing, highly suggestive, suggestive, or weak evidence
174 (class I to IV), following a grading scheme that has already been applied in various fields, as
175 reported in Supplementary Table S2 [29-32]. According to credibility assessment criteria for
176 meta-analyses of observational studies, the level of evidence was determined as follows:
177 **Convincing (class I):** High significant association; large sample size, having the event of
178 interest; the largest component study reporting a nominal statistically significant result; a 95%
179 PI that excluded the null; no large heterogeneity; no evidence of small-study effect; no excess
180 significance bias. **Highly suggestive (class II):** High significant association, large sample size
181 having the event of interest; the largest component study reporting a statistically significant

182 result. **Suggestive (class III):** Significant Associations and large sample size having the event of
183 interest. **Weak (class IV):** Remaining statistically significant associations with $P < .05$.

184 We assessed the methodological quality of the included meta-analyses of observational studies
185 using AMSTAR-2 that ranks the quality of a meta-analysis from critically low to high
186 according to 16 predefined items [20]. The items are about whether it contains research
187 questions and inclusion criteria for the review; the protocol; explaining selection of the study
188 design; using a comprehensive literature search strategy; performing study selection data
189 extraction in duplicate; providing a list of excluded studies and justify the exclusions;
190 describing the included studies in adequate detail; assessing the risk of bias; reporting on the
191 sources of funding; use appropriate methods for statistical combination of results; assessing the
192 potential impact of risk of bias on the results; accounting for bias in primary studies when
193 interpreting/discussing the results of the review; investigation of publication bias and
194 heterogeneity, and conflict of interest. The critical domains and non-critical domains were
195 evaluated by using AMSTAR-2, which is shown in Table 2 in detail [20].

196

197 **RESULTS**

198 *Literature review*

199 Overall, we identified 3172 papers. After removing the duplicates (985), 2187 title/abstracts
200 were screened with 57 eligible full-texts. Of them, 9 studies were finally included in our
201 umbrella review (Figure 1).

202

203 *Findings of the case-control and cross-sectional studies*

204 The median number of studies of meta-analyses for each outcome was 5 (range 2-66), the
205 median number of participants was 814 (range 388 to 321,939), and the median number of
206 cases was 363. A total of 41 independent outcomes was finally included.

207 Overall, 37 out of the 41 outcomes reported nominally significant summary results ($p < 0.05$)
208 ($=90.2\%$), with 22 associations surviving the application of a more stringent p-value ($P < 10^{-6}$),
209 as shown in Table 1.

210 Heterogeneity among studies was high, with only 14 out of 41 outcomes having low
211 heterogeneity ($I^2 < 50\%$). On the contrary, 10 out of 41 had moderate heterogeneity (I^2 between
212 50 and 75%) and the other 17 had high heterogeneity. Eight associations presented 95% PIs
213 excluding the null value. The small-study effect was present in 10 out of 41 outcomes included,
214 three outcomes reported an excess significance bias, and 36 out of 41 outcomes had their largest
215 study reporting statistically significant results.

216 As shown in Table 1, using the criteria mentioned above, UI was associated with significantly
217 worse scores than controls in Female Sexual Function in five studies including 436 females
218 before and after overactive bladder treatment, as shown in Figure 2 (lubrication, orgasm,
219 satisfaction scores supported by a class II evidence; pain and total score sustained by a class
220 III)[33]. The evidence supporting the role of UI in Mid-Urethral Sling (MUS) surgery for SUI
221 at 6 months post-operative in female sexual function was supported by a weak strength of
222 evidence, similar to the evidence in MUS surgery for SUI at 12 months post-operative[34].

223 Regarding mental health, UI was associated with a higher prevalence of depression
224 ($OR=1.815$; 95%CI: 1.551-2.124), and anxiety ($OR=1.498$; 95%CI: 1.273-1.762) than people
225 without UI (Figure 3)[35, 36]. However, given the susceptibility of this evidence to excess
226 significance bias, these outcomes should be interpreted as having a weak strength of evidence.

227 Moreover, UI was associated with poorer QoL[36], higher presence of Grade II+ pressure ulcer
228 development[37], higher rate of mortality (n=66 studies, HR=2.392; 95%CI: 2.053-2.787, class
229 IV)[38] and an increased presence of falls and frailty than their counterparts, although these
230 were graded as class IV evidence (Figure 4)[39-41]. Finally, urgency UI was associated with
231 an increased presence of falls and fecal incontinence (FI), arthritis, diabetes supported by class
232 IV evidence[39], as shown in Table 1.

233 As reported in Supplementary Table S3, only one meta-analysis was rated “moderate”, three
234 rated “low”, whilst the other meta-analyses were rated as “critically low” according to the
235 AMSTAR-2 criteria.

236

237 **DISCUSSION**

238 This umbrella review summarized the findings of ten previous meta-analyses of the association
239 between UI and 41 independent outcomes. Highly suggestive (i.e., class II) evidence was found
240 for associations between UI and female sexual dysfunction. The other outcomes including
241 higher prevalence of depression and anxiety, poorer QoL, higher mortality, falls, fecal
242 incontinence, pressure ulcers, and frailty were found to have weak evidence.

243

244 *Female Sexual Dysfunction (FSD)*

245 In this umbrella review, it was determined that the only result, reaching the class II level of
246 evidence among the UI-related health outcomes, was female sexual dysfunction. UI worsened
247 the parameters of desire, arousal, lubrication, orgasm, and satisfaction scores of FSD.
248 Overactive bladder (OAB) treatment improved both OAB-wet and sexual functions
249 simultaneously[33]. It has been shown that placement of a MUS, performed for stress

250 incontinence in females, positively affects sexual function with class IV evidence level due to
251 improvement in UI at 6th and 12th months follow-up[34].

252 UI and FSD are two common conditions that are typically underdiagnosed and undertreated.
253 Although it seems clear that these two urogenital conditions may be interrelated, they are
254 generally considered as two separate problems[42]. For example, one study showed that nearly
255 three-quarters of women who attended a urology clinic for UI or other lower urinary tract
256 symptoms were not asked about their sexual health problems[43]. Another important issue is
257 that although UI and FSD are commonly thought to be conditions related to aging, they can be
258 seen widely throughout life, including the premenopausal period[42]. The prevalence of FSD
259 in females with UI was at least 25% and of women with UI who were sexually active, 23% to
260 56% had FSD⁴³. Even without UI, FSD has negative effects on self-esteem, well-being, and
261 the establishment of strong relationships with a partner[42]. Negative effects of UI on sexual
262 function resulting in coital incontinence may increase the intensity of these harmful effects and
263 cause a decrease in the frequency of sexual activity in women due to smell, embarrassment,
264 loss of self-confidence, and fear of repetition of UI in subsequent sexual intercourse[44, 45].
265 Considering that sexual activity is associated with a range of benefits for psychological and
266 physiological well-being, such as improved QoL and mental health, heart rate variability, and
267 lower risk of certain cancers and fatal coronary events, the relationship between UI and sexual
268 dysfunction should be further investigated[46-48].

269 All three UI types (stress, urgency, and mixed) worsen sexual functions[49-51]. However, the
270 reports on the response of sexual function following the treatment of UI are conflicting[33, 34,
271 42]. In this umbrella review, it was found that MUS surgery applied for stress UI improves
272 postoperative desire, arousal, orgasm, lubrication, satisfaction, and pain scores and
273 significantly reduces coital incontinence[34]. On the other hand, urge UI (UUI, OAB-wet) was
274 reported to be the most influencing factor on sexuality. OAB treatments showed improvement

275 in both OAB-wet and sexual function, and OAB therapies significantly improved OAB and
276 reduced female sexual dysfunction[33]. Therefore, it is important to realize that UI is a treatable
277 risk factor for FSD, which is known to be multifactorial.

278

279 ***Mental health***

280 UI not only causes physical problems but can also cause mental health problems such as
281 depression and anxiety[52]. Although it is not a life-threatening illness, those with UI may feel
282 high levels of stress, embarrassment, and discomfort from the odor. Moreover, UI can
283 sometimes occur so quickly and in large volumes that it can seriously affect one's
284 socialization[53]. UI causes a decrease in performance in daily life activities owing to frequent
285 visits to the toilet, while general health problems such as UI-related sleep loss and daytime
286 fatigue may increase psychological and emotional distress[54, 55]. Just like FSD, urinary
287 system problems can deter males from leaving the house and cause limitations in their social
288 relationships[53]. For males, urinary problems are stigmatizing and can reduce their masculine
289 identity, causing internalization of negative self-worth and low self-esteem[56]. UI can lead to
290 a lower sense of self-control, leading to the belief that those with UI have a lower coping
291 capacity and reduce their sense of self-efficacy[57]. Due to all these reasons, depression and
292 anxiety can be observed frequently in those with UI.

293 In addition, UI is common in those with multimorbidity and frailty, and within these patients
294 UI is highly likely to negatively affect their mental health, above that of multimorbidity,
295 potentially resulting in the development of clinical depression or anxiety[4, 41, 58, 59].
296 Therefore, common factors in etiology may explain the relationship between UI and mental
297 health. It is worth noting findings from a previous cross-sectional study that UI-related
298 depression and anxiety did not benefit from anti-depressants[60], while another study found

299 that treatments applied to those with UI significantly improved UI, anxiety, and depression
300 with significant correlations in their symptoms[61].

301

302 *Quality of life (QoL)*

303 Often, individuals with UI deny and hide UI, causing physical and psychosocial restrictions in
304 daily life[36]. In fact, key consequences include loss of self-esteem and social isolation, in
305 addition to other negative outcomes such as anxiety, depression, sexual impairment, and
306 decreased physical activity[36]. All these conditions are associated with poor QoL.

307 UI can be associated with poor QoL through a variety of mechanisms. First, people with UI
308 usually exhibit more co-morbidities than those without. Although several risk factors of UI
309 have been reported, the most specifically related are gender, age, dementia, and mobility[62].
310 Moreover, fluid intake, self-motility, and diuretic therapy can also affect diuresis and thus
311 UI[62]. All these factors are commonly known to be associated with poor QoL in older people.
312 Second, it is possible for people with UI to use diapers, and the use of these tools, in certain
313 circumstances, can lead to Incontinence-Associated Dermatitis (IAD)[63], defined as "skin
314 rash and edema, sometimes accompanied by blisters with serous exudate, erosion, or secondary
315 infection"[64]. IAD, like other dermatological conditions, is associated with a poor QoL[36].
316 Finally, we believe that the poor QoL in UI can be justified by the presence of shame leading
317 to a change in lifestyles and habits (i.e., reduced or suppressed physical activity) and mental
318 disorders in these people (i.e., depression and anxiety)[2, 60].

319

320 *Mortality*

321 The relationship between UI and premature death is probably multifactorial. First, the risk
322 factors that cause UI development may themselves negatively affect survival. Conditions
323 known as risk factors for UI such as age, multimorbidity, cognitive impairment, frailty, and
324 disability are likely to increase mortality. For example, frail and elderly patients are at the
325 highest risk of developing UI[41]. Therefore, the mortality rate of patients suffering from UI
326 is expected to be higher than those who do not suffer from this condition. A recent study
327 revealing that the relationship between UI and death may be comprehended based on increased
328 frailty in incontinent individuals supports this hypothesis[65]. Second, UI may shorten survival
329 by increasing multiple unfavorable outcomes. For example, UI increases the risk of falls and
330 fall-related injuries in both genders[11, 66, 67], and it is well known that hip fracture may
331 increase the risk of low mobility by more than 4 times, the risk of rehospitalization by 2.5 times
332 and mortality by 1.8 times, especially in the elderly[68]. On the other hand, complications such
333 as depression, anxiety, sexual dysfunction, and decreased QoL caused by UI may indirectly
334 lead to a shortening of survival. In addition, a recent retrospective study reported that drugs
335 used for UI increase the risk of mortality by 50% due to their anticholinergic effects[69].
336 However, future studies may reveal the truth of this hypothesis.

337

338 ***Grade II+ pressure ulcer development***

339 Pressure ulcers cause severe pain, physical and psychological discomfort, and limitations in
340 activities, and also lead to a prolonged hospital stay, healthcare utilization, and mortality[37].

341 The skin surface microclimate includes temperature and humidity. Exposure to moisture can
342 cause moisture-related skin damage in the sacral area as a consequence of inflammation of the
343 epidermis and dermis. Moisture-related damage, often incorrectly classified as a type of
344 pressure ulcer, includes sweating-related intertrigo, skin damage around the wound resulting

345 from wound exudate, or effluent. The most common cause of moisture-related skin damage is
346 IAD[64]. Skin irritants from incontinence include urine, feces, double incontinence, and liquid
347 feces[37, 64]. Moisture from incontinence increases the vulnerability of the skin and superficial
348 tissue layers to pressure-induced blood flow reduction[70]. Moisture also attenuates the skin
349 and makes it more exposed to the effects of pressure and shear[64, 71]. Exposure to urine and
350 feces results in skin hyperhydration and an increase in skin pH, reducing tissue tolerance and
351 increasing the risk of local infection by enabling microorganisms to multiply[71].

352

353 *Falls*

354 The link between urinary incontinence and falls is likely to be related to the need to rush to the
355 toilet and the distress and anxiety related to the aftermath of not being able to get to the toilet
356 in time[62]. This hypothesis is in line with a previous report that falls usually happen in the
357 bathroom[72]. Furthermore, older patients with UUI may be particularly vulnerable because
358 UI severity can increase over time, mobility decreases with age, and older individuals are more
359 susceptible to injury from falls[11]. Cognitive demands of carrying out multiple tasks at the
360 same time, such as rushing to the toilet quickly, focusing on controlling urine flow, and
361 overcoming obstacles at home, can also have a harmful effect on sustaining balance in older
362 people[73]. Therefore, the UUI rate was higher in those who reported falls (34.5%) than those
363 who did not (19.6%) with increased urine leakage volume related to a higher risk of falling[74].
364 Although it is thought that controlling the UI through medication may be beneficial in
365 preventing falls, which is an important health problem, these drugs also have the potential to
366 increase falls[75].

367

368 *Frailty*

369 It has been demonstrated that UI leads to a wide variety of adverse effects in older people,
370 including falls, urinary tract infections, skin complications, functional decline, psychosocial
371 limitations, poor quality of life, and poor health perception[41]. These complications of UI may
372 cause an accumulation of defects that are well known to create a state of frailty. Conversely,
373 individuals with frailty frequently experience homeostatic dysregulations leading to
374 impairments in physical functioning, mobility, gait and balance, and cognition, which might
375 result in UI. An important distinction in UI etiology between healthy and weak older people is
376 the presence of conditions and factors outside of the lower urinary tract - such as cognitive
377 impairment, poor mobility, and polypharmacy - that can precipitate the loss of continence and
378 exacerbate the main urinary symptoms[41, 76]. Furthermore, this finding justifies the need for
379 a multidimensional approach and clinical evaluation of the UI in older people to assess the key
380 risk factors and the etiology of the UI, with the inclusion of new components to determine
381 frailty, and then the integral and patient-specific treatment plan[41, 76].

382

383 *Diabetes, Fecal incontinence, and Arthritis*

384 Diabetes is an independent risk factor for UI. As a matter of fact, hyperglycemia is likely to be
385 associated with earlier and more common adverse outcomes compared to other microvascular
386 complications such as retinopathy, neuropathy, or nephropathy[77]. A plausible cause of UI is
387 microvascular harm to the innervations of the bladder and urethral sphincter, sphincter
388 dysfunction, bladder instability, urinary retention, and elevated postvoid residual urine volume
389 adding to overflow UI, chronic bacterial colonization, and urinary tract infections (UTIs)[78,
390 79]. Many different patient backgrounds and clinical characteristics have been recognized as
391 risk factors for UI. Aging is an important risk factor for UI as it is associated with decreased
392 sensation, detrusor muscle mass, elasticity, and bladder capacity[77]. Obesity also increases

393 the risk of UI among diabetic women because it causes increased intra-abdominal and pelvic
394 pressure[77]. The presence of frequent UTIs and the number of normal vaginal deliveries
395 (parity) are other important risk factors for UI[80]. Clinical variables of patients have also been
396 specified as important risk factors for UI in epidemiological studies and include diabetes
397 duration, hemoglobin A1c level, and the presence of diabetic long-term complications[81].
398 Some investigations demonstrate that up to 50% of severe incontinence could be avoided by
399 precluding type 2 diabetes[77]. This underscores that there is an important link between both
400 health conditions.

401 The lower bowel (LB) and the lower urinary tract (LUT) are closely related. Both organs
402 originate from the embryological cloaca and have a similar function: storage and evacuation of
403 feces and urine, respectively[82]. Peripheral innervation plays a similar role in the function of
404 both viscera. The central processing and perception of afferent activity ensue in the same brain
405 areas. The close relationship between the LB and the LUT also has clinical relevance as the
406 pathology of both co-exist[82]. UI and FI are two highly prevalent pelvic floor disorders. About
407 20% of women with UUI symptoms may also have symptoms of FI, referred to as double
408 incontinence[83]. Women who suffer from both diseases have greater impairment regarding
409 their physical and psychosocial wellbeing than do women suffering from isolated UI or FI,
410 resulting in social isolation and reduced quality of life[84]. The usefulness of the agents used
411 in the treatment of UUI in FI may be a clear indication of the close relationship between these
412 two conditions[85].

413 Arthritis may cause restricted mobility and the ability to disrobe quickly, leading to UI. Those
414 with UUI may be more likely to have painful osteoarthritis symptoms due to increased physical
415 demands from responding to the greater frequency of urgent urination episodes on their already
416 pre-existing osteoarthritis. Arthritis may also limit the ability to change positions in order to
417 prevent stress incontinence[86].

418 In this review, we found that the discrepancy between the results presented and the commonly
419 reported associations between urinary incontinence and health outcomes, except for FSD. The
420 possible reasons for this can be explained by the limitations. First, the eligible meta-analyses
421 included studies with significantly different designs, populations, and other basic
422 characteristics (e.g., women, men, older and middle-aged adults), which may have contributed
423 to the large heterogeneity in some meta-analyses. However, a common estimate of
424 heterogeneity ($I^2 < 50\%$) was used as one of the criteria for grading convincing outcomes, even
425 if the use of the same I^2 is still discussed [87]. Second, according to AMSTAR-2, most of the
426 reviews had low or critically low-level quality, because most of them did not report funding,
427 or did not have a pre-registering protocol. However, to estimate if these biases can modify our
428 outcomes is hard to hypothesize. Another factor was that they did not assess the potential
429 impact of risk of bias in individual studies on the results of the meta-analysis or the authors did
430 not account for bias in primary studies when interpreting/discussing the results of the reviews.
431 Large heterogeneity and risk of bias were responsible for both grading of the evidence and low
432 quality. Therefore, future studies should be done by eliminating them with a better research
433 strategy.

434 **Conclusions**

435 UI seems to be significantly associated with several negative health outcomes, although only
436 the association with FSD is supported by highly suggestive evidence. However, the present
437 review does not allow firm conclusions to be made on whether UI can be considered as a risk
438 factor for other medical conditions including depression, anxiety, poorer QoL, higher
439 mortality, falls, pressure ulcers, diabetes, arthritis, fecal incontinence, and frailty. A
440 multidimensional approach to clinical assessment and treatment of UI taking account of both
441 physical and mental health symptoms is warranted.

443 **Authors' Contributions:** All authors made substantial contributions to all of the following;
444 (1) conception and design of the study, data acquisition, or analysis and interpretation of data;
445 (2) drafting or critical revision of the article for intellectual content; and (3) final approval of
446 the version to be submitted.

447 **Ethical Approval Statement:** Not applicable since this manuscript uses a methodology of
448 umbrella review.

449 **Informed Consent Statement:** Not applicable since this manuscript is a secondary analysis.

450

451

452

453 **FIGURE LEGENDS**

454 **Figure 1.** PRISMA flow diagram of the literature search.

455 **Figure 2.** Summary estimates of meta-analyses regarding sexual function (desire, arousal,
456 lubrication, orgasm, satisfaction, pain), female coital incontinence, and treatment of UI.

457 **Figure 3.** Summary estimates of meta-analyses regarding mental health (depression, anxiety,
458 and SF-36 score) and UI.

459 **Figure 4.** Summary estimates of meta-analyses regarding quality of life and systemic disease
460 and UI

461

462 **TABLE LEGENDS**

463 **Table 1.** Main findings of the case-control and cross-sectional studies

464 **Table 2.** AMSTAR-2 quality assessment of systematic reviews and meta-analysis.

465

466 **SUPPLEMENTARY TABLE LEGENDS**

467 **Supplementary Table S1.** PRISMA Checklist

468 **Supplementary Table S2.** Credibility assessment criteria for meta-analyses of observational
469 studies.

470

471 **References:**

- 472 1. Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, et al. The standardisation
473 of terminology in lower urinary tract function: report from the standardisation sub-committee
474 of the International Continence Society. *Urology*. 2003;61(1):37-49. doi: 10.1016/s0090-
475 4295(02)02243-4.
- 476 2. de Macedo Dantas TH, Castaneda L, Correia GN, Campelo C, de Sousa Dantas D.
477 Functioning and disability of premenopausal women with urinary incontinence: An
478 assessment by using the World Health Organization Disability Assessment Schedule-
479 WHODAS 2.0. *Neurourol Urodyn*. 2019;38(6):1767-74. doi: 10.1002/nau.24073.
- 480 3. Abufaraj M, Xu T, Cao C, Siyam A, Isleem U, Massad A, et al. Prevalence and trends in
481 urinary incontinence among women in the United States, 2005-2018. *Am J Obstet Gynecol*.
482 2021;225(2):166.e1-.e12. doi: 10.1016/j.ajog.2021.03.016.
- 483 4. Jacob L, López-Sánchez GF, Oh H, Shin JI, Grabovac I, Soysal P, et al. Association of
484 multimorbidity with higher levels of urinary incontinence: a cross-sectional study of 23 089
485 individuals aged ≥ 15 years residing in Spain. *Br J Gen Pract*. 2021;71(702):e71-e7. doi:
486 10.3399/bjgp20X713921.
- 487 5. Nazarko L. Male urinary incontinence management: penile sheaths. *Br J Community Nurs*.
488 2018;23(3):110-6. doi: 10.12968/bjcn.2018.23.3.110.
- 489 6. Jachan DE, Müller-Werdan U, Lahmann NA. Impaired Mobility and Urinary Incontinence
490 in Nursing Home Residents: A Multicenter Study. *J Wound Ostomy Continence Nurs*.
491 2019;46(6):524-9. doi: 10.1097/won.0000000000000580.
- 492 7. Nam JY, Park EC, Cho E. Does Urinary Incontinence and Mode of Delivery Affect
493 Postpartum Depression? A Nationwide Population-Based Cohort Study in Korea. *Int J*
494 *Environ Res Public Health*. 2021;18(2). doi: 10.3390/ijerph18020437.

- 495 8. Fultz N, Girts T, Kinchen K, Nygaard I, Pohl G, Sternfeld B. Prevalence, management and
496 impact of urinary incontinence in the workplace. *Occup Med (Lond)*. 2005;55(7):552-7. doi:
497 10.1093/occmed/kqi152.
- 498 9. Bogner HR, Gallo JJ, Sammel MD, Ford DE, Armenian HK, Eaton WW. Urinary
499 incontinence and psychological distress in community-dwelling older adults. *J Am Geriatr*
500 *Soc*. 2002;50(3):489-95. doi: 10.1046/j.1532-5415.2002.50115.x.
- 501 10. Brown JS, Vittinghoff E, Wyman JF, Stone KL, Nevitt MC, Ensrud KE, et al. Urinary
502 incontinence: does it increase risk for falls and fractures? Study of Osteoporotic Fractures
503 Research Group. *J Am Geriatr Soc*. 2000;48(7):721-5. doi: 10.1111/j.1532-
504 5415.2000.tb04744.x.
- 505 11. Dokuzlar O, Koc Okudur S, Smith L, Soysal P, Yavuz I, Aydin AE, et al. Assessment of
506 factors that increase risk of falling in older women by four different clinical methods. *Aging*
507 *Clin Exp Res*. 2020;32(3):483-90. doi: 10.1007/s40520-019-01220-8.
- 508 12. Gray M. Incontinence-related skin damage: essential knowledge. *Ostomy Wound*
509 *Manage*. 2007;53(12):28-32.
- 510 13. Nuotio M, Jylhä M, Luukkaala T, Tammela TL. Urinary incontinence in a Finnish
511 population aged 70 and over. Prevalence of types, associated factors and self-reported
512 treatments. *Scand J Prim Health Care*. 2003;21(3):182-7.
- 513 14. Tamanini JT, Santos JL, Lebrão ML, Duarte YA, Laurenti R. Association between
514 urinary incontinence in elderly patients and caregiver burden in the city of Sao Paulo/Brazil:
515 Health, Wellbeing, and Ageing Study. *Neurourol Urodyn*. 2011;30(7):1281-5. doi:
516 10.1002/nau.21040.
- 517 15. Thom DH, Haan MN, Van Den Eeden SK. Medically recognized urinary incontinence
518 and risks of hospitalization, nursing home admission and mortality. *Age Ageing*.
519 1997;26(5):367-74. doi: 10.1093/ageing/26.5.367.

- 520 16. Coyne KS, Wein A, Nicholson S, Kvasz M, Chen CI, Milsom I. Economic burden of
521 urgency urinary incontinence in the United States: a systematic review. *J Manag Care Pharm.*
522 2014;20(2):130-40. doi: 10.18553/jmcp.2014.20.2.130.
- 523 17. Milsom I, Coyne KS, Nicholson S, Kvasz M, Chen CI, Wein AJ. Global prevalence and
524 economic burden of urgency urinary incontinence: a systematic review. *Eur Urol.*
525 2014;65(1):79-95. doi: 10.1016/j.eururo.2013.08.031.
- 526 18. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The
527 PRISMA statement for reporting systematic reviews and meta-analyses of studies that
528 evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.*
529 2009;62(10):e1-34. doi: 10.1016/j.jclinepi.2009.06.006.
- 530 19. Ioannidis JP. Integration of evidence from multiple meta-analyses: a primer on umbrella
531 reviews, treatment networks and multiple treatments meta-analyses. *Cmaj.* 2009;181(8):488-
532 93. doi: 10.1503/cmaj.081086.
- 533 20. Shea BJ, Reeves BC, Wells G, Thuku M, Hamel C, Moran J, et al. AMSTAR 2: a critical
534 appraisal tool for systematic reviews that include heterised or non-randomised studies of
535 healthcare interventions, or both. *Bmj.* 2017;358:j4008. doi: 10.1136/bmj.j4008.
- 536 21. IntHout J, Ioannidis JP, Borm GF. The Hartung-Knapp-Sidik-Jonkman method for
537 random effects meta-analysis is straightforward and considerably outperforms the standard
538 DerSimonian-Laird method. *BMC Med Res Methodol.* 2014;14:25. doi: 10.1186/1471-2288-
539 14-25.
- 540 22. Higgins JP, Thompson SG, Spiegelhalter DJ. A re-evaluation of random-effects meta-
541 analysis. *J R Stat Soc Ser A Stat Soc.* 2009;172(1):137-59. doi: 10.1111/j.1467-
542 985X.2008.00552.x.

- 543 23. IntHout J, Ioannidis JP, Rovers MM, Goeman JJ. Plea for routinely presenting prediction
544 intervals in meta-analysis. *BMJ Open*. 2016;6(7):e010247. doi: 10.1136/bmjopen-2015-
545 010247.
- 546 24. Serghiou S, Goodman SN. Random-Effects Meta-analysis: Summarizing Evidence With
547 Caveats. *Jama*. 2019;321(3):301-2. doi: 10.1001/jama.2018.19684.
- 548 25. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*.
549 2002;21(11):1539-58. doi: 10.1002/sim.1186.
- 550 26. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a
551 simple, graphical test. *Bmj*. 1997;315(7109):629-34. doi: 10.1136/bmj.315.7109.629.
- 552 27. Ioannidis JP, Trikalinos TA. An exploratory test for an excess of significant findings.
553 *Clin Trials*. 2007;4(3):245-53. doi: 10.1177/1740774507079441.
- 554 28. Ioannidis J. Clarifications on the application and interpretation of the test for excess
555 significance and its extensions. *Journal of Mathematical Psychology*. 2013;57:184–7. doi:
556 10.1016/j.jmp.2013.03.002.
- 557 29. Aromataris E, Fernandez R, Godfrey CM, Holly C, Khalil H, Tungpunkom P.
558 Summarizing systematic reviews: methodological development, conduct and reporting of an
559 umbrella review approach. *Int J Evid Based Healthc*. 2015;13(3):132-40. doi:
560 10.1097/xeb.0000000000000055.
- 561 30. Belbasis L, Savvidou MD, Kanu C, Evangelou E, Tzoulaki I. Birth weight in relation to
562 health and disease in later life: an umbrella review of systematic reviews and meta-analyses.
563 *BMC Med*. 2016;14(1):147. doi: 10.1186/s12916-016-0692-5.
- 564 31. Soysal P, Hurst C, Demurtas J, Firth J, Howden R, Yang L, et al. Handgrip strength and
565 health outcomes: Umbrella review of systematic reviews with meta-analyses of observational
566 studies. *J Sport Health Sci*. 2021;10(3):290-5. doi: 10.1016/j.jshs.2020.06.009.

- 567 32. Soysal P, Veronese N, Smith L, Torbahn G, Jackson SE, Yang L, et al. Orthostatic
568 hypotension and health outcomes: an umbrella review of observational studies. *Eur Geriatr*
569 *Med.* 2019;10(6):863-70. doi: 10.1007/s41999-019-00239-4.
- 570 33. Balzarro M, Rubilotta E, Mancini V, Trabacchin N, Oppeduzzi L, Li Marzi V, et al. Impact
571 of Overactive Bladder-Wet Syndrome on Female Sexual Function: A Systematic Review and
572 Meta-Analysis. *Sex Med Rev.* 2019;7(4):565-74. doi: 10.1016/j.sxmr.2019.05.002.
- 573 34. Lai S, Diao T, Zhang W, Seery S, Zhang Z, Hu M, et al. Sexual Functions in Women
574 With Stress Urinary Incontinence After Mid-Urethral Sling Surgery: A Systematic Review
575 and Meta-Analysis of Prospective Randomized and Non-Randomized Studies. *J Sex Med.*
576 2020;17(10):1956-70. doi: 10.1016/j.jsxm.2020.07.003.
- 577 35. Cheng S, Lin D, Hu T, Cao L, Liao H, Mou X, et al. Association of urinary incontinence
578 and depression or anxiety: a meta-analysis. *J Int Med Res.* 2020;48(6):300060520931348.
579 doi: 10.1177/0300060520931348.
- 580 36. Pizzol D, Demurtas J, Celotto S, Maggi S, Smith L, Angiolelli G, et al. Urinary
581 incontinence and quality of life: a systematic review and meta-analysis. *Aging Clin Exp Res.*
582 2021;33(1):25-35. doi: 10.1007/s40520-020-01712-y.
- 583 37. Beeckman D, Van Lancker A, Van Hecke A, Verhaeghe S. A systematic review and
584 meta-analysis of incontinence-associated dermatitis, incontinence, and moisture as risk
585 factors for pressure ulcer development. *Res Nurs Health.* 2014;37(3):204-18. doi:
586 10.1002/nur.21593.
- 587 38. John G, Bardini C, Combescure C, Dällenbach P. Urinary Incontinence as a Predictor of
588 Death: A Systematic Review and Meta-Analysis. *PLoS One.* 2016;11(7):e0158992. doi:
589 10.1371/journal.pone.0158992.

- 590 39. Coyne KS, Wein A, Nicholson S, Kvasz M, Chen CI, Milsom I. Comorbidities and
591 personal burden of urgency urinary incontinence: a systematic review. *Int J Clin Pract.*
592 2013;67(10):1015-33. doi: 10.1111/ijcp.12164.
- 593 40. Noguchi N, Chan L, Cumming RG, Blyth FM, Naganathan V. A systematic review of the
594 association between lower urinary tract symptoms and falls, injuries, and fractures in
595 community-dwelling older men. *Aging Male.* 2016;19(3):168-74. doi:
596 10.3109/13685538.2016.1169399.
- 597 41. Veronese N, Soysal P, Stubbs B, Marengoni A, Demurtas J, Maggi S, et al. Association
598 between urinary incontinence and frailty: a systematic review and meta-analysis. *Eur Geriatr*
599 *Med.* 2018;9(5):571-8. doi: 10.1007/s41999-018-0102-y.
- 600 42. Duralde ER, Rowen TS. Urinary Incontinence and Associated Female Sexual
601 Dysfunction. *Sex Med Rev.* 2017;5(4):470-85. doi: 10.1016/j.sxmr.2017.07.001.
- 602 43. Salonia A, Zanni G, Nappi RE, Briganti A, Dehò F, Fabbri F, et al. Sexual dysfunction is
603 common in women with lower urinary tract symptoms and urinary incontinence: results of a
604 cross-sectional study. *Eur Urol.* 2004;45(5):642-8; discussion 8. doi:
605 10.1016/j.eururo.2003.11.023.
- 606 44. Aslan G, Köseoğlu H, Sadik O, Gimen S, Cihan A, Esen A. Sexual function in women
607 with urinary incontinence. *Int J Impot Res.* 2005;17(3):248-51. doi: 10.1038/sj.ijir.3901296.
- 608 45. Wehbe SA, Kellogg S, Whitmore K. Urogenital complaints and female sexual
609 dysfunction. Part 2. *J Sex Med.* 2010;7(7):2304-17; quiz 18-9. doi: 10.1111/j.1743-
610 6109.2010.01951.x.
- 611 46. Costa RM, Brody S. Sexual satisfaction, relationship satisfaction, and health are
612 associated with greater frequency of penile-vaginal intercourse. *Arch Sex Behav.*
613 2012;41(1):9-10. doi: 10.1007/s10508-011-9847-9.

- 614 47. Ebrahim S, May M, Ben Shlomo Y, McCarron P, Frankel S, Yarnell J, et al. Sexual
615 intercourse and risk of ischaemic stroke and coronary heart disease: the Caerphilly study. *J*
616 *Epidemiol Community Health*. 2002;56(2):99-102. doi: 10.1136/jech.56.2.99.
- 617 48. Smith L, Yang L, Veronese N, Soysal P, Stubbs B, Jackson SE. Sexual Activity is
618 Associated with Greater Enjoyment of Life in Older Adults. *Sex Med*. 2019;7(1):11-8. doi:
619 10.1016/j.esxm.2018.11.001.
- 620 49. Bilgic D, Kizilkaya Beji N. How do urinary incontinence types affect sexual function and
621 quality of life for Turkish women? *Low Urin Tract Symptoms*. 2020;12(3):253-9. doi:
622 10.1111/luts.12314.
- 623 50. Handa VL, Harvey L, Cundiff GW, Siddique SA, Kjerulff KH. Sexual function among
624 women with urinary incontinence and pelvic organ prolapse. *Am J Obstet Gynecol*.
625 2004;191(3):751-6. doi: 10.1016/j.ajog.2003.11.017.
- 626 51. Nilsson M, Lalos O, Lindkvist H, Lalos A. Impact of female urinary incontinence and
627 urgency on women's and their partners' sexual life. *Neurourol Urodyn*. 2011;30(7):1276-80.
628 doi: 10.1002/nau.21039.
- 629 52. Reis AM, Brito LGO, Lunardi ALB, Pinto ESMP, Juliato CRT. Depression, anxiety, and
630 stress in women with urinary incontinence with or without myofascial dysfunction in the
631 pelvic floor muscles: A cross-sectional study. *Neurourol Urodyn*. 2021;40(1):334-9. doi:
632 10.1002/nau.24563.
- 633 53. Chiu MYL, Wong HT, Yang X. Distress Due to Urinary Problems and Psychosocial
634 Correlates among Retired Men in Hong Kong. *Int J Environ Res Public Health*. 2020;17(7).
635 doi: 10.3390/ijerph17072533.
- 636 54. Ge TJ, Vetter J, Lai HH. Sleep Disturbance and Fatigue Are Associated With More
637 Severe Urinary Incontinence and Overactive Bladder Symptoms. *Urology*. 2017;109:67-73.
638 doi: 10.1016/j.urology.2017.07.039.

- 639 55. Omli R, Hunskaar S, Mykletun A, Romild U, Kuhry E. Urinary incontinence and risk of
640 functional decline in older women: data from the Norwegian HUNT-study. *BMC Geriatr.*
641 2013;13:47. doi: 10.1186/1471-2318-13-47.
- 642 56. Wong SY, Hong A, Leung J, Kwok T, Leung PC, Woo J. Lower urinary tract symptoms
643 and depressive symptoms in elderly men. *J Affect Disord.* 2006;96(1-2):83-8. doi:
644 10.1016/j.jad.2006.05.013.
- 645 57. Nicolson P, Kopp Z, Chapple CR, Kelleher C. It's just the worry about not being able to
646 control it! A qualitative study of living with overactive bladder. *Br J Health Psychol.*
647 2008;13(Pt 2):343-59. doi: 10.1348/135910707x187786.
- 648 58. Liu X, Cao H, Zhu H, Zhang H, Niu K, Tang N, et al. Association of chronic diseases
649 with depression, anxiety and stress in Chinese general population: The CHCN-BTH cohort
650 study. *J Affect Disord.* 2021;282:1278-87. doi: 10.1016/j.jad.2021.01.040.
- 651 59. Soysal P, Veronese N, Thompson T, Kahl KG, Fernandes BS, Prina AM, et al.
652 Relationship between depression and frailty in older adults: A systematic review and meta-
653 analysis. *Ageing Res Rev.* 2017;36:78-87. doi: 10.1016/j.arr.2017.03.005.
- 654 60. Felde G, Engeland A, Hunskaar S. Urinary incontinence associated with anxiety and
655 depression: the impact of psychotropic drugs in a cross-sectional study from the Norwegian
656 HUNT study. *BMC Psychiatry.* 2020;20(1):521. doi: 10.1186/s12888-020-02922-4.
- 657 61. Kinjo M, Masuda K, Nakamura Y, Taguchi S, Tambo M, Okegawa T, et al. Effects on
658 Depression and Anxiety After Mid-Urethral Sling Surgery for Female Stress Urinary
659 Incontinence. *Res Rep Urol.* 2020;12:495-501. doi: 10.2147/rru.S270915.
- 660 62. Offermans MP, Du Moulin MF, Hamers JP, Dassen T, Halfens RJ. Prevalence of urinary
661 incontinence and associated risk factors in nursing home residents: a systematic review.
662 *Neurourol Urodyn.* 2009;28(4):288-94. doi: 10.1002/nau.20668.

- 663 63. Beele H, Smet S, Van Damme N, Beeckman D. Incontinence-Associated Dermatitis:
664 Pathogenesis, Contributing Factors, Prevention and Management Options. *Drugs Aging*.
665 2018;35(1):1-10. doi: 10.1007/s40266-017-0507-1.
- 666 64. Rodríguez-Palma M, Verdú-Soriano J, Soldevilla-Agreda JJ, Pancorbo-Hidalgo PL,
667 García-Fernández FP. Conceptual Framework for Incontinence-Associated Dermatitis Based
668 on Scoping Review and Expert Consensus Process. *J Wound Ostomy Continence Nurs*.
669 2021;48(3):239-50. doi: 10.1097/won.0000000000000754.
- 670 65. Matta R, Hird AE, Saskin R, Radomski SB, Carr L, Kodama RT, et al. Is There an
671 Association between Urinary Incontinence and Mortality? A Retrospective Cohort Study. *J*
672 *Urol*. 2020;203(3):591-7. doi: 10.1097/ju.0000000000000574.
- 673 66. Dokuzlar O, Koc Okudur S, Soysal P, Kocyigit SE, Yavuz I, Smith L, et al. Factors that
674 Increase Risk of Falling in Older Men according to Four Different Clinical Methods. *Exp*
675 *Aging Res*. 2020;46(1):83-92. doi: 10.1080/0361073x.2019.1669284.
- 676 67. Liu PS, Huang HK, Ding DC. Association of lower urinary tract symptoms and hip
677 fracture in adults aged ≥ 50 years. *PLoS One*. 2021;16(3):e0246653. doi:
678 10.1371/journal.pone.0246653.
- 679 68. Trevisan C, Bedogni M, Pavan S, Shehu E, Piazzani F, Manzato E, et al. The impact of
680 second hip fracture on rehospitalization and mortality in older adults. *Arch Gerontol Geriatr*.
681 2020;90:104175. doi: 10.1016/j.archger.2020.104175.
- 682 69. Kachru N, Holmes HM, Johnson ML, Chen H, Aparasu RR. Risk of Mortality Associated
683 with Non-selective Antimuscarinic medications in Older Adults with Dementia: a
684 Retrospective Study. *J Gen Intern Med*. 2020;35(7):2084-93. doi: 10.1007/s11606-020-
685 05634-3.
- 686 70. Mayrovitz HN, Sims N. Biophysical effects of water and synthetic urine on skin. *Adv*
687 *Skin Wound Care*. 2001;14(6):302-8. doi: 10.1097/00129334-200111000-00013.

- 688 71. Koudounas S, Mugita Y, Minematsu T, Nakagami G, Weller C, Sanada H. Does the
689 presence of bacterial urinary infection contribute to the development of incontinence-
690 associated dermatitis? A scoping review. *J Tissue Viability*. 2021;30(2):256-61. doi:
691 10.1016/j.jtv.2021.01.008.
- 692 72. Aminzadeh F, Edwards N, Lockett D, Nair R. Utilization of bathroom safety devices,
693 patterns of bathing and toileting and bathroom falls in a sample of community living older
694 adults. *Technology and Disability*. 2001;13. doi: 10.3233/TAD-2000-13202.
- 695 73. Wolf SL, Riolo L, Ouslander JG. Urge incontinence and the risk of falling in older
696 women. *J Am Geriatr Soc*. 2000;48(7):847-8. doi: 10.1111/j.1532-5415.2000.tb04765.x.
- 697 74. Foley AL, Loharuka S, Barrett JA, Mathews R, Williams K, McGrother CW, et al.
698 Association between the Geriatric Giants of urinary incontinence and falls in older people
699 using data from the Leicestershire MRC Incontinence Study. *Age Ageing*. 2012;41(1):35-40.
700 doi: 10.1093/ageing/afr125.
- 701 75. Green AR, Segal J, Boyd CM, Huang J, Roth DL. Patterns of Potentially Inappropriate
702 Bladder Antimuscarinic Use in People with Dementia: A Retrospective Cohort Study. *Drugs*
703 *Real World Outcomes*. 2020;7(2):151-9. doi: 10.1007/s40801-020-00181-z.
- 704 76. Wagg A, Gibson W, Ostaszkievicz J, Johnson T, 3rd, Markland A, Palmer MH, et al.
705 Urinary incontinence in frail elderly persons: Report from the 5th International Consultation
706 on Incontinence. *Neurourol Urodyn*. 2015;34(5):398-406. doi: 10.1002/nau.22602.
- 707 77. Danforth KN, Townsend MK, Lifford K, Curhan GC, Resnick NM, Grodstein F. Risk
708 factors for urinary incontinence among middle-aged women. *Am J Obstet Gynecol*.
709 2006;194(2):339-45. doi: 10.1016/j.ajog.2005.07.051.
- 710 78. Brown JS, Nyberg LM, Kusek JW, Burgio KL, Diokno AC, Foldspang A, et al.
711 Proceedings of the National Institute of Diabetes and Digestive and Kidney Diseases

- 712 International Symposium on Epidemiologic Issues in Urinary Incontinence in Women. *Am J*
713 *Obstet Gynecol.* 2003;188(6):S77-88. doi: 10.1067/mob.2003.353.
- 714 79. Nazzal Z, Khatib B, Al-Quqa B, Abu-Taha L, Jaradat A. The prevalence and risk factors
715 of urinary incontinence amongst Palestinian women with type 2 diabetes mellitus: A cross-
716 sectional study. *Arab J Urol.* 2020;18(1):34-40. doi: 10.1080/2090598x.2019.1699340.
- 717 80. Bani-issa WA, Halabi JO, Abdullah AR, Hasan HA, Raigangar VL. Prevalence and risk
718 factors for incontinence among Emirati women with diabetes. *J Transcult Nurs.*
719 2014;25(1):42-50. doi: 10.1177/1043659613503873.
- 720 81. Jackson SL, Scholes D, Boyko EJ, Abraham L, Fihn SD. Urinary incontinence and
721 diabetes in postmenopausal women. *Diabetes Care.* 2005;28(7):1730-8. doi:
722 10.2337/diacare.28.7.1730.
- 723 82. Wyndaele M, De Winter BY, Pelckmans P, Wyndaele JJ. Lower bowel function in
724 urinary incontinent women, urinary continent women and in controls. *Neurourol Urodyn.*
725 2011;30(1):138-43. doi: 10.1002/nau.20900.
- 726 83. Sultan AH, Monga A, Lee J, Emmanuel A, Norton C, Santoro G, et al. An International
727 Urogynecological Association (IUGA)/International Continence Society (ICS) joint report on
728 the terminology for female anorectal dysfunction. *Int Urogynecol J.* 2017;28(1):5-31. doi:
729 10.1007/s00192-016-3140-3.
- 730 84. Ratto C, Doglietto GB. *Fecal Incontinence: Diagnosis and Treatment.* 2007.
- 731 85. Kissane LM, Martin KD, Meyer I, Richter HE. Effect of darifenacin on fecal
732 incontinence in women with double incontinence. *Int Urogynecol J.* 2021;32(9):2357-63. doi:
733 10.1007/s00192-020-04369-3.
- 734 86. Kim H, Yoshida H, Hu X, Saito K, Yoshida Y, Kim M, et al. Association between self-
735 reported urinary incontinence and musculoskeletal conditions in community-dwelling elderly

736 women: a cross-sectional study. *Neurourol Urodyn.* 2015;34(4):322-6. doi:
737 10.1002/nau.22567.

738 87. Melsen WG, Bootsma MC, Rovers MM, Bonten MJ. The effects of clinical and statistical
739 heterogeneity on the predictive values of results from meta-analyses. *Clin Microbiol Infect.*
740 2014;20(2):123-9. doi: 10.1111/1469-0691.12494.

741

