



## Quantification of the temporal change of diffusion patterns following mild traumatic brain injury

Jing, M., McGinnity, TM., Coleman, SA., Fuchs, A., & Kelso, J. A. S. (2012). Quantification of the temporal change of diffusion patterns following mild traumatic brain injury. In *Unknown Host Publication* IEEE.

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Unknown Host Publication

**Publication Status:**  
Published (in print/issue): 10/06/2012

**Document Version**  
Author Accepted version

### General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

### Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk)

2012 International Joint Conference on Neural Networks

**Title**

Quantification of the temporal change of diffusion patterns following mild traumatic brain injury

**Authors**

M. Jing\*<sup>1</sup>, T. M. McGinnity<sup>1</sup>, S. Coleman<sup>1</sup>, A. Fuchs<sup>2</sup>, J.A.S. Kelso<sup>2</sup>

<sup>1</sup>Intelligent Systems Research Centre, School of Computing and Intelligent Systems, University of Ulster, UK.

<sup>2</sup>Human Brain and Behavior Laboratory, Center for Complex Systems and Brain Sciences, Florida Atlantic University, USA.

Despite the emerging applications of diffusion tensor imaging (DTI) [1,4], there is a lack of longitudinal investigation on the evolution of diffusion based patterns in mild traumatic brain injury (mTBI), which may help to predict the recovery and assist in studying the long term impact caused by mTBI. Although the newly developed DTI techniques, such as voxel-wise analysis, region of interest (ROI) based analysis and tractography [3], have demonstrated their capability to detect mTBI related regions, they are not effective solutions for quantification of the temporal changes of mTBI. Furthermore, these methods rely on statistical analysis based on reasonably large data samples, which may not be suitable for case studies, such as tracking recovery of function in an individual.

In this work, we proposed an effective approach for longitudinal DTI study of mTBI, in which a group based independent component analysis (GICA) [2] is employed to a group of diffusion scalar maps arranged according to their data collection time points post injury. The DTI datasets from three mTBI subjects and six control subjects were used and four types of diffusion scalar maps were calculated, including fractional anisotropy (FA), mean diffusivity, axial diffusivity and transverse diffusivity. The results show that the GICA decomposes the group maps into a number of spatially independent components (ICs). The desired component is the one that represents the common spatial pattern within the group data, which is separated from noise, artifacts and unknown spatial patterns. The timecourse obtained from GICA represents the distribution of the selected IC in the group data, which reveals the temporal changes of the diffusion patterns during recovery of mTBI. In addition, we proposed to define a ROI by a skeleton of mean FA maps created by FSL [5], which allows quantitative comparisons to be performed in tract-focused regions, further revealing the trends of recovery in ROI based analysis. Our approach fully utilizes the outcomes of the spatial components and timecourse from the GICA model and provides a way to quantify pathways of change in longitudinal studies of mTBI.

**References**

- [1] Basser, P.J., Basser, P.J., Mattiello, J., Le Bihan, D., 1994. MR diffusion tensor spectroscopy and imaging, *Journal of Biophys*, 66, 259-267.
- [2] Calhoun, V.D., Adali, T., Pearlson, G.D., Pekar, J. J., 2001. A method for making group inferences from functional MRI data using independent component analysis. *Human. Brain Mapping*, 14(3), 140-151.
- [3] Rutgers, D.R., Fillard, P., Parodot, G., Tadie, M., Lasjaunias, P., D. Ducreux, D. 2008. Diffusion tensor imaging characteristics of the corpus callosum in mild, moderate, and severe traumatic brain injury. *American Journal of Neuroradiology*, 29(9), 1730-1735.
- [4] Jing, M., McGinnity, T.M., Coleman, S., Zhang, H., Fuchs, A., Kelso, S., 2012. Enhancement of Fibre Orientation Distribution Reconstruction in Diffusion Weighted Imaging by Single Channel Blind Source Separation, *IEEE Trans. Bio. Eng.*, 59(2), 371-383.
- [5] Smith, S.M., Jenkinson, M., Woolrich, M.W., et al. 2004. Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*, 23(S1), 208-219.