



Data report on three datasets: Mortality patterns between agricultural and non-agricultural ward areas

Trearty, K., Bunting, B., & Mallett, J. (2023). Data report on three datasets: Mortality patterns between agricultural and non-agricultural ward areas. *Frontiers in Genetics*, 13, 1-7. [953167].
<https://doi.org/10.3389/fgene.2022.953167>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Frontiers in Genetics

Publication Status:
Published online: 04/01/2023

DOI:
[10.3389/fgene.2022.953167](https://doi.org/10.3389/fgene.2022.953167)

Document Version
Publisher's PDF, also known as Version of record

General rights
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.



OPEN ACCESS

EDITED BY

Domenica D'Elia,
Institute of Biomedical Technologies
(CNR), Italy

REVIEWED BY

Bassirou Bonfoh,
Swiss Centre for Scientific Research
(Côte d'Ivoire), Côte d'Ivoire
Bogdan Vasile Ileanu,
Bucharest Academy of Economic
Studies, Romania

*CORRESPONDENCE

Kelly Trearty,
✉ trearty_33@hotmail.com

†PRESENT ADDRESS

Kelly Trearty,
Frontiers Media SA, Lausanne,
Switzerland

SPECIALTY SECTION

This article was submitted to Statistical
Genetics and Methodology,
a section of the journal
Frontiers in Genetics

RECEIVED 25 May 2022

ACCEPTED 06 December 2022

PUBLISHED 04 January 2023

CITATION

Trearty K, Bunting B and Mallett J (2023),
Data report on three datasets: Mortality
patterns between agricultural and non-
agricultural ward areas.
Front. Genet. 13:953167.
doi: 10.3389/fgene.2022.953167

COPYRIGHT

© 2023 Trearty, Bunting and Mallett.
This is an open-access article
distributed under the terms of the
[Creative Commons Attribution License
\(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original
publication in this journal is cited, in
accordance with accepted academic
practice. No use, distribution or
reproduction is permitted which does
not comply with these terms.

Data report on three datasets: Mortality patterns between agricultural and non-agricultural ward areas

Kelly Trearty ^{†*}, Brendan Bunting and John Mallett

Department of Psychology, Ulster University, Coleraine, United Kingdom

The health of the farming community in Northern Ireland (NI) requires further research as previous mortality studies have reported contradictory results regarding farmers' health outcomes compared against other occupations and the general population. This study collated the NINIS area-level farm census with the population census information across 582 non-overlapping wards of NI to compile three mortality datasets (2001, 2011, and pooled dataset) (NISRA 2019). These datasets allow future researchers to investigate the influence of demographic, farming, and economic predictors on all-cause mortality at the ward level. The 2001 and 2011 mortality datasets were compiled for cross-sectional analyses and subsequently pooled for longitudinal analyses. Findings from these datasets will provide evidence of the influence of Farming Intensity scores influence on death risk within the wards for future researchers to utilise. This data report will aid in the understanding of socio-ecological variables' additive contribution to the risk of death at the ward level within NI. This data report is of interest to the One Health research community as it standardises the environment–human–animal data to pave the way towards a new One Health research paradigm. For example, future researchers can use this nationally representative data to investigate whether agriculturally saturated wards have a higher mortality risk than non-agriculturally based wards of NI.

KEYWORDS

death trends, aggregated data, area-level data, farm census, population census, Northern Ireland, Northern Ireland Neighbourhood Service (NINIS), agricultural versus non-agricultural wards

Introduction

Agricultural workers have the highest work intensification of all occupations as they merge home-life with work-life and in principle are at their enterprise 365 days per year (Furey et al., 2016). The health of the farming community requires further research as previous mortality studies have reported contradictory results regarding farmers' health outcomes compared against other occupations and the general population (Stiernström et al., 2001; Smyth et al., 2012; Cushen et al., 2016). It is still unclear which underlying

mechanisms are increasing farmers' mortality rates; therefore, this study's compilation of the three datasets allows for various types of analyses and aims to fill some of this research gap.

The economy of Northern Ireland (NI) is dominated by the agricultural industry, as farming sectors employ approximately 1.2 million hectares or three-quarters of its acreage (NISRA 2019). In addition, NI's physical landscape is diverse, ranging from high mountains and fertile plains to a jagged coastline, making it ideal for accommodating a vast range of agricultural activities (Abson et al., 2013). The number of NI farmers (farmers, spouses, and other farm workers) in 2001 ($n = 56,362$) decreased by 17% in 2011 ($n = 46,848$) (NISRA 2019), thus following global trends in farm number reductions (Smyth et al., 2012).

Furthermore, farmers aged between 16 and 64 years old accounted for 10% in 2001 and 7% in 2011 of the economically active NI population (HSENI, 2019b; NISRA 2019). The number of deaths in NI decreased by 2% between 2001 ($n = 14,513$) and 2011 ($n = 14,204$) (NISRA 2019). Ireland's fatality rate in agriculture, forestry, and fishing is 5–7 times greater than any other Irish economic sector (Phelan 2018; Health and Safety Authority 2021). Farming had an average of eight on-farm deaths per year in NI until 2010; however, this increased by 50% to 12 fatalities in 2011–2012 and continues to increase (HSENI, 2019a).

Building on these statistics and the recommendations of previous research (Havelka et al., 2009; Som 2010; Allik et al., 2016; Wilkinson et al., 2016), this study linked the 2001 and 2011 population census with the farm census information across 582 non-overlapping wards of NI (NISRA 2013). This was approached by utilising the Northern Ireland Neighbourhood Information Service (NINIS) online repository to compile three mortality datasets (2001, 2011, and pooled datasets) (NISRA 2019). The current study's database set-up emulated that of Luke and Krauss (2004), converting area-level variables into the proportion of their summed total for ease of interpretation. Findings from these datasets will provide evidence of the influence of Farming Intensity scores on death risk within wards for future researchers to utilise, and will aid in the understanding of socio-ecological variables and their additive contribution to the risk of deaths at the ward level within NI.

Data and methods

The NINIS is a web-based service that hosts nationally representative population census and farm census information within its open-data portal (NISRA, 2019). These data were publicly available, and in January of 2019, ethical approval was granted by Ulster University Research Ethics Committee.

Table 1 identifies the origin index codes for the underlying data of 2001 and 2011 datasets (NISRA 2019). This study downloaded each variable 1) at the ward level, 2) in count form, and 3) for the census years of 2001 and 2011.

Ten of the 17 variables were transformed to emulate a dichotomized measure (*age, males, below degree, living alone, limiting long-term illness, unpaid care, full-time workers, farmers, and grass*). Building on the previous research (Ruiz-Martinez et al., 2015), Farming Intensity was computed as a composite factor score to assess agricultural activity within each ward, using six indicator items from the farm census (farms, farmers, grass, cattle, pigs, and poultry) (NISRA 2019).

Deaths: The outcome variable was *all-cause mortality*, which represented the total number of *deaths* observed over a 12-month period. The causes of death included in this mortality variable were as follows: malignant neoplasms C00-C97 (e.g. cancers, etc.); circulatory diseases I00-I97 (e.g. heart disease, stroke, blood clots, etc.); respiratory diseases J00-J99 (e.g. pneumonia, asthma, bronchitis, influenza, etc.); and external causes V01-Y98, X60- X84, Y10-Y34, Y87.0, Y87.2 (e.g. road traffic accidents, falls, suicide, undetermined intent, fires, poisoning, assault etc.). The *death* variable for each ward remained in its raw form, as a count.

Ward Population: The 582 wards contained different population sizes ranging from approximately 740–9,500 residents. To operationalise its usefulness, the *natural log of the population* was calculated from the *ward population* variable (McGranahan 1999). This study expected wards with larger *natural log of the population* scores to be at a higher risk of *death* than wards with lower *natural log of the population* scores.

Age: Derived from the date of birth question, an individual's *age* was their age on their last birthday by that census day. *Age* was available from the NINIS databank in 2011 as a continuum (0–100+ year-olds in single years) and in 2001 as seven structured age group variables (categorised at different life stages). It was compiled into those aged *below 64 years* and *over 65 years*. This study hypothesised that wards with larger amounts of residents aged *65 to 100+ years* were at a higher risk of *death* than wards with higher amounts of *below 64-year-olds* (Doebler & Glasgow 2016; Lee et al., 2017; van Doorn et al., 2018). *Age* was converted into the proportion of *65 to 100+ year-olds* in a ward relative to the total resident population per ward.

Gender was self-reported as *male* or *female* on the census day. This study postulated that wards with larger amounts of *male* inhabitants were at a higher risk of *mortality* than wards with higher amounts of *females*. The number of *males* per ward was converted into their proportion relative to the total resident population per ward.

Marital Status represented a person's legal marital status on the census day. It was dichotomized into *living alone* (summed: single, separated, divorced, and widowed) and *cohabitating* (summed: married, remarried, and same sex civil partnership). This study posited that wards with larger amounts of people *living alone* were at a higher risk of *death* than wards with higher amounts of dwellers *cohabitating*. The number of those *living alone* was converted into its proportion relative to the residents 16 years and older per ward.

TABLE 1 Themes, subsets, names, and codes of raw variables extracted from the NINIS website to create 2001 and 2011 mortality datasets.

Theme	Subset	Variable	Codes	
			2011	2001
Population	Deaths	Deaths	2011	2001
Census	Demography	Ward population	QS103NI	KS02
		Age	QS103NI	KS02
		Gender	QS105NI	KS01
		Marital status	KS103NI	KS04
		Qualifications	Qualifications and students	KS501NI
	Labour market	Hours worked	DT602NI	UV041
	Health	Unpaid care	QS301NI	UV021
		Limiting long-term illness	QS303NI	UV022
	Agriculture	Farm census	Farms	2011
Area farmed			2011	2001
Grass			2011	2001
Cattle			2011	2001
Pigs			2011	2001
Poultry			2011	2001
Agricultural labour			2011	2001
Average standard output/standard gross margin			2010	2001
People and places			Community	Ward size (hectares)
Deprivation	NIMDM	Northern Ireland Multiple Deprivation Measure	2010	2005

Note: codes provide unique reference codes for each variable per year. *Deprivation* reflected the overall weighed score of its seven deprivation domains: i) income, ii) employment, iii) health, iv) education, v) proximity to services, vi) environment, and vii) crime.

Qualifications represented the highest level of professional and vocational education one obtained by the census day. It was dichotomised into *below degree* (summed: no qualifications, level 1, level 2, apprenticeship, level 3, and others) and *above degree* (level 4 and level 5). This study theorized that wards with larger amounts of *below degree* qualifications were at a higher risk of *mortality* than wards with higher amounts of *above degree* qualifications. The total number of individuals with *below degree* qualifications was converted into its proportion relative to residents over 16 years old per ward.

Limiting long-term illness (LLTI): It was self-reported on whether an individual had an *LLTI* or disability lasting or expected to last a minimum of 12 months from the census day. It was used to create *no LLTIs* and *yes LLTIs present* (summed: daily activities limited a little; daily activities limited a lot). The eleven types of *LLTIs* encapsulated within this variable were a combination of physical and mental health conditions. The six physical health issues were: long-term pain; mobility issues; deaf or partial hearing; blind or visually impaired; breathing issues; and chronic illness (e.g. cancer, diabetes, etc.). The five mental health issues were: communication

difficulties; mental health issues (e.g. depression, schizophrenia, etc.); memory loss issues; learning or behavioural difficulties; and other conditions. This study presupposed that wards with larger amounts of residents with *LLTIs* were at a higher risk of *death* than wards with higher amounts of residents with *no LLTIs*. The number of people with an *LLTI* was converted into its proportion relative to the total number of residents per ward.

Unpaid carers represented the number of hours one provides unpaid care to a care recipient (family members, friends, neighbours, and others) weekly because of their poor physical/mental health or old age. It was dichotomised into *0 h (no care provided)* and *1 or more hours* (summed: 1–19 h, 20–49 h, and 50+ hours) of unpaid care provided weekly. This study considered that wards with larger amounts of *unpaid carers* were at a higher risk of *mortality* than wards with higher amounts of *no unpaid carers*. The total number of people providing *unpaid care (1 or more hours)* was converted into its proportion relative to the total residents per ward.

Hours worked weekly represented the number of individuals aged between 16 and 74 years who worked the week before the census in their main job, including paid and unpaid overtime. It

was dichotomised into *part-time workers* (summed: 1–15 h; 16–30 h) and *full-time workers* (summed: 31–48 h; 49+ hours). This study regarded that wards with larger amounts of inhabitants working *full-time hours* were at a higher risk of *death* than wards with higher amounts of *part-time workers*. The total number of people working *full-time hours* was converted into its proportion relative to the employed residents aged 16–74 years per ward.

Northern Ireland Multiple Deprivation Measure (NIMDM) reflected the overall weighed score of its seven deprivation domains. Each of the seven dimensions had different influences relating to an area's disadvantage, such as 1) income (25%); 2) employment (25%); 3) health (15%); 4) education (15%); 5) proximity to services (10%); 6) environment (5%); and 7) crime (5%). NIMDM provided an indication of the spatial rank order for each area from the most deprived (1) to the least deprived (582) ward. This measure was able to identify if one area had higher/lower deprivation than another, but it was not able to quantify how much more/less deprived one area was to another area. This study surmised that wards with lower NIMDM rankings (more deprivation in that area) were at a higher risk of *mortality* than those with higher NIMDM rankings (less deprivation in that area). The overall 2005 and 2010 NIMDM deprivation rankings were utilised as they corresponded with the census years under analysis. NIMDM remained unchanged in their raw numerical form.

Average standard gross margin (SGM) represented the annual farming profit earned from arable and/or livestock production per ward. The average SGM changed in 2011 to the average standard output (SO) within the farm census. Therefore, an approximate equivalence was assumed by including the 2001 and 2010 SGM information within datasets as a proxy for farm profit. The SGM represented the average profit earned from farm production within each ward (output minus production costs) and was expressed in European Size Unit (or euros). One SGM unit represented €1,200, which can be used to indicate the economic size of farming outputs within wards (NISRA, 2019). This study expected that wards with lower *farming profit* were at a higher risk of *death* than wards with higher *farming profit*. The 2001 and 2010 variables remained unchanged in their count form per ward.

Farms represented the total number of active farm holdings per ward from the farm census. Active farms were defined as spatial units encapsulating 1+ hectare(s) of farmland, used for crops and/or livestock production. This study hypothesised that wards with larger amounts of *farms* were at a higher risk of *death* than wards with no farms. *Farms* remained as a count per ward.

Agricultural labour (farmers) represented the total number of active farmers (16–100+ years old) employed within the agricultural industry for 20 or more weeks (39%) per year, from the farm census. This study merged farmers, spouses, and other workers to create the summed total of *agricultural labourers (or farmers)* per ward. This study postulated that wards with larger amounts of *farmers* were at a higher risk of *death* than wards with no

farmers. The total number of *farmers* was converted into its proportion relative to the total population (16–100+ years old) per ward.

Grass: NINIS defines the farming practice of producing *grass* within the farm census as the total grass in hectares, excluding rough grazing, farm woodlands, and non-agricultural land. This study posited that wards with larger amounts of *grass-based farming* were at a higher risk of *death* than wards with no grass farming. It was converted into the proportion of *grass* hectares relative to the *total hectare area farmed* per ward.

Cattle represented the total number of cattle in each ward, from the Farm Census. This study theorised that wards with larger amounts of *cattle farming* were at a higher risk of *death* than wards with no *cattle*.

Pigs represented the total number of pigs per ward, from the farm census. This study presupposed that wards with larger amounts of *pig farming* were at a higher risk of *death* than wards with no *pig farming*.

Poultry represented the total number of poultries in each ward, from the farm census. This study surmised that wards with larger amounts of *poultry farming* were at a higher risk of *death* than wards with no *poultry farming*.

The livestock variables (number of *cattle*, *pigs*, and *poultry* within each ward) were rescaled from large continuous variables into four-level categorical variables (scored 0–3). Wards with 0–3 head of each livestock variable were recoded as 0 to represent *none of that animal* within that ward. The remaining amount of each livestock variable was divided by 3 to create equally banded categories across wards, representing small (1), medium (2), and large (3) quantities of each animal per ward.

Dataset compilation

The Statistical Package for Social Science (SPSS, Version 25) software was used to construct the three mortality-based datasets (IBM Corp, 2017). The ward-level variables were individually extracted in count form *via* an Excel spreadsheet, imported into SPSS-25 to collate the data, and then imported into MPlus 8.1 (Muthén and Muthén, 1998–2019) for further analysis (see supplementary material: NINIS data extraction procedure). The three datasets were set up in wide format. Geographical locations of wards were used as the linking identifier *via* a deterministic one-to-one matching algorithm (Zhu et al., 2015). The farm census variables formed the spine of the 2001 and 2011 datasets. All other variables from the population census were downloaded in an identical format to ensure harmonious merging and configuration. Data cleaning and missing value analyses indicated that there were no missing data present within the variables (Dziadkowiec et al., 2016).

Strengths

The NINIS digital ecosystem stores agricultural and administrative records (collected routinely for different purposes) independently and in isolation, which minimises the usefulness of the data. The current study provides evidence that the NINIS information is findable, accessible, interoperable, reusable, and expandable (Table 1). The innovative design of these datasets encourages future researchers to utilise publicly available macro and/or micro data to unlock secondary value from the data.

This paper presents information on NINIS data in a way that is accessible to readers who may not share the author's particular specialism. In addition, this research will encourage future investigators to conduct interdisciplinary research as no other study has applied this approach, combining NINIS agricultural (agro-ecological science) and administrative data (health science) to facilitate new research questions within NI wards (Morrison et al., 1993). Conclusions drawn from these multiple data sources will be stronger with a reduced risk of error, in contrast to those drawn from only a single source.

This study used the two most up-to-date population censuses (2001 and 2011) and then collated the farm census data from the same years. The two distinct periods (2001 and 2011) covered identical areas but are not necessarily the same people, while the 2011 database provides the most up-to-date information currently available. This paper provides the methodology on how to analyse the next census once it becomes available online. However, the current census for 2021 has been delayed due to COVID-19 and has been rescheduled for enumeration in April 2022. The authors of this study intend to merge the next census information (once available on the NINIS website) to the current datasets.

The adjustment of the SGM to SO in 2011 does not affect the current study's reproducibility. Therefore, an extension of this study may utilise SGM (2001 and 2010) and SO (2011 and 2021) variables to compare the differences in farming outputs over 10 years as this would allow one to analyse the changes in the agricultural economic size between wards. NI policymakers within the Department of Agriculture, Environment, and Rural Affairs (DAERA), National Health Service (NHS), and Health and Safety Executive Northern Ireland (HSENI) could collaboratively use these datasets to identify agriculturally based wards (with higher deaths in 2011 than in 2001) and tailor farmers' healthcare needs accordingly.

Limitations

As a result of NISRA's aggregation procedures, a small amount of data loss occurred during the summarisation of

information. Moreover, the three datasets within this paper do not possess the ability to analyse the associations of individual-level variables, as the information used for compilation was aggregated (or clustered) at the ward level. Lokar et al. (2019) stated that this means researchers must be cautious to avoid making an *ecological fallacy* when interpreting results, bearing in mind that individual- and ward-level predictor associations with deaths are different (Robinson 1950). Jacob (2016) reported that some researchers remain hesitant to analyse the aggregated data as there is an underlying misconception that clustered data have restricted use; however, NINIS aggregated data results will provide the benefits of efficiency and power. Moreover, despite these variables being aggregated at the ward level, there were robust and vast amounts of literature supporting their use within this study of mortality (Haan et al., 1987; Yen & Kaplan, 1999; Chan et al., 2014).

Conclusion

In order to fill the gaps in the current research, this data report collated the NINIS information at ward level from the population and farm censuses (2001 and 2011) to compile three mortality-based datasets (NISRA 2019). The 2001 and 2011 mortality datasets were compiled for future cross-sectional analyses and subsequently pooled into a third dataset for longitudinal analyses. This study expanded on the design, methodology, and analytic limitations of a neighbourhood social environment study conducted by Yen and Kaplan (1999). This NI study was conducted in response to: i) corresponding reductions in the number of deaths of farmers between census years and ii) no nationally representative datasets available with the ability to analyse economic, environmental, and social influences on mortality patterns between agricultural and non-agricultural wards.

The purpose of this paper is to aid researchers who would like to use these datasets for future analyses by making the methodology that underpins it transparent and easy to replicate. For example, future researchers can use these data to investigate whether agriculturally saturated wards have a higher mortality risk than non-agriculturally based wards of NI. For didactic purposes, this data report provides the descriptive, structural, and administrative metadata (Davenhall 2011) of the datasets so that external investigators who want to replicate, reuse, adapt, or expand it can do so competently by utilising this approach.

This data report is of interest to the One Health research community as it standardizes the environment-human-animal

data to pave the way towards a new One Health research paradigm. In addition, these three datasets use novel data linkage methods, and their integrative analyses will enhance the understanding of the interconnectedness of the environment and multiple species within NI. Furthermore, the current amalgamation of data within this paper allows for an early release of summary results, alongside an extensive array of academic research uses.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary Material; further inquiries can be directed to the corresponding author.

Author contributions

KT used these datasets in her PhD 2021. BB was the primary and JM was the secondary supervisor, and they oversaw each element within the databases.

References

- Abson, D. J., Fraser, E. D., and Benton, T. G. (2013). Landscape diversity and the resilience of agricultural returns: A portfolio analysis of land-use patterns and economic returns from lowland agriculture. *Agric. Food Secur.* 2 (2), 2–15. doi:10.1186/2048-7010-2-2
- Allik, M., Brown, D., Dundas, R., and Leyland, A. H. (2016). Developing a new small-area measure of deprivation using 2001 and 2011 census data from Scotland. *Health Place* 39 (1), 122–130. doi:10.1016/j.healthplace.2016.03.006
- Chan, K. S., Roberts, E., McCleary, R., Buttorff, C., and Gaskin, D. J. (2014). Community characteristics and mortality: The relative strength of association of different community characteristics. *Am. J. Public Health* 104 (9), 1751–1758. doi:10.2105/AJPH.2014.301944
- Cushen, B., Sulaiman, I., Donoghue, N., Langan, D., Cahill, T., Nic Dhonncha, E., et al. (2016). High prevalence of obstructive lung disease in non-smoking farmers: The Irish farmers lung health study. *Respir. Med.* 115, 13–19. doi:10.1016/j.rmed.2016.04.006
- Davenhall, C. (2011). DCC I Digital curation reference manual. [Online] Available at: https://www.dcc.ac.uk/sites/default/files/documents/Scientific%20Metadata_2011_Final.pdf.
- Doebler, S., and Glasgow, N. (2016). Relationship between deprivation and the self-reported health of older people in Northern Ireland. *J. Agine Health* 2016 1–33. doi:10.1177/0898264316641079
- Dziadkowiec, O., Callahan, T., Ozkaynak, M., Reeder, B., and Welton, J. (2016). Using a data quality framework to clean data extracted from the electronic health record: A case study. *eGEMs (Generating Evid. Methods improve patient outcomes)* 4 (1), 1201–1215. doi:10.13063/2327-9214.1201
- Furey, E. M., O'Hora, D., McNamara, J., Kinsella, S., and Noone, C. (2016). The roles of financial threat, social support, work stress, and mental distress in dairy farmers' expectations of injury. *Front. Public Health* 4 (126), 126–211. doi:10.3389/fgene.2016.00126
- Haan, M., Kaplan, G. A., and Camacho, T. (1987). Poverty and health: Prospective evidence from the Alameda county study. *Am. J. Epidemiol.* 125 (6), 989–998. doi:10.1093/oxfordjournals.aje.a114637
- Havelka, M., Lučanin, J. D., and Lučanin, D. (2009). Biopsychosocial Model: The integrated approach to health and disease. *Coll. Antropol.* 33 (1), 303–310.
- Health and Safety Authority (2021). Agriculture and forestry. [Online] Available at: https://www.hsa.ie/eng/Your_Industry/Agriculture_Forestry/Further_Information/Fatal_Accidents/.
- HSENI (2019a). Farm safety week statistics. [Online] Available at: <https://www.hseni.gov.uk/news/farm-safety-week-statistics>.
- HSENI (2019b). HSENI annual report and accounts 2018-2019: Controlling risk together. Available at: https://www.hseni.gov.uk/sites/hseni.gov.uk/files/hseni-annual-report-and-statement-of-accounts-2018-19_2.pdf
- IBM Corp (2017). *IBM SPSS statistics for windows*. Armonk: IBM Corp.
- Jacob, R. (2016). OPRE Report #2016-91. Washington DC: OPRE.Using aggregate administrative data in social policy research
- Lee, M. J., Cawley, D. T., Ng, J. P., and Kaar, K. (2017). Trends in fractures and fatalities of farmyard injuries in Ireland: A 10 year analysis. *Ir. Med. J.* 1–5, 494.
- Lokar, K., Zagar, T., and Zadnik, V. (2019). Estimation of the ecological fallacy in the geographical analysis of the association of socio-economic deprivation and cancer incidence. *Int. J. Environ. Res. Public Health* 16 (3), 296–318. doi:10.3390/ijerph16030296
- Luke, D., and Krauss, M. (2004). Where there's smoke there's money: Tobacco industry campaign contributions and U.S. Congressional voting. *Am. J. Prev. Med.* 27, 363–372. doi:10.1016/j.amepre.2004.08.014
- McGranahan, D. A. (1999). *Natural amenities drive rural population change*. Washington: Department of Agriculture. doi:10.22004/ag.econ.33955
- Morrison, H., Savitz, D., Semenciw, R., Hulka, B., Mao, Y., Morison, D., et al. (1993). Farming and prostate cancer mortality. *Am. J. Epidemiol.* 137 (3), 270–280. doi:10.1093/oxfordjournals.aje.a116674
- Muthén, L. K., and Muthén, B. O. (1998-2019). *Mplus user's guide*. Eighth Edition. Los Angeles(CA): Muthén & Muthén.
- NISRA (2019). NINIS northern Ireland neighbourhood information service. [Online] Available at: <https://www.ninis2.nisra.gov.uk/public/Home.aspx>.
- NISRA (2013). *NISRA geography fact sheet: Northern Ireland neighbourhood information system*. Belfast: NISRA.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fgene.2022.953167/full#supplementary-material>

- Phelan, S. (2018). Fatalities in agriculture '5 to 7 times greater' than all other sectors. [Online] Available at: <https://www.agriland.ie/farming-news/agricultural-workplace-risks-analysed-over-14-year-period-in-new-research/>.
- Robinson, W. S. (1950). Ecological correlations and the behavior of individuals. *Am. Sociol. Rev.* 15 (3), 351–357. doi:10.2307/2087176
- Ruiz-Martinez, I., Marraccini, E., Debolini, M., and Bonari, E. (2015). Indicators of agricultural intensity and intensification: A review of the literature. *Ital. J. Agron.* 10 (656), 74–84. doi:10.4081/ija.2015.656
- Smyth, B., Evnas, D. S., Kelly, A., Cullen, L., and Donovan, D. O. (2012). The farming population in Ireland: Mortality trends during the "Celtic Tiger" years. *Eur. J. Public Health* 2012, 50–55. doi:10.1093/eurpub/cks017
- Som, H. (2010). *World programme for the census of agriculture 2010*. Rome, Italy: FAO Statistics Division.
- Stiernström, E.-L., Holmberg, S., Thelin, A., and Svärdsudd, K. (2001). A prospective study of morbidity and mortality rates among farmers and rural and urban nonfarmers. *J. Clin. Epidemiol.* 54 (2), 121–126. doi:10.1016/S0895-4356(00)00287-0
- van Doorn, D., Richardson, N., Storey, A., Osborne, A., Cunningham, C., Blake, C., et al. (2018). Farming characteristics and self-reported health outcomes of Irish farmers. *Occup. Med.* 68 (3), 199–202. doi:10.1093/occmed/kqy020
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., et al. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Sci. Data* 3 (1), 160018–160019. doi:10.1038/sdata.2016.18
- Yen, I. H., and Kaplan, G. A. (1999). Neighborhood social environment and risk of death: Multilevel evidence from the alameda county study. *Am. J. Epidemiol.* 149 (10), 898–907. doi:10.1093/oxfordjournals.aje.a009733
- Zhu, Y., Matsuyama, Y., Ohashi, Y., and Setoguchi, S. (2015). When to conduct probabilistic linkage vs. deterministic linkage? A simulation study. *J. Biomed. Inf.* 56, 80–86. doi:10.1016/j.jbi.2015.05.012