



Fish consumption and polyunsaturated fatty acid status of pregnant women in the Seychelles Child Development and Nutrition Study Cohort 2

Yeates, A. J., Mulhern, M. S., Ueland, P. M., McSorley, E. M., Wallace, J. M. W., Watson, G. E., Myers, G. E., Van Wijngaarden, E., Shamlaye, C., Davidson, P. W., & STRAIN, J.J. (2013). Fish consumption and polyunsaturated fatty acid status of pregnant women in the Seychelles Child Development and Nutrition Study Cohort 2. In *Unknown Host Publication* (Vol. 72 (OC)). Cambridge University Press. Advance online publication. <https://doi.org/10.1017/S0029665113001183>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Unknown Host Publication

Publication Status:
Published online: 13/05/2013

DOI:
[10.1017/S0029665113001183](https://doi.org/10.1017/S0029665113001183)

Document Version
Publisher's PDF, also known as Version of record

General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

Scottish Section Meeting, 18–19 March 2013, Polyunsaturated fatty acid mediators: implications for human health

Fish consumption and polyunsaturated fatty acid status of pregnant women in the Seychelles Child Development and Nutrition Study Cohort 2

A. J. Yeates¹, M. S. Mulhern¹, E. M. McSorley¹, J. M. W. Wallace¹, G. E. Watson², G. J. Myers², E. van Wijngaarden², C. Shamlaye³, J. Henderson³, S. W. Thurston², P. W. Davidson² and J. J. Strain¹

¹Northern Ireland Centre for Food and Health, University of Ulster, Coleraine, BT52 1SA, ²University of Rochester, School of Medicine and Dentistry, NY, USA and ³Ministry of Health, Victoria, Mahé, Republic of Seychelles

Fish is the major dietary source of long chain polyunsaturated fatty acids (PUFA), including docosahexaenoic acid (DHA), which is essential for optimal child neurodevelopment⁽¹⁾. In the first nutrition cohort of the Seychelles Child Development Study (SCDS), we showed no significant relationship between fish consumption (average 12 meals per week) and DHA status of pregnant women⁽²⁾. We hypothesised that altered fatty acid metabolism during pregnancy could be obscuring any correlation between dietary intake data and physiological measures. In the present study, we repeated the analysis to look at associations between fish consumption and PUFA status in the larger second nutrition SCDS cohort.

Fish consumption was assessed as frequency of fish meals consumed per week using a fish use questionnaire (FUQ) administered during pregnancy. Lipid extractions were carried out on serum samples collected at 28 weeks gestation following the Folch *et al.* method⁽³⁾ and fatty acid methyl esters were determined by GC-MS, as previously described⁽²⁾. A total of $n = 1474$ women completed the FUQ and provided a blood sample for PUFA measurement. Descriptive statistics and Pearson correlation analysis was carried out using IBM SPSS Statistics *v.* 20.

| | Mean | SD | 5 th , 95 th centile |
|-----------------------|--------|-------|--|
| Fish meals per week | 8.59 | 4.58 | 3.00, 17.00 |
| <i>n-6 PUFA mg/ml</i> | | | |
| LA | 0.902 | 0.255 | 0.526, 1.340 |
| AA | 0.203 | 0.082 | 0.061, 0.334 |
| Total <i>n-6</i> | 1.105 | 0.295 | 0.659, 1.608 |
| <i>n-3 PUFA mg/ml</i> | | | |
| ALA | 0.037 | 0.006 | 0.035, 0.047 |
| EPA | 0.052 | 0.008 | 0.049, 0.069 |
| DHA | 0.185 | 0.086 | 0.049, 0.329 |
| Total <i>n-3</i> | 0.274 | 0.091 | 0.147, 0.438 |
| <i>n-6:n-3</i> ratio | 4.358* | 1.636 | 2.506, 7.697 |

* Values were significantly correlated with fish meals per week in Pearson correlation ($P < 0.05$)

The mean age of pregnant women at enrolment to the study was 26 (SD = 6) years and the mean number of fish meals consumed per week during pregnancy was reported as 8.59 (SD = 4.58). Fish consumption was not significantly correlated with either individual or summed concentrations of *n-3* or *n-6* PUFA; however it was significantly negatively associated with the ratio of *n-6:n-3* ($r = -0.06$, $P = 0.04$).

These results agree with those of a previous SCDS cohort, where fish consumption, albeit assessed by food diary and in a smaller cohort ($n = 229$)⁽²⁾, was not significantly associated with serum PUFA status in pregnant women. This finding highlights a potential limitation of using fish consumption data as a biomarker for PUFA status, particularly during pregnancy. Further studies to identify a more robust biomarker for fish consumption are warranted.

1. Carlson SE & Neuringer (1999) *Lipids* **34**, 171–178.
2. Bonham MP, Duffy EM, Wallace JMW *et al.* (2008) *Prostag Leukotr Ess* **78**, 343–350.
3. Folch J, Lees M & Stanley GH (1957) *J Biol Chem* **226**, 497–509.