



Is there a common pathway to developing ASD and PTSD symptoms?

Hansen, M., Armour, C., Wittmann, L., Elklit, A., & Shevlin, M. (2014). Is there a common pathway to developing ASD and PTSD symptoms? *Journal of Anxiety Disorders*, 28(8), 865-872.
<https://doi.org/10.1016/j.janxdis.2014.09.019>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Journal of Anxiety Disorders

Publication Status:
Published (in print/issue): 01/01/2014

DOI:
[10.1016/j.janxdis.2014.09.019](https://doi.org/10.1016/j.janxdis.2014.09.019)

Document Version
Author Accepted version

General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

Table 1

Descriptive statistics and correlations between control factors, risk factors, and dependent factors (Pearson's r).

	<i>M</i>	<i>SD</i>	<i>r</i>	1	2	3	4	5	6	7	8	9	10	11
1. Age	42.26	12.54	19-65	-										
2. Sex ¹	-	-	-	-.12*	-									
3. Prior trauma	1.59	1.46	0-10	.09	-.04	-								
4. Prior rob ²	-	-	-	.33**	.05	.11*	-							
5. PDEQ	17.00	7.36	10-45	-.12**	-.23**	.07	-.17**	-						
6. PRS	5.55	7.27	0-43	.05	-.25**	.01	-.11*	.60**	-					
7. TIS	7.67	6.98	0-24	-.11	-.19**	-.06	-.13**	.53**	.48**	-				
8. ASI	10.36	9.36	0-48	-.03	-.08	.02	-.03	.41**	.50**	.36**	-			
9. NCS	32.43	14.27	21-104	-.14	-.21**	.07	-.05	.47**	.49**	.37**	.45**	-		
10. NCW	16.13	7.94	7-46	-.08	-.15**	.04	-.03	.41**	.43**	.29**	.41**	.67**	-	
11. ASDS	34.86	13-53	19-86 ³	-.14	-.27**	.00	-.06	.70**	.63**	.48**	.53**	.59**	.54**	-
12. HTQ	25.23	8.09	17-62	-.09	-.23**	.08	-.01	.47**	.49**	.40**	.43**	.63**	.53**	.68**

Note. *M* = mean, *SD* = standard deviation, *R* = range, PDEQ (Peritraumatic Dissociation Questionnaire), PRS (Physical Reaction Scale), TIS (Tonic Immobility Scale), ASI (Anxiety Sensitivity Index), NCS (negative cognitions about self), NCW (negative cognitions about the world), ASDS (Acute Stress Disorder Scale), HTQ (Harvard Trauma Questionnaire), *N* = 450.

¹0 = females, 1 = males, 60.9 % females, ²0 = no, 1 = yes, 62.9 % previously exposed to robbery. ³ASDS range in victims with ASD = 43-86 (*n* = 53), HTQ range in victims with PTSD = 34-62 (*n* = 27).