



Multiplex determination of analytes related to metabolic syndrome after weight loss with Evidence biochip arrays

Al-Fadda, A. A., Turjoman, A. A., Newell, K., Gibson, D. S., Lamont, J. V., & Duncan, M. W. (2011). Multiplex determination of analytes related to metabolic syndrome after weight loss with Evidence biochip arrays. In *Unknown Host Publication* (Vol. 49, pp. O385). De Gruyter. <http://www.degruyter.com/view/j/cclm.2011.49.issue-s1/cclm.2011.510/cclm.2011.510.xml>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Unknown Host Publication

Publication Status:
Published (in print/issue): 01/01/2011

Document Version
Publisher's PDF, also known as Version of record

General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

0384

TOTAL ANTIOXIDANT CAPACITY, SUPEROXIDE DISMUTASE AND CATALASE IN DIABETIC POLYNEUROPATHY

S. Kundalic¹, B. Kundalic², V. Cosic¹, L. Zvezdanovic-Celebic¹, S. Madic¹, D. Stankovic-Ferlez¹, T. Djordjevic¹, T. Ristic¹, M. Stanjokovic¹

¹Center for Medical Biochemistry, Clinical Center, Nis, Serbia, ²Faculty of Medicine, Nis, Serbia

E-mail: slavacakun@gmail.com

Background. Oxidative stress resulting from enhanced free-radical formation and/or a defect in antioxidant defenses has been implicated in the pathogenesis of experimental diabetic polyneuropathy (DP). The antioxidant capacity is always decreased in diabetic patients, but it seems necessary to measure all the components to ascertain the reasons. The aim of this study was to determine the plasma total antioxidant capacity (TAC) and changes in the activities of superoxide dismutase (SOD) and catalase (CAT) in patients with diabetic polyneuropathy.

Methods. We evaluated TAC, SOD, CAT in samples obtained from 30 DP patients and 30 healthy sex and age matched subjects as control group. Laboratory analyses involved fasting blood glucose and glycated hemoglobin (HbA1c) levels. The activities of SOD and CAT were determined by standard spectrophotometric Methods. Total antioxidant status was measured using Randox kit.

Results. Serum glucose and HbA1c levels were significantly higher in DP patients versus the control group ($p < 0.001$). The TAC was significantly depleted in the diabetic group than healthy donors ($p < 0.001$). Average SOD and CAT activity were significantly lower in patients with DP than control group ($p < 0.01$). There was significant negative correlation between the TAC and serum glucose level, TAC and HbA1c ($p < 0.001$), but not between TAC and duration of diabetes. Significant correlation in the case of SOD and CAT was not marked.

Conclusions. Examination of the data from diabetes and total antioxidant status, SOD and CAT strongly implicates hyperglycemia-induced oxidative stress in DP. We conclude that striving for superior antioxidative therapies remains essential for prevention of neuropathy in diabetic patients.

0385

MULTIPLEX DETERMINATION OF ANALYTES RELATED TO METABOLIC SYNDROME AFTER WEIGHT LOSS WITH EVIDENCE BIOCHIP ARRAYS

Al-Fadda, A.A.¹; Turjoman, A.A.^{1,2}; Newell, K.²; Gibson, D.^{2,3}; Lamont, J.V.⁴; Duncan, M.W.^{1,2}

¹Obesity Research Center, College of Medicine, King Saud University, P.O. Box 2925, Riyadh 11461, Saudi Arabia. ²Division of Endocrinology, Diabetes and Metabolism, Department of Medicine, School of Medicine, University of Colorado Denver, Aurora CO 80045, USA. ³Arthritis Research Group, Queen's University, Centre for Infection and Immunity, Health Sciences Building, Belfast BT9 7BL, UK. ⁴Randox Laboratories Ltd. 55 Diamond Road, Crumlin, Co Antrim BT29 4QY, UK.

E-mail: scientific.publications@randox.com

Background. Variations in the levels of adiponectin, C-peptide, CRP, cystatin C, ferritin, IL-1 α , IL-6, insulin, leptin, PAI-1, resistin, TNF α have been associated to metabolic dysfunctions. Biochip array technology enables their multiplex determination from a single sample, at a single point in time. This generates a patient's profile, which provides more information than single analyte determination. The aim of this study was to determine these analytes in plasma of obese subjects following weight loss.

Methods. Plasma samples from obese subjects ($n=12$, average $IBM > 30 \text{ kg/m}^2$) were analysed. For weight reduction, 5 subjects followed dietary restriction and 7 subjects underwent bariatric surgery. Two different biochip arrays were used for the multiplex determination. Simultaneous chemiluminescent immunoassays applied to the Evidence Investigator analyser were used. Statistical significance between groups was calculated using the paired Student's t-test (two tailed).

Results. Following dietary restriction, significant reductions were observed in plasma insulin and CRP levels ($p=0.045$ and 0.030 respectively). C-peptide, IL-6, leptin and TNF α exhibited moderate but no statistically significant reduction. A marginal increase was observed in PAI-1 and resistin. Adiponectin and IL-1 α remained unchanged. Following bariatric surgery, a highly significant reduction was observed in plasma leptin ($p=0.005$) and a significant increase was observed in plasma adiponectin ($p=0.014$). C-peptide and PAI-1 were reduced and IL-1 α , resistin and TNF α increased but did not reach statistical significance. No significant changes were detected in CRP, IL-6 and insulin levels.

Conclusions. This multi-analytical approach using biochip arrays shows differences between the groups undergoing weight loss and represents a valuable analytical tool in research settings.