

RESEARCH ARTICLE

# Personalised Prescription of Scalable High Intensity Interval Training to Inactive Female Adults of Different Ages

Jacqueline L. Mair<sup>1,3\*</sup>, Alan M. Nevill<sup>2</sup>, Giuseppe De Vito<sup>1</sup>, Colin A. Boreham<sup>1</sup>

**1** UCD Institute for Sport & Health, Newstead, University College Dublin, Belfield, Dublin, Co Dublin, Ireland, **2** School of Sport, Performing Arts and Leisure, University of Wolverhampton, Wolverhampton, West Midlands, United Kingdom, **3** School of Sport, Ulster University, Shore Road, Newtownabbey, Co Antrim, United Kingdom

☞ These authors contributed equally to this work.

\* [j.mair@ulster.ac.uk](mailto:j.mair@ulster.ac.uk)



## Abstract

Stepping is a convenient form of scalable high-intensity interval training (HIIT) that may lead to health benefits. However, the accurate personalised prescription of stepping is hampered by a lack of evidence on optimal stepping cadences and step heights for various populations. This study examined the acute physiological responses to stepping exercise at various heights and cadences in young ( $n = 14$ ) and middle-aged ( $n = 14$ ) females in order to develop an equation that facilitates prescription of stepping at targeted intensities. Participants completed a step test protocol consisting of randomised three-minute bouts at different step cadences (80, 90, 100, 110 steps·min<sup>-1</sup>) and step heights (17, 25, 30, 34 cm). Aerobic demand and heart rate values were measured throughout. Resting metabolic rate was measured in order to develop female specific metabolic equivalents (METs) for stepping. Results revealed significant differences between age groups for METs and heart rate reserve, and within-group differences for METs, heart rate, and metabolic cost, at different step heights and cadences. At a given step height and cadence, middle-aged females were required to work at an intensity on average  $1.9 \pm 0.26$  METs greater than the younger females. A prescriptive equation was developed to assess energy cost in METs using multi-level regression analysis with factors of step height, step cadence and age. Considering recent evidence supporting accumulated bouts of HIIT exercise for health benefits, this equation, which allows HIIT to be personally prescribed to inactive and sedentary women, has potential impact as a public health exercise prescription tool.

## OPEN ACCESS

**Citation:** Mair JL, Nevill AM, De Vito G, Boreham CA (2016) Personalised Prescription of Scalable High Intensity Interval Training to Inactive Female Adults of Different Ages. PLoS ONE 11(2): e0148702. doi:10.1371/journal.pone.0148702

**Editor:** Nir Eynon, Victoria University, AUSTRALIA

**Received:** October 19, 2015

**Accepted:** January 20, 2016

**Published:** February 5, 2016

**Copyright:** © 2016 Mair et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** The authors have no support or funding to report.

**Competing Interests:** The authors have declared that no competing interests exist.

## Introduction

The health and fitness benefits of stair climbing have been established in both epidemiological and intervention studies. Habitual stair climbing has been associated with reduced BMI [1], reduced risk of premature mortality [2–4] and stroke [5] as well as increased longevity [6]. Additionally, short-term intervention trials have found stair climbing to be an effective, time efficient strategy for improvements in aerobic fitness [7–9] and certain health parameters, such

as lipid profiles [8, 10]. For example, Kennedy and colleagues [9] report that 6 minutes of stair climbing per day resulted in comparable improvements in aerobic fitness as walking for 45 minutes per day. It is likely that such health outcomes are a result of the accumulation of short bouts of vigorous intensity stair climbing exercise (8.8 METs; [11]), which is substantiated by investigations linking short duration, high intensity interval training (HIIT) with health benefits [12–14].

Despite the desirable health and fitness outcomes associated with stair climbing, the activity does present a number of barriers and is not suitable for some populations. For example, older individuals have a higher risk of falling while climbing stairs [15], resulting in a higher incidence of such falls [16]. Furthermore, many people may not have access to multiple flights of stairs during their everyday routine and of those who do, several report a stair climbing threshold of no more than four flights of stairs before they will choose to take the lift [17]. Thus, a similar exercise modality, which overcomes the barriers to stair climbing while replicating its benefits, may be of value for public health interventions. Bench stepping is an analogous activity to stair climbing which may fulfil these criteria.

Bench stepping requires an individual to ascend and descend a single step of specified height by completing a four-step cycle (two steps upwards onto the bench, and two steps downwards off the bench). It has been classified as vigorous intensity exercise of approximately 8.5 METs [11], which is similar to stair climbing at a moderate to brisk pace [8, 18, 19]. It is currently unclear, however, what step height and step cadence combination is required to achieve this intensity, especially in populations of different ages. Additionally, the MET intensity allocated to this activity is based on a standardised resting  $\text{VO}_2$  of  $3.5\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  [20] a reference value that is insensitive to differences in age, gender and body mass index (BMI; [21]).

It is important to establish an equivalency between stepping and stair climbing so that the latter (with its proven health benefits [1–10]) may be replicated with bench stepping. This is particularly important if stepping is to be promoted in populations of different ages, as it is well established that physical work capacity reduces with age [22, 23]. Furthermore, previous work from our group has helped to characterise the neuromuscular demands of bench stepping [24], as well as improvements in cardiorespiratory fitness, and mobility arising from short-term bench step training [25, 26]. However, there remains a need to further refine the prescription of step exercise especially considering its potential as a scalable HIIT model that may be promoted on a public health level.

This study, therefore, aims to describe and compare the physiological responses to bench stepping exercise in young and middle-aged females, in order to develop an equation that will allow the prescription of step exercise (using step height and step cadence) at a target intensity known to enhance health and fitness in a given population. The chosen step heights for this study are commonly found in public and residential settings (17 cm; or 34 cm if two are taken at a time) and compare with the heights from a standard, commercially available aerobic bench (25 cm and 30 cm). Our choice of step height therefore facilitates the prescription of stepping exercise based on an individual's everyday environment and promotes activity that may be easily incorporated into the daily routine.

## Methods

The study was approved by the ethics committee of University College Dublin and was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. Fourteen young (aged 19–30 years) and 14 middle-aged (aged 50–63 years) females were recruited for the study through advertisements in local area and university newsletters (Table 1). Participants were provided with a full explanation of the study and written informed

Table 1. Participant characteristics.

	Young	Middle-Aged	t	Cohen's d
Age (years)	23.4 ± 2.74	56.6 ± 3.99		
Stature (m)	1.68 ± 0.66	1.65 ± 0.04	1.78	0.67
Body Mass (kg)	64.8 ± 9.65	64.7 ± 6.57	0.01	0.01
BMI (kg·m <sup>2</sup> )	23.0 ± 3.35	23.9 ± 2.47	-0.87	0.33
Leg Length (cm)	87.4 ± 3.19	86.0 ± 2.71	1.24	0.47
RMR (kcal)	1363.1 ± 288.12	1041.2 ± 153.57	3.64*	1.37
VO <sub>2</sub> rest (mL·kg <sup>-1</sup> ·min <sup>-1</sup> )	3.08 ± 0.59	2.36 ± 0.36	3.90*	1.47

\*significant difference between-groups (p = 0.001). RMR = resting metabolic rate. BMI = body mass index. Values are means ± SD, n = 14 in each group, df = 26.

doi:10.1371/journal.pone.0148702.t001

consent was obtained prior to testing. Inclusion criteria were: females aged 18–30 years and 50–64 years who are physically inactive (as defined by a failure to meet recommended activity levels [27]), but medically stable [28]. Exclusion criteria included cigarette smoking, anabolic steroid ingestion, a history of cardiovascular disease, diabetes (type I or type II), hypertension, or any other metabolic disease or illness requiring the ingestion of medications that affect carbohydrate or lipid metabolism.

## Research Design

The experimental design required participants to report to the exercise physiology laboratory on three occasions, separated by a minimum of seven days. During the first session participants were measured for anthropometric variables and resting metabolic rate (RMR), and on the second and third sessions, participants completed a step test protocol (described below).

## Anthropometric variables

Stature was determined with a stadiometer (Holtain Ltd, Pembs, UK) to the nearest 0.1cm, and body mass was measured on a digital scale (Seca, Hamburg, Germany) to the nearest 0.1kg with the participant wearing minimal clothing and no shoes. BMI was calculated using the equation  $body\ mass/height^2$ . Leg length was measured to the nearest 0.1cm with the subject in a supine position taking a line from the anterior superior iliac spine to the medial malleolus using a standard tape measure.

## Resting Metabolic Rate

RMR measurements were carried out following a recommended protocol [29]. Participants were instructed to abstain from food and beverages (except water) for a minimum of 12 hours and to avoid strenuous exercise for 24 hours prior to testing. In addition, all participants travelled by car or public transport on the day of testing so as not to raise metabolism significantly above resting levels. RMR was measured using a Quark b<sup>2</sup> indirect calorimeter (Cosmed Srl, Rome, Italy) following standard manufacturer calibration procedures, and resting HR was measured using a telemetric heart rate monitor (Polar Electro Oy, Kempele, Finland). On arrival at the laboratory, participants rested quietly in a supine position for 20 minutes in a temperature-controlled room. Measurement commenced immediately following the rest period, using a facemask protocol, and respiratory quotient values were monitored throughout the test to ensure accurate gas measurements. Data were collected at 30-second intervals for a period of 15 minutes and analysed offline for steady state.

## Dietary Control Measures

All participants completed a two-day nutritional diary on the days leading up to the second testing session. This was then replicated prior to the third testing session to reduce variability in metabolic measures that may have resulted from a change in diet. Participants were instructed to abstain from food and beverages three hours before testing, not to drink coffee or beverages containing caffeine for at least eight hours and to avoid intense exercise for 48 hours prior to testing.


## Stepping Exercise Test

The stepping test protocol (Fig 1) consisted of randomised three-minute stepping bouts at four different cadences (80, 90, 100, 110 steps·min<sup>-1</sup>) interspersed by three-minutes of rest to allow heart rate to return to within 10 beats of baseline values. This protocol was repeated using four different step heights (17, 25, 30, 34 cm), two completed during session two and two completed during session three, also selected in a random order. Participants therefore completed a total of 16 three-minute stepping bouts. The stepping cycle consisted of two steps up onto the bench to full extension of the legs and two steps back down, maintaining the same leading leg throughout. Therefore a cadence of 100 steps·min<sup>-1</sup> equated to 25 step cycles. Step cadence was regulated using a pre-selected music track with the required tempo. All participants performed a familiarisation trial at each treatment combination of step height and step cadence prior to the commencement of testing on both days. A handrail was provided as an added safety precaution in the event that the participant should lose balance during the test, but care was taken to ensure that the handrail was not used to aid the stepping cycle.

Throughout the stepping protocol, the participant wore a face-mask (Hans Rudolf Inc, Shawnee, KS, USA) connected to an indirect calorimeter (Cosmed Srl, Rome, Italy) for breath-by-breath measurements of gas exchange, and a telemetric heart rate monitor (Polar Electro Oy, Kempele, Finland) for continuous measurement of heart rate (HR). The gas analyser was calibrated with known concentrations of O<sub>2</sub> and CO<sub>2</sub> and the turbine was calibrated with a three-litre syringe prior to every testing session. Baseline resting values of VO<sub>2</sub> and HR were collected for two minutes immediately preceding commencement of the stepping protocol.

## Data Analysis

Resting metabolic rate was calculated from 10 minutes of steady state data (defined as variation in the volume of oxygen consumed of no more than 10% [30]) and in all cases, the initial five minutes of test data were discarded. Heart rate reserve (HRr) was calculated [31] using maximum HR determined from a female specific equation [32], and mean resting HR obtained during the RMR test. Mean values for VO<sub>2</sub> (ml·kg<sup>-1</sup>·min<sup>-1</sup>), HR (beats·min<sup>-1</sup>), HRr (%) and metabolic cost (Kcal·min<sup>-1</sup>) were calculated from the final 30 s of filtered data at each step height/step cadence combination during the stepping test, and METs were calculated by dividing mean VO<sub>2</sub> by the corresponding individual resting VO<sub>2</sub> value obtained during the RMR test.



**Fig 1. Stepping test protocol.**

doi:10.1371/journal.pone.0148702.g001

## Statistical Analysis

Statistical procedures were carried out using SPSS 20 (IBM Corporation, New York, USA). Sample size was established by performing an *a priori* power calculation based on previously published data [33]. Using a two-tail multiple regression model with a significance level of 5%, an effect size of 0.82 (derived from an  $R^2$  value of 0.45), and 3 predictor variables, this analysis suggested a minimum of 14 participants per group would achieve a power of 85%.

Normal distributions were inspected using the Shapiro-Wilk test and normal Q-Q plots prior to analysis. An independent samples t-test was used to test between-group differences for anthropometric variables and RMR values and effect size values were calculated using Cohen's  $d$ , whereby 0.2 is considered a small effect, 0.5 a medium effect, and 0.8 a large effect [34]. A three-way ANOVA with repeated measures was used to analyse dependant variables (METs, HR, HRr, metabolic cost), with the three factors being a between-subjects factor of 'age group' (the young vs. middle-aged) and two within-subjects factors of 'step height' and 'step cadence'. *Post-hoc* pairwise comparisons with Bonferroni correction were used to identify where any detected differences lay and effect sizes were calculated using partial eta squared ( $\eta_p^2$ ) where 0.01 is a small effect, 0.06 is a medium effect, and 0.14 is a large effect [34]. A multilevel regression analysis [35] on the combined data (young and middle-aged groups) was performed using Multilevel Models Project (MLn) software in order to establish a prediction equation for the purpose of prescribing stepping exercise at a target intensity (METs). The two-level hierarchical regression model used to predict METs incorporated three fixed-factor predictor variables; step height, step cadence and age, with participants or individuals, assumed to be a random sample, being the level two units, and with the subjects' repeated measurements, being the level one units. Standard errors obtained from maximum-likelihood estimation procedures were used to establish the significance of the model. Statistical significance was accepted as  $P < 0.05$ .

## Results

There were no differences between groups for any measured anthropometric variables. RMR was significantly lower in the middle-aged females compared with the young (Table 1). Furthermore, resting  $\text{VO}_2$  was 12% and 33% lower than the standard resting value (equivalent to 1 MET) of  $3.5 \text{ ml} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  in the young and middle-aged group, respectively.


### Between-Group Analysis

The repeated measures ANOVA revealed significant between-group main effects across step height and step cadence combinations for METs ( $P < 0.01$ ,  $\eta_p^2 = .350$ ; Fig 2) and %HRr ( $P < 0.01$ ,  $\eta_p^2 = .366$ ), but there were no observable differences for HR ( $P = 0.84$ ,  $\eta_p^2 = .002$ ), or metabolic cost ( $P = 0.81$ ,  $\eta_p^2 = .002$ ). At a given step height and cadence, the middle-aged females were required to work at an intensity on average  $1.9 \pm 0.26$  METs greater than the younger females.

### Within-Group Analysis

Within-group comparisons (Table 2) revealed a significant main effect for step height across all dependent variables; METs ( $P < 0.001$ ,  $\eta_p^2 = .904$ ), HR ( $P < 0.001$ ,  $\eta_p^2 = .864$ ), %HRr ( $P < 0.001$ ,  $\eta_p^2 = .864$ ) and metabolic cost ( $P < 0.001$ ,  $\eta_p^2 = .932$ ). Similarly, a significant main effect for step cadence was observed for all dependent variables; METs ( $P < 0.001$ ,  $\eta_p^2 = .893$ ), HR ( $P < 0.001$ ,  $\eta_p^2 = .940$ ), %HRr ( $P < 0.001$ ,  $\eta_p^2 = .942$ ) and metabolic cost ( $P < 0.001$ ,  $\eta_p^2 = .924$ ).

There was a significant third order interaction effect (step height\*step cadence\*age group) for METs ( $P = 0.02$ ,  $\eta_p^2 = .094$ ) indicating a slight non-linear response. As evident in Fig 2, for



**Fig 2. METs across step height and step cadence combinations.** Values are means  $\pm$  SE; Y = young females; M = middle-aged females; SC = step cadence.

doi:10.1371/journal.pone.0148702.g002

the middle-aged group linear trends begin to converge at higher step heights and rates, showing a slight plateau in the metabolic response. Significant interaction effects for HR (step height\* - step cadence:  $P = 0.046$ ,  $\eta_p^2 = .080$ ) and %HRr (step height\*step cadence:  $P = 0.043$ ,  $\eta_p^2 = .081$ ; step cadence\*group:  $P < 0.001$ ,  $\eta_p^2 = .351$ ) and metabolic cost (step height\*group:  $P = 0.036$ ,  $\eta_p^2 = .103$ ) were also detected.

### Multilevel Regression Analysis

A model for estimating the MET intensity of stepping exercise was developed from step height, step cadence and age (Eq 1). The percentage of additional variance accounted for by the

**Table 2. Physiological responses to differing step heights and step cadences, in young and middle-aged females.**

Step Height (cm)	Middle-Aged				Young			
	Step Cadence (steps·min <sup>-1</sup> )							
	80	90	100	110	80	90	100	110
<i>Heart Rate (beats·min<sup>-1</sup>)*</i>								
17	116.4 $\pm$ 14.74	121.7 $\pm$ 15.75	126.8 $\pm$ 15.73	134.9 $\pm$ 14.89	115.2 $\pm$ 20.72	121.6 $\pm$ 23.13	124.3 $\pm$ 22.03	130.5 $\pm$ 23.83
25	126.8 $\pm$ 12.26	135.0 $\pm$ 13.37	141.9 $\pm$ 15.28	149.4 $\pm$ 15.51	128.5 $\pm$ 22.70	134.0 $\pm$ 22.86	137.6 $\pm$ 22.12	146.5 $\pm$ 21.09
30	138.0 $\pm$ 16.38	147.9 $\pm$ 15.95	152.4 $\pm$ 15.69	161.8 $\pm$ 14.70	136.4 $\pm$ 22.89	146.5 $\pm$ 25.04	155.1 $\pm$ 24.73	159.9 $\pm$ 23.04
34	147.6 $\pm$ 13.92	154.6 $\pm$ 14.34	160.1 $\pm$ 10.60	166.8 $\pm$ 9.21	148.5 $\pm$ 21.32	151.1 $\pm$ 20.87	159.3 $\pm$ 22.31	166.7 $\pm$ 20.45
<i>Heart rate reserve (%)* †</i>								
17	59.7 $\pm$ 14.04	65.1 $\pm$ 14.63	69.9 $\pm$ 15.53	78.4 $\pm$ 14.12	45.0 $\pm$ 14.32	50.2 $\pm$ 16.53	52.2 $\pm$ 15.67	57.2 $\pm$ 17.34
25	70.1 $\pm$ 11.58	78.0 $\pm$ 12.92	85.5 $\pm$ 13.83	93.0 $\pm$ 14.19	54.9 $\pm$ 16.73	59.9 $\pm$ 16.82	62.9 $\pm$ 15.96	69.9 $\pm$ 15.44
30	80.5 $\pm$ 15.40	90.9 $\pm$ 15.56	96.05 $\pm$ 14.85	105.7 $\pm$ 13.52	62.2 $\pm$ 16.65	70.3 $\pm$ 18.20	74.8 $\pm$ 18.07	80.7 $\pm$ 17.23
34	91.2 $\pm$ 13.76	98.5 $\pm$ 14.75	104.0 $\pm$ 10.86	110.6 $\pm$ 8.80	72.1 $\pm$ 16.04	73.6 $\pm$ 14.75	79.9 $\pm$ 16.61	86.0 $\pm$ 15.61
<i>Metabolic Cost (Kcal·min<sup>-1</sup>)*</i>								
17	5.25 $\pm$ 0.81	5.66 $\pm$ 0.95	6.15 $\pm$ 1.01	6.76 $\pm$ 1.13	5.13 $\pm$ 0.86	5.70 $\pm$ 0.87	6.08 $\pm$ 0.96	6.63 $\pm$ 1.03
25	6.14 $\pm$ 0.90	6.87 $\pm$ 0.94	7.43 $\pm$ 0.90	8.01 $\pm$ 0.94	6.21 $\pm$ 1.06	6.81 $\pm$ 0.97	7.23 $\pm$ 1.04	7.93 $\pm$ 1.12
30	7.32 $\pm$ 1.12	7.88 $\pm$ 1.18	8.20 $\pm$ 1.09	8.70 $\pm$ 1.25	7.08 $\pm$ 1.02	7.90 $\pm$ 1.16	8.43 $\pm$ 1.12	9.00 $\pm$ 1.28
34	7.95 $\pm$ 1.23	8.36 $\pm$ 0.98	8.93 $\pm$ 0.93	9.43 $\pm$ 1.14	8.01 $\pm$ 1.03	8.75 $\pm$ 1.25	9.35 $\pm$ 1.3	10.16 $\pm$ 1.44

\*Within-group differences are significant ( $p < 0.01$ ).

†Between-group differences are significant ( $p > 0.01$ ).

Values are means  $\pm$  SD.

doi:10.1371/journal.pone.0148702.t002

inclusion of leg length ( $-0.066 \pm 0.086$ ) was not significant. Similarly, the addition of stature ( $-0.087 \pm 0.045$ ) to the model did not account for additional variance in METs and did not significantly alter the percentage variance explained by the initial three predictors. The estimated parameters from the multilevel regression analysis including the three fixed factors and the two sources of random variation between- and within-individuals are given in [Table 3](#).

$$\text{METs} = (a + b \cdot \text{step height}) + (c \cdot \text{step cadence}) + (d \cdot \text{age}) \tag{Eq 1}$$

### Discussion

The aim of the present study was to describe the physiological responses to bench stepping exercise in young and middle-aged females in order to develop an equation that will allow the prescription of step exercise (using step height and step cadence) at a target intensity known to enhance health and fitness in a given population. It was found that METs, HR and metabolic cost vary considerably in response to differing step heights and cadences, which supports previous research in the area of aerobic dance bench stepping [33, 36, 37]. Our results show that both step height and step cadence have a significant independent effect on the physiological responses to stepping exercise. In both the young and middle-aged groups, each 10 beat.min<sup>-1</sup> increase in step cadence resulted in a 0.6-MET increase in intensity, while each mean increase of 6 cm in step height resulted in a 1-MET increase in intensity. We further extend this line of research by reporting that the acute responses to stepping at different heights and cadences vary across different age groups. Results revealed that at a given step height and cadence, the older females were required to work at an intensity on average  $1.9 \pm 0.26$  METs greater than the younger females. For example, a step height and cadence combination that corresponded to an intensity of 9 METs (75%HRr) in the younger group resulted in 10.8 METs (96% HRr) in the middle-aged group, an intensity that is close to maximal exertion. Since ageing is accompanied by a decline in the efficiency of the cardiovascular system [38] and in skeletal muscle quantity [39], these results are not entirely unexpected. At a step height of 30 cm, the middle-aged steppers are already operating close to their maximum capability suggesting that further increases in step height are unnecessary and impractical for this age group. Furthermore, these age-related differences provide additional evidence that previously published equations for step exercise [40] may significantly underestimate the metabolic cost of this activity in older populations, resulting in inappropriate exercise prescriptions. Our results, therefore, lend credence to the importance of tailored exercise programmes for individuals of different ages.

**Table 3. Multilevel regression analysis of METs, using step height, step cadence, and age as predictor variables.**

Fixed Explanatory Variables	Value
Constant (a)	-3.743 ± 0.695
Step Height (b)	0.188 ± 0.004
Step Cadence (c)	0.060 ± 0.002
Age (d)	0.053 ± 0.015
Variance-Covariance Matrix of Random Variables	
	Constant (a)
	<i>Level 1 (within individuals)</i>
Constant (a)	0.332 ± 0.023
	<i>Level 2 (between individuals)</i>
Constant (a)	1.731 ± 0.468

Values are means ± SE

doi:10.1371/journal.pone.0148702.t003

Stepping exercise has the potential to be promoted on a public health level as a scalable HIIT model, as it is low-cost, simple to perform, and may be particularly appealing to those who have restricted access to exercise facilities or little time to exercise. With this, as well as the need for personalised prescriptions in mind, a further aim of our study was to develop a useful means of prescribing intensity-specific stepping exercise to female adults of different ages. By choosing a step height that is readily accessible to the individual, it is possible to calculate the step cadence required to reach a desired age-adjusted vigorous intensity level (Eq 1). As previously mentioned, stair climbing exercise is associated with a number of short and long-term positive health outcomes [1–10], and so replicating the physiological responses of this everyday activity through bench stepping exercise should result in similar benefits. It is therefore recommended that users of the prescription equation select an intensity of 8.8 METs (equivalent to stair climbing) in order to prescribe an appropriate stepping cadence based on their age and the step height available to them. Furthermore, to reflect a scaled HIIT protocol, stepping bouts of 3 minutes, interspersed with up to 3 minutes of recovery, are suggested. Personalised exercise prescription such as this may improve adherence to exercise programmes in inactive populations [41]. By addressing a number of common barriers to exercise participation using stepping as a vehicle for scalable HIIT, there is a greater likelihood of longer term adherence to the exercise. Furthermore, high intensity exercise has been linked with heightened mood post exercise [42] and higher levels of enjoyment compared with moderate intensity exercise [43, 44]. For example, Bartlett and co-workers [43] have shown higher ratings of perceived enjoyment following 3 minute bouts of HIIT compared with 50 minutes of continuous exercise. Further examination of training responses, as well as comparison with traditional exercise programmes and more contemporary HIIT protocols, is required in order to elucidate the long term benefits of this scalable exercise prescription. Those wishing to use this equation to prescribe exercise training to individuals should consider the principle of overload to ensure continued cardiorespiratory fitness adaptations as training progresses.

By including an accurate measurement of RMR into the calculation of MET intensity, the quality of the prediction for the individual work performed by participants has been improved. This measurement also allowed comparison of resting metabolism between the age groups, and with the standard resting  $\text{VO}_2$  value of  $3.5\text{ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$  [20], which is still commonly cited in research. Our results show that young and older females have a 12% and 33% lower RMR than the standard value, respectively, further highlighting the inadequacy of this standardisation in certain populations [21, 45]. It should be noted that the RMR measurement used in this study did not account for differences in lean body mass between groups, which may have further refined our results. Byrne and colleagues [21] have shown BMI to be strongly correlated with fat mass in a heterogeneous sample. Therefore, it is possible that the slightly higher average BMI in our middle-aged group could have reflected a higher fat mass and contributed to the lower RMR values compared with our younger group. Importantly, this study introduces a more accurate representation of the relative MET intensity of stepping exercise across two different age groups.

## Limitations

The study sample was restricted to healthy Caucasian females, within two age ranges, and therefore the reported results are not truly representative of the general population. Lean muscle mass was not measured in the present study, data from which may have helped to explain age-related difference in RMR. Further, due to impracticalities it was not possible to complete all testing on the same day. While every effort was made to present participants for testing in the same physiological state, day-to-day variability in may have introduced some error to calculations.



## Conclusions

The ability to prescribe stepping exercise at a specific intensity that replicates stair climbing means that the health benefits of the latter can potentially be promoted to a larger population. This study confirms that stepping can replicate the physiological demands of stair climbing in both a young and middle-aged female population, however younger females require higher step heights and faster cadences compared with their middle-aged counterparts, to reach higher levels of exercise intensity. The importance of innovative exercise methods to improve health and fitness in a practical, time-efficient manner across various populations is growing. This study provides a simple means of prescribing an intensity-specific step exercise programme, which may be performed in small amounts throughout the day, as a scalable form of HIIT, to females of different ages.

## Supporting Information

**S1 Dataset. Study data.**  
(SAV)

## Author Contributions

Conceived and designed the experiments: CAB GDV. Performed the experiments: JLM. Analyzed the data: JLM AMN. Contributed reagents/materials/analysis tools: CAB GDV AMN. Wrote the paper: JLM CAB GDV AMN.

## References

1. Shenassa ED, Frye M, Braubach M, Daskalakis C. Routine stair climbing in place of residence and body mass index: a pan-European population based study. *Int J Obesity (Lond)*. 2008; 32: 490–494.
2. Yu S, Yarnell JW, Sweetnam PM, Murray L. What level of physical activity protects against premature cardiovascular death? The Caerphilly Study. *Heart*. 2003; 89: 502–506. PMID: [12695452](#)
3. Paffenbarger RS, Hyde RT, Wing AL, Lee IM, Jung DL, Kampert JB. The association of changes in physical-activity level and other lifestyle characteristics with mortality among men. *New Engl J Med*. 1993; 328(8): 538–545. PMID: [8426621](#)
4. Myers J, Kaykha A, George S, Albella J, Zaheer N, Lear S, Yamazaki T, Froelicher V. Fitness versus physical activity patterns in predicting mortality in men. *Am J Med*. 2004; 117: 912–918. PMID: [15629729](#)
5. Lee IM, Paffenbarger RS. Physical activity and stroke incidence: the Harvard Alumni Health Study. *Stroke*. 1998; 29: 2049–2054. PMID: [9756580](#)
6. Lee IM, Paffenbarger RS Jr. Associations of light, moderate, and vigorous intensity physical activity with longevity. The Harvard Alumni Health Study. *Am J Epidemiol*. 2000; 151: 293–299. PMID: [10670554](#)
7. Fardy PS, Ilmarinen J. Evaluating the effects and feasibility of an at work stairclimbing intervention program for men. *Med Sci Sports*. 1975; 7: 91–93. PMID: [1152630](#)
8. Boreham CA, Wallace WF, Nevill A. Training effects of accumulated daily stair-climbing exercise in previously sedentary young women. *Prev Med*. 2000; 30: 277–281. PMID: [10731455](#)
9. Kennedy RA, Boreham CA, Murphy MH, Young I, Mutrie N. Evaluating the effects of a low volume stair-climbing programme on measures of health-related fitness in sedentary office workers. *J Sports Sci Med*. 2007; 6: 448–454. PMID: [24149477](#)
10. Engbers LH, van Poppel MN, van Mechelen W. Modest effects of a controlled worksite environmental intervention on cardiovascular risk in office workers. *Prev Med*. 2007; 44: 356–362. PMID: [17187852](#)
11. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR, Tudor-Locke C, et al. 2011 Compendium of Physical Activities: A Second Update of Codes and MET Values. *Med Sci Sports Exerc*. 2011; 43: 1575–1581. doi: [10.1249/MSS.0b013e31821ece12](#) PMID: [21681120](#)
12. Babraj JA, Vollaard NB, Keast C, Guppy FM, Cottrell G, Timmons JA. Extremely short duration high intensity interval training substantially improves insulin action in young healthy males. *BMC Endocr Disord*. 2009; 9(1): 3–10.

13. Hood MS, Little JP, Tarnopolsky MA, Myslik F, Gibala MJ. Low-volume interval training improves muscle oxidative capacity in sedentary adults. *Med Sci Sports Exerc.* 2011; 43: 1849–1856. doi: [10.1249/MSS.0b013e3182199834](https://doi.org/10.1249/MSS.0b013e3182199834) PMID: [21448086](https://pubmed.ncbi.nlm.nih.gov/21448086/)
14. Higgins TP, Baker MD, Evans SA, Adams RA, Cobbold C. Heterogeneous responses of personalised high intensity interval training on type 2 diabetes mellitus and cardiovascular disease risk in young healthy adults. *Clin Hemorheol Micro.* 2014; 59(4): 365–377.
15. Studenski S, Duncan PW, Chandler J, Samsa G, Prescott B, Hogue C, et al. Predicting falls: the role of mobility and nonphysical factors. *J Am Geriatr Soc.* 1994; 42: 297–302. PMID: [8120315](https://pubmed.ncbi.nlm.nih.gov/8120315/)
16. Hemenway D, Solnick SJ, Koeck C, Kytir J. The incidence of stairway injuries in Austria. *Accid Anal Prev.* 1994; 26(5): 675–679. PMID: [7999212](https://pubmed.ncbi.nlm.nih.gov/7999212/)
17. Adams J, White M. A systematic approach to the development and evaluation of an intervention promoting stair use. *Health Education Journal.* 2002; 61(3): 272–286.
18. Bassett DR, Vachon JA, Kirkland AO, Howley ET, Duncan GE, Johnson KR. Energy cost of stair climbing and descending on the college alumnus questionnaire. *Med Sci Sports Exerc.* 1997; 29: 1250–1254. PMID: [9309638](https://pubmed.ncbi.nlm.nih.gov/9309638/)
19. Teh KC, Aziz AR. Heart rate, oxygen uptake, and energy cost of ascending and descending the stairs. *Med Sci Sports Exerc.* 2002; 34: 695–699. PMID: [11932581](https://pubmed.ncbi.nlm.nih.gov/11932581/)
20. Balke B. Biodynamics: Human. In: Glasser O, editor. *Medical physics.* Chicago: Year Book Medical Publishers Inc; 1960. pp. 50–52.
21. Byrne NM, Hills AP, Hunter GR, Weinsier RL, Schutz Y. Metabolic equivalent: one size does not fit all. *J Appl Physiol.* 2005; 99: 1112–1119. PMID: [15831804](https://pubmed.ncbi.nlm.nih.gov/15831804/)
22. Bouchard C, Shephard RJ, Stephens T, Sutton JR. *Exercise, fitness and health: a consensus of current knowledge.* Champaign: Human Kinetics; 1990.
23. Betik AC, Hepple RT. Determinants of VO<sub>2</sub> max decline with aging: an integrated perspective. *Appl Physiol Nutr Metab.* 2008; 33: 130–140. doi: [10.1139/H07-174](https://doi.org/10.1139/H07-174) PMID: [18347663](https://pubmed.ncbi.nlm.nih.gov/18347663/)
24. Mair JL, Laudani L, Vannozzi G, De Vito G, Boreham C, Macaluso A. Neuromechanics of repeated stepping with external loading in young and older women. *Eur J Appl Physiol.* 2014; 114(5): 983–994. doi: [10.1007/s00421-014-2826-9](https://doi.org/10.1007/s00421-014-2826-9) PMID: [24504651](https://pubmed.ncbi.nlm.nih.gov/24504651/)
25. Doheny EP, McGrath D, Ditroilo M, Mair JL, Greene BR, Caulfield B, et al. Effects of a low-volume, vigorous intensity step exercise program on functional mobility in middle-aged adults. *Ann Biomed Eng.* 2013; 241(8): 1748–1757.
26. Mair JL, Boreham C, Ditroilo M, McKeown D, Lowery MM, Caulfield B, et al. Benefits of a worksite or home-based bench stepping intervention for sedentary middle-aged adults—a pilot study. *Clin Physiol Funct Imaging.* 2013; 32(1): 10–17.
27. Department of Health. *Start active, stay active: a report on physical activity from the four home countries. A report from the Chief Medical Officer.* London: Department of Health; 2011.
28. Greig CA, Young A, Skelton DA, Pippet E, Butler FM, Mahmud SM. Exercise studies with elderly volunteers. *Age Ageing.* 1994; 23: 185–189. PMID: [8085501](https://pubmed.ncbi.nlm.nih.gov/8085501/)
29. Compher C, Frankenfield D, Keim N, Roth-Yousey L. Best practice methods to apply to measurement of resting metabolic rate in adults: a systematic review. *J Am Diet Assoc.* 2006; 106: 881–903. PMID: [16720129](https://pubmed.ncbi.nlm.nih.gov/16720129/)
30. Horner NK, Lampe JW, Patterson RE, Neuhaus ML, Beresford SA, Prentice RL. Indirect calorimetry protocol development for measuring resting metabolic rate as a component of total energy expenditure in free-living postmenopausal women. *J Nutr.* 2001; 131: 2215–2218. PMID: [11481420](https://pubmed.ncbi.nlm.nih.gov/11481420/)
31. Karvonen MJ, Kentala E, Mustala O. The effects of training on heart rate; a longitudinal study. *Ann Med Exp Biol Fenn.* 1957; 35: 307–315. PMID: [13470504](https://pubmed.ncbi.nlm.nih.gov/13470504/)
32. Gulati M, Shaw LJ, Thisted RA, Black HR, Bairey Merz CN, et al. Heart rate response to exercise stress testing in asymptomatic women: the St. James women take heart project. *Circulation.* 2010; 122: 130–137. doi: [10.1161/CIRCULATIONAHA.110.939249](https://doi.org/10.1161/CIRCULATIONAHA.110.939249) PMID: [20585008](https://pubmed.ncbi.nlm.nih.gov/20585008/)
33. Grier TD, Lloyd LK, Walker JL, Murray TD. Metabolic cost of aerobic dance bench stepping at varying cadences and bench heights. *J Strength Cond Res.* 2002; 16: 242–249. PMID: [11991777](https://pubmed.ncbi.nlm.nih.gov/11991777/)
34. Cohen J. *Statistical power analysis for the behavioral sciences.* Hillsdale: Lawrence Erlbaum Associates; 1988.
35. Nevill AM, Holder RL, Baxter-Jones A, Round JM, Jones DA. Modelling developmental changes in strength and aerobic power in children. *J Appl Physiol.* 1998; 84: 963–970. PMID: [9480958](https://pubmed.ncbi.nlm.nih.gov/9480958/)
36. Scharff-Olson MR, Williford HN. The energy cost associated with selected step training exercise techniques. *Res Q Exerc Sport.* 1996; 67: 465–468. PMID: [9016489](https://pubmed.ncbi.nlm.nih.gov/9016489/)

37. Zaletel P, Furjan-Mandic G, Zagorc M. Differences in heart rate and lactate levels at three different workloads in step aerobics. *Kinesiology*. 2009; 41: 97–104.
38. Lakatta EG. Changes in cardiovascular function with aging. *Eur Heart J*. 1990; 11: 22–29. PMID: [2188839](#)
39. Janssen I, Heymsfield SB, Wang ZM, Ross R. Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J Appl Physiol*. 2000; 89:81–88. PMID: [10904038](#)
40. American College of Sports Medicine. *ACSM's guidelines for exercise testing and prescription*. 7th ed. Baltimore: Lippincott Williams & Wilkins; 2006.
41. Buford TW, Roberts MD, Church TS. Toward exercise as personalized medicine. *Sports Med*. 2013; 43(3): 157–165. doi: [10.1007/s40279-013-0018-0](#) PMID: [23382011](#)
42. Hall EE, Ekkekakis P, Petruzzello SJ. The affective beneficence of vigorous exercise revisited. *Brit J Health Psych*. 2002; 7: 47–66.
43. Bartlett JD, Close GL, MacLaren DPM, Gregson W, Drust B, Morton JP. High-intensity interval running is perceived to be more enjoyable than moderate-intensity continuous exercise: Implications for exercise adherence. *J Sports Sci*. 2011; 29(6): 547–553. doi: [10.1080/02640414.2010.545427](#) PMID: [21360405](#)
44. Jung ME, Bourne JE, Little JP. Where Does HIT Fit? An Examination of the Affective Response to High-Intensity Intervals in Comparison to Continuous Moderate- and Continuous Vigorous-Intensity Exercise in the Exercise Intensity-Affect Continuum. *PLoS ONE*. 2014; 9(12): e114541. doi: [10.1371/journal.pone.0114541](#) PMID: [25486273](#)
45. Sergi G, Coin A, Sarti S, Perissinotto E, Peloso M, Mulone S, et al. Resting VO<sub>2</sub>, maximal VO<sub>2</sub> and metabolic equivalents in free-living healthy elderly women. *Clin Nutr*. 2010; 29: 84–88. doi: [10.1016/j.clnu.2009.07.010](#) PMID: [19713012](#)