



## Clinical pharmacy interventions in an Austrian hospital: a report highlights the need for the implementation of clinical pharmacy services

Guntschnig, S., Courtenay, A., Abuelhana, A., & Scott, M. (2023). Clinical pharmacy interventions in an Austrian hospital: a report highlights the need for the implementation of clinical pharmacy services. *European Journal of Hospital Pharmacy*, Article ejhpharm-2023-003840. <https://doi.org/10.1136/ejhpharm-2023-003840>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
European Journal of Hospital Pharmacy

**Publication Status:**  
Published (in print/issue): 25/09/2023

**DOI:**  
[10.1136/ejhpharm-2023-003840](https://doi.org/10.1136/ejhpharm-2023-003840)

**Document Version**  
Author Accepted version

**Document Licence:**  
CC BY-NC

### General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

### Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk)

# Clinical Pharmacy Interventions in an Austrian hospital: A report highlights the need for the implementation of clinical pharmacy services

Sonja Guntschnig, corresponding author: [sonja.guntschnig@gmail.com](mailto:sonja.guntschnig@gmail.com);  
Clinical Pharmacist at Tauernklinikum Zell am See, Austria; PhD Student at  
Ulster University, NI; ORCID: 0000-0003-2128-5693

Aaron Courtenay Clinical pharmacy Lecturer at Ulster University, NI;  
ORCID: 0000-0002-2651-7839

Ahmed Abuelhana, Clinical pharmacy Lecturer at Ulster University, NI;  
ORCID: 0000-0003-0823-4736

Michael Scott, Director of regional medicines optimisation innovation  
centre, Antrim, NI; Visiting Professor at Ulster University, NI; ORCID:  
0000-0003-4048-0937

### 3 Abstract

4 **Title:** Clinical Pharmacy Interventions in an Austrian hospital: A report highlights the need for  
5 the implementation of clinical pharmacy services

6 **Background:** Clinical pharmacy services face challenges in Austria due to limited implementation  
7 and acceptance, outdated legislation, and a lack of guidelines and training, despite the evidence  
8 from global studies of the positive impact of clinical pharmacists on patient care.

9 **Objectives:** This study aims to identify the necessary types of clinical pharmacy interventions  
10 required at a 360-bed hospital located in Austria. The second aim is to evaluate the extent to  
11 which physicians accept the suggestions made by clinical pharmacists.

12 **Methods:** Over a period of 27 months, a clinical pharmacist made a series of interventions,  
13 which were evaluated using a six-point clinical significance scale. To determine the inter-rater  
14 reliability, a subset of 25 interventions was assessed for their clinical significance by four  
15 independent internal medicine physicians.

16 **Results:** A total of 1064 interventions were made by the pharmacist. Clinical pharmacy input  
17 was deemed necessary for 72.3% (986 out of 1364) of patients, with an average of 1.08  
18 interventions per patient. The prompt acceptance rate of these interventions by physicians was  
19 83.5% (888 out of 1064), while 12.9% (137 out of 1064) were considered by physicians but not  
20 immediately acted upon. The average clinical significance intervention rating was 2.15. The inter-  
21 rater reliability agreement between the four MDs and between the four MDs and the pharmacist  
22 was classified as 'good' to 'moderate'.

23 **Conclusion:** This study in a secondary care Austrian hospital demonstrates the requirement for  
24 clinical pharmacy services, which are highly valued by other healthcare professionals. The clinical  
25 pharmacist is a key member of the multidisciplinary ward team, playing a vital role in reducing  
26 drug related problems and enhancing patient safety. This work should now be scaled and tested  
27 in other Austrian hospitals.

## 28 Key messages

29

### 30 What is already known on this topic?

31 A wealth of research underscores the global benefits of clinical pharmacy services for patient  
32 safety and well-being. However, the acceptance of these interventions varies significantly,  
33 contingent upon diverse factors.

### 34 What this study adds?

35 This study addresses a notable gap by highlighting the absence of Austrian publications on this  
36 subject. It emphasizes the imperative of introducing clinical pharmacy services in a small rural  
37 Austrian hospital and gauges their acceptance among medical staff.

### 38 How might this study affect research, practice or policy?

39 This study has the potential to inspire Austrian clinical pharmacy researchers to share their  
40 findings, fostering a culture of knowledge dissemination. Moreover, it could stimulate increased  
41 scientific engagement, staffing, and policy adjustments within hospitals, ultimately impacting  
42 healthcare quality and policy decisions.

## 43 Keywords

- 44 • Clinical competence
- 45 • Drug-related side effects and adverse reactions
- 46 • Evidence-based medicine
- 47 • Quality of health care
- 48 • Patient safety
- 49 • Medical errors
- 50 • Public health
- 51 • Pharmacy service, hospital

## 52 Introduction

53 According to the European Society of Clinical Pharmacy (ESCP), “Clinical pharmacy aims to  
54 optimise the utilisation of medicines through practice and research in order to achieve person-  
55 centered goals“[1]. Clinical pharmacy services (CPS) have seen significant progress in recent  
56 years, but their implementation and acceptance varies greatly among countries. In Austria, only  
57 15,8% of hospitals have a pharmacy department with even less providing CPS on a regular  
58 basis[2]. The traditional roles and perceptions of medical and pharmacy staff make it challenging  
59 to expand the scope of CPS[3–5]. The positive impact clinical pharmacists have on patient care  
60 is already accepted by medical and nursing staff, however, more needs to be done to change  
61 stakeholders' perspectives and implement necessary legislative changes. Efforts made by clinical  
62 pharmacists in Austria to highlight the necessity of expanding CPS throughout the country  
63 remain unacknowledged[3,4], partly, because larger local multicentre studies are still missing but  
64 also because politicians and stakeholders still do not understand the significance of clinical  
65 pharmacy in promoting patient well-being and improving the cost efficiency of medication  
66 management. This is due to current legislation in Austria failing to acknowledge the importance  
67 of integrating CPS into Austrian hospitals[6]. Furthermore, there is no defined role for clinical  
68 pharmacists in the legislation, and no guidelines for the clinical pharmacist to patient bed ratio  
69 that should be implemented in each hospital[7]. Compared to other European countries,  
70 pharmacists in Austria are not utilised to the same extent as their education and training would  
71 suggest. Urbanczyk et al. have described similar problems in other Central and Eastern European  
72 countries and advocate for broad implementation of CPS[8]. Austria is currently grappling with a  
73 significant deficit in both medical and nursing personnel. However, politicians are both currently  
74 not addressing nor considering other professional groups as potential solutions. This problem  
75 could be partially alleviated by shifting some responsibilities to pharmacists and technicians, as  
76 has been demonstrated in numerous other countries both recently and in the past[9–12]. This

77 study aims to highlight the need for CPS in hospitals across Austria, the acceptance rate of  
78 clinical pharmacy interventions by physicians and the impact on patient safety by having a  
79 clinical pharmacist on the ward team.

80

## 81 **Methods**

82

### 83 **General description**

84 This study was conducted at the Tauernklinikum in Zell am See, Austria, a 360-bed rural clinic in  
85 the state of Salzburg. The clinical pharmacy interventions (CPIs) took place on a 72-bed medical  
86 ward and a 68-bed orthopaedics/traumatology ward. The patients were selected via convenience  
87 sampling by the clinical pharmacist during ward rounds over a 27 month collection period. All  
88 sampled patients were on polypharmacy (i.e. those with five or more prescribed medications)  
89 and over the age of 18. Patient rooms based on ward round groups were chosen and medication  
90 reviews conducted for those patients. The pharmacist would coordinate with the medical staff to  
91 determine when the ward round would start and then join in. Not all patients in the selected  
92 rooms were seen due to time constraints. A similar approach was used for remote type 2b  
93 medication reviews according to the Pharmaceutical Care Network Europe (PCNE)[13], where  
94 the pharmacist would make suggestions over the phone or through electronic patient records,  
95 instead of joining the ward round in person.

96

### 97 **Data collection**

98 The data for this study were collected during the routine work of the hospital's clinical  
99 pharmacist. A document was used to record the CPIs, which was adapted from a French  
100 publication[14] for use in Austria by the Austrian Association of Hospital Pharmacists[15]. The  
101 document contains records of the pharmacist's initials, the date and details of the intervention,

102 the doctor's initials, and the patient's gender and age. The number of patients reviewed for  
 103 medication appropriateness was documented on days when CPIs occurred. Patients without  
 104 polypharmacy or with clearly absent medication-related issues were excluded from the review.  
 105 The pharmacist rated the interventions using a 6-point significance scale according to  
 106 Hatoum[16], as can be seen in table 1.

107 *Table 1 Scale for Recommendations' Potential Impact on Patient Care - Reproduced from Hatoum et al. Evaluation of the contribution of clinical*  
 108 *pharmacists: inpatient care and cost reduction*

Rating	Explanation
X	Adverse significance: Recommendation may lead to adverse outcomes
0	No significance: Recommendation is informational (not specifically related to patient in question)
1	Somewhat significant: Benefit of recommendation to patient could be neutral, depending on professional interpretation
2	Significant: Recommendation could bring care to more acceptable and appropriate level
3	Very significant: Recommendation qualified by potential or existing major organ dysfunction
4	Extremely significant: Information qualified by life-and-death situation

109  
 110 To reduce bias, a random sample of interventions was rated by two medical consultants and two  
 111 medical registrars. The ratings were then evaluated, and a mean was calculated for each coder.  
 112 SPSS was used to perform a two-way model inter-rater reliability (IRR) analysis for the four  
 113 different raters' assessments to determine the intra-class correlation (ICC). The pharmacist's  
 114 rating was then correlated with the ICC to determine if it could be extrapolated to all 1064  
 115 pharmacist intervention ratings.

116  
 117 **Ethics approval**

118 Ethical approval was sought from the Salzburg ethics committee but was not necessary for this  
 119 study, as all patient and staff data has been fully anonymized.

120

## 121 Results

122

### 123 General results

124 A total number of 1064 CPIs were made from the first of December 2020 until the 16<sup>th</sup> of  
125 February 2023. Of these, 866 were undertaken on the medical ward and 198 on the  
126 orthopaedics/traumatology ward. Pharmaceutical orthopaedics/traumatology ward round  
127 participation was started later, from September 2022, with only very few CPIs being made  
128 before that date and only on request. The average age of patients needing an intervention was  
129 73.4 ( $\pm$  13.5) years, internal medical patients being 74 ( $\pm$ 12.5) and orthopaedics/traumatology  
130 surgery patients 70.8 ( $\pm$ 16.9), respectively. In total, female patients accounted for 502 out of 1064  
131 (47.2%) and male patients needing an intervention were 562 out of 1064 (52.8%).

132

### 133 Acceptance rates

134 The pharmacist worked with 37 different physicians.  
135 Of all interventions undertaken, prompt acceptance rate by the physicians involved was 83.5%  
136 (888/1064). In 12.9% (137/1064) of all interventions a change was considered by the physician  
137 but not promptly followed through with (i.e. where laboratory reports were still missing to make  
138 an informed decision or where they wanted to discuss with a medical colleague first). Only six of  
139 all CPIs were immediately declined where:

- 140 • Nicorandil was prescribed at 10mg 1-0-0; the pharmacist suggested to split the dose to  
141 b.i.d. as per SmPC, because of the short elimination half-life of Nicorandil. The  
142 recommendation was rejected because the physician felt that the patient's coronary  
143 artery disease was well adjusted with their current medication.
- 144 • Patient with hyperkalaemia and previously prescribed combination of spironolactone and  
145 furosemide. Re-initiation of furosemide to lower potassium levels was suggested by the



- 146 pharmacist, but the patient was not oedematous at that point and terminal. All  
147 medication apart from analgesia and sedatives were stopped at that juncture.
- 148 • Patient with extensive polypharmacy with a high risk for bleeds was prescribed Pradaxa  
149 at 79 years at 150mg 1-0-1. Pradaxa dose reduction was discussed but the patient's serum  
150 creatinine was 0.7, which indicated good kidney function, so the dose reduction was  
151 rejected.
  - 152 • Pantoprazole was prescribed in a therapeutic dose without indication. The pharmacist  
153 suggested a reduction to 20mg per day (prophylactic dose) as per guideline[17]. The  
154 recommendation was not accepted because of a higher bleeding risk due to the advanced  
155 age of the patient and left at 40mg.
  - 156 • The pharmacist recommended ECG for QTc interval control where a patient had  
157 duloxetine and trazodone prescribed concomitantly with a previous QTc of over 500ms.  
158 The physician stated that the patient had a left bundle branch block and therefore QTc  
159 prolongation was not dangerous and no need for ECG control.
  - 160 • ECG for QTc interval prolongation control was recommended for a patient who was  
161 prescribed Alfuzosine previously with the new addition of pantoprazole, amitriptyline  
162 and prothipendyl. The recommendation was rejected because taking ECGs is an  
163 uncommon measure on the orthopaedics/traumatology ward and would need input from  
164 internal medicine.

165 The remaining 33 interventions were informational (i.e. where the intervention was undertaken  
166 after the patient had left the hospital or the drug(s) concerned were already deprescribed) or not  
167 assessable for acceptance.


168

169 Medication related problems (MRPs)

170 The distribution of types CPIs to address MRPs can be seen in the Figure 1. Drug-drug  
171 interactions were sub-categorised into “to be considered”, “use with caution”, “avoid  
172 combination” and “combination contraindicated”. Altogether, 150 interventions involved drug-  
173 drug interactions, where 4 were categorized as “to be considered”, 21 “use with caution”, 104  
174 “avoid combination” and 21 as “combination contraindicated”.

175

### Distribution of medication related problems (MRP)



176

177 *Figure 1 Bar chart showing distribution of reported MRPs*

178

### 179 Interventions

180 The different types of CPIs that were undertaken can be seen in Table 2.

<b>Types of Interventions</b>	<b>Of 1064 total interventions</b>
New medication prescribed	75
Medication stopped	322
Medication changed to different medication	69
Route of administration altered	4
Medication patient monitoring suggested	111
Optimisation of administration	107
Dose-adjustment undertaken	252
More detailed information provided	101
Organisational-administrative support provided	25
Support provided for document optimisation	32

182 *\*double-categorisation was possible*


183

184 **Significance of CPIs**

185 The overall average score for all 1064 CPIs taken was 2.15. Figure 2 shows the distribution of  
186 these scores. Of all interventions, 99% (1053/1064) were rated “2” or higher; which means a  
187 „significant“, “very significant” or “extremely significant” pharmaceutical intervention. This  
188 indicates that CPIs implemented at Tauernklinikum in Zell am See have been found to  
189 potentially enhance patient care and have the potential to prevent medication errors that may  
190 result in organ failure or fatal outcomes, as measured by the Hatoum scale[16]. Interventions  
191 were rated for significance by the clinical pharmacist. To demonstrate IRR, 5 out of the  
192 interventions rated 1 and 4, respectively and 10 out of the ones rated 2 and 3, respectively were  
193 chosen randomly in Excel functions for rating by two registrar medical doctors and two  
194 consultant medical doctors as illustrated in Figure 3. Of these randomly chosen interventions for  
195 IRR, 5 repetitive interventions have been dismissed before the interventions were listed for MDs


196 to rate them. The clinical pharmacist rated the 25 interventions with a mean of 2.68. The MDs  
197 rated with a mean of 2.45, where one mean was 1.64, one 2.36, one 2.84 and one 2.96.  
198  
199  
200

**SIGNIFICANCE RATING DISTRIBUTION FOR ALL 1064 INTERVENTIONS**



201  
202 *Figure 2 Bar chart showing significance rating distribution for all 1064 interventions*

### SIGNIFICANCE RATINGS FOR 25 INTERVENTIONS RATED BY FOUR MD'S AND ONE PHARMACIST



203

204 *Figure 3 Significance rating for 25 interventions rated by four MDs and one pharmacist*

205

206 The average measurement of ICC for the four consultant coders was 0.749, with a 95%  
 207 confidence interval ranging from 0.538 to 0.878. The average measurement for "absolute  
 208 agreement" was 0.624, with a 95% confidence interval ranging from 0.280 to 0.820. These  
 209 calculated ICC values indicate "good" agreement for average measurement and "moderate"  
 210 agreement for absolute agreement among the four coders. The single measures were low, with  
 211 consistency at 0.428 and absolute agreement at 0.294. However, when correlated with the  
 212 pharmacist's scores, the ICC values for consistency and absolute agreement were 0.780 and  
 213 0.693, respectively, indicating "good" and "moderate" agreement. The Cronbach's alpha of  
 214 0.780 suggests high reliability, especially considering the two different professional rater  
 215 groups[18]. Overall, the high percentage of significant pharmaceutical interventions and the  
 216 agreement between raters suggest that CPIs at Tauernklinikum in Zell am See are effective in  
 217 improving patient welfare and reducing the risk of medication errors and potentially fatal  
 218 outcomes.


219

220 **Cost reduction**

221 In Figure 4, the potential for cost reduction resulting from the suggested pharmaceutical  
222 interventions is shown. These interventions included medication discontinuation, which  
223 accounted for 31.9% (339/1064) of the cases, dose reduction, which accounted for 19.5%  
224 (207/1064) of the cases, and the remaining interventions (48.6%, 518/1064) which did not  
225 directly lead to cost reduction.

226

**COST REDUCTION POTENTIAL**



227

228 *Figure 4 Cost reduction potential*

229 **Discussion**

230 **General results**

231 Considering the aging population and medical specialties, this study suggests that there is a  
232 significant demand for clinical pharmacy input in both, medical and orthopaedics/traumatology  
233 wards. The average age of patients who required interventions was high, which is consistent with  
234 the demographic trend of an ageing population. Additionally, the gender distribution indicates

235 that both male and female patients require CPIs, emphasizing the need for gender-inclusive  
236 healthcare services.

237

### 238 **Acceptance of CPIs**

239 Numerous studies have shown significant variation in acceptance for CPIs across Europe and  
240 the UK[19–24]. In Tauernklinikum Zell am See, approximately 72.3% (986 out of 1064) of  
241 patients sampled require CPIs, indicating a clear need for medicines reconciliation and medicines  
242 review. However, due to staffing constraints, this remains an issue in most Austrian hospitals.

243 The absence of clinical pharmacists in the multidisciplinary ward team poses a gap in optimising  
244 medicine use, reducing MRPs, and improving patient safety. According to the Chief Executive of  
245 the Society of Hospital Pharmacists of Australia, in 2018, the recommended case load for one  
246 hospital pharmacist is a maximum of 30 patients[25]. For a 360-bed hospital, like the one in this  
247 study, this translates to 12 full-time clinical pharmacists. However, the actual staffing level is only  
248 0.75 WTE (whole-time equivalent).

249 The numerous ward rounds with both disciplines represented, the pharmacist and the rounding  
250 physicians, demonstrates valuable collaborative efforts between pharmacists and physicians in  
251 patient care. Multidisciplinary ward round teams have been highlighted to be beneficial for the  
252 patient in many publications[26–29].

253

254 Most recommendations made by the pharmacist were accepted and implemented by physicians.

255 This demonstrates the value of the pharmacist as important member of the healthcare team,  
256 providing expertise in medication management and helping to improve patient outcomes.

257 Not only promptly accepted interventions were recorded but also in 12.9% of interventions, a  
258 change in medication therapy after following up laboratory reports and patients' and/or  
259 colleagues' consultation, was considered by physicians based on the pharmacist's

260 recommendations. This shows that the pharmacist was able to identify and address medication-  
261 related issues that required further attention and intervention.

262

263 In some instances, pharmacist recommendations were declined by physicians immediately.  
264 Reasons for refusal included patient preference, clinical status, and concerns about bleeding risk  
265 or other medical conditions. These cases highlight the importance of effective communication  
266 and collaboration between pharmacist and physicians in the decision-making process for  
267 medication therapy and taking patient preferences into consideration when reviewing  
268 medications as per the Scottish polypharmacy guidance for realistic prescribing[30].

269

270 A small percentage (3.1%) of interventions were purely informational, meaning that the  
271 intervention was undertaken after the patient had left the hospital or the drug(s) concerned had  
272 already been deprescribed. This highlights the challenges of timely communication and  
273 coordination among healthcare providers in a hospital setting, and the need for effective  
274 communication channels and protocols for medication management.

275

## 276 **Types of CPIs**

277 The main type of CPI observed was overdosing, which can be explained by non compliance with  
278 PPI prescribing guidelines. Halving the therapeutic dose to the prophylactic PPI dose is not a  
279 common practice and was addressed on ward rounds many times.

280 However, non-compliance with guidelines was the second most common drug related problem  
281 to be addressed. The interventions in this section included expanding the scope of individual  
282 patient treatment, addressing unfamiliarity with guidelines, and failure to adjust dosages in cases  
283 of organ dysfunction.

284 The third largest category of interventions was inappropriate or inadequate drug administration.

285 The fourth category involved drug-drug interactions.



286 Consultants and registrars were receptive to and accepting of discussions with the pharmacist  
287 regarding MRPs which supports previous findings from work undertaken in Northern Ireland  
288 that highlight the complementary role of pharmacists in the multidisciplinary team on the  
289 ward[10].

290

291

## 292 **Significance of CPIs**

293 According to their independent medical and pharmaceutical significance ratings CPIs undertaken  
294 in this report have the potential to enhance patient care and prevent medication errors that may  
295 result in significant morbidity. It strongly indicates that clinical pharmacist input is an essential  
296 component of multidisciplinary ward teams, leading to more comprehensive and holistic patient  
297 therapy, optimising medication use and ensuring patient safety.

298 By incorporating various professional opinions, including those of pharmacists, psychologists,  
299 physiotherapists, and others, into their patient care plans, physicians are able to broaden their  
300 perspective beyond purely medical considerations, resulting in improved patient care. This  
301 interdisciplinary approach has been supported by numerous studies that have shown its  
302 benefits.[9,11,12,31–33]

303

## 304 **Cost reduction**

305 Discontinuation of medication and dose reductions are important strategies for reducing costs  
306 through CPIs and ensuring prudent medicines use. Schumock et al. have evidenced in 2003 that,  
307 in addition to enhancing patient care, CPIs on the wards have the potential to yield cost savings  
308 in medication expenditure[34].

309

310 Other CPIs that do not directly lead to cost reduction may still have significant impact on  
311 hospitalisation and sickness prevention in the long term due to various reasons. Scott et al. have  
312 showcased between 2000 and 2014 in Northern Ireland that CPS can reduce length of hospital  
313 stays by 2 days, increase time to hospital readmission by 20 days, decrease ward round time by  
314 more than 25 minutes and decrease discharge time by over 90 minutes. Not only were CPS  
315 indirectly cost saving by saving time but also via error reduction (admission drug history error  
316 reduction per patient by 4.2 and improvement of discharge medication accuracy was <1%  
317 compared with 25% by medical staff). [10]

318 These interventions may prevent disease progression or complications, improve patient  
319 outcomes, and enhance patient satisfaction and adherence. It is crucial to consider the  
320 multifactorial nature of cost reduction in healthcare and evaluate various outcomes when  
321 assessing the value and impact of CPIs.

322 Similarly, in 2022 Urbanczyk et al. have demonstrated that CPIs on surgical wards demonstrate  
323 cost-avoidance via prevention of adverse drug events and a cost-benefit ratio of 1:9.5 in the  
324 Polish hospital setting[7].

325

## 326 [Limitations](#)

- 327 • **Limited Generalizability:** The findings of a single site report may not be generalisable to  
328 some other settings, as patient populations and clinical practices can vary widely across  
329 different sites.
- 330 • **Limited Scope of Practice:** The interventions proposed by the pharmacist may be limited  
331 by their scope of practice, which may not encompass all potential MRP or interventions.
- 332 • **Lack of Blinding:** Since the pharmacist is the only individual proposing interventions,  
333 there is no blinding in the study, which may introduce bias in the assessment of  
334 outcomes.

- 335       • Limited Impact Assessment: There was no ability to look at the impact on other  
336       healthcare resource use.

337

338

## 339 Conclusions

340 Overall, this study concludes that the CPIs implemented at Tauernklinikum in Zell am See have  
341 been found to enhance patient care, reduce medication errors, and have the potential to result in  
342 direct cost reduction. The interventions were well accepted by physicians and were rated as  
343 significant by both the clinical pharmacist and medical doctors involved in the study. It  
344 showcases the significant potential and urgent need for the development and expansion of CPS  
345 in a small Austrian hospital.

346 Since it is only a single site report the authors hope it will inform stakeholders and peers to  
347 expand such observational research on the significance, acceptance rates, cost reduction and  
348 need of CPS to other Austrian hospitals. One of the main reasons for the underdevelopment of  
349 CPS in Austria is the drug-oriented curriculum in undergraduate pharmacy degrees with limited  
350 emphasis on patient-centric pharmacotherapy, leading many clinical pharmacists to seek self-  
351 education or additional clinical degrees from overseas[35,36]. Additionally, lack of funding for  
352 CPS through insurance systems hinders their growth in Austria. The current demographic  
353 development leads to a lack of medical and allied health professionals; utilizing CPS could  
354 significantly alleviate medical personnel in certain areas (medicines reconciliation, medication  
355 reviews, patient admission and discharge, communication with extramural interfaces, stocking on  
356 wards, medication preparation for administration, amongst others). Politically, these issues would  
357 best be addressed by obtaining enough data to support CPS implementation in Austrian  
358 hospitals.

## 359 Acknowledgments and affiliations

360 The authors express their gratitude to Dr. Hans-Joachim Miska, Dr. Helene AchRAINER, Dr.  
361 Katharina Heugenhauer, and Dr. Wolfgang Rank for rating the CPIs for IRR. They also thank  
362 Dr. René Wenzel and Dr. Michael Hofer for allowing the pharmacist to work on their wards.  
363 The authors acknowledge the unwavering support, kindness, and patience of Prof. Rudolph  
364 Pointner, the medical director at Tauernklinikum Zell am See. Finally, they extend their  
365 appreciation to all physicians at Tauernklinikum Zell am See for their collaboration and friendly  
366 approach to interdisciplinary co-working, as well as their work culture with a flat hierarchy.

## 367 Funding, conflict of interest and raw data accessibility

368 The authors declare no conflicts of interest and no funding for this study. Raw data are with the  
369 corresponding scientist and can be requested.

## 370 Contributorship

371 SG was involved in all stages of preparing this manuscript. AC, AA and MS helped with  
372 planning, conception, data analysis and interpretation of data and also reviewing the manuscript.  
373

## 374 References

- 375 1 Dreischulte T, van den Bemt B, Steurbaut S. European Society of Clinical Pharmacy  
376 definition of the term clinical pharmacy and its relationship to pharmaceutical care: a  
377 position paper. *Int J Clin Pharm* 2022;**44**:837–42. doi:10.1007/s11096-022-01422-7
- 378 2 Österreichische Apothekerkammer: Apotheke in Zahlen 2020.  
379 <https://www.apothekerkammer.at/infothek/zahlen-daten-fakten/apotheke-in-zahlen-2020>  
380 (accessed 5 Apr 2021).
- 381 3 Stemer G, Laml-Wallner G, Kuegler I, *et al.* Comprehensive evaluation of clinical  
382 pharmacists' interventions in a large Austrian tertiary care hospital. *European Journal of Hospital*  
383 *Pharmacy: Science and Practice* 2012;**19**:529–34. doi:10.1136/ejhpharm-2012-000131
- 384 4 Stemer G, Lemmens-Gruber R. The clinical pharmacist's contributions within the  
385 multidisciplinary patient care team of an intern nephrology ward. *Int J Clin Pharm*  
386 2011;**33**:759–62. doi:10.1007/s11096-011-9548-4

- 387 5 Deibl S, Mueller D, Kirchdorfer K, *et al.* Self-reported clinical pharmacy service provision in  
388 Austria: an analysis of both the community and hospital pharmacy sector-a national study.  
389 *Int J Clin Pharm* 2020;**42**:1050–60. doi:10.1007/s11096-020-01066-5
- 390 6 RIS - Apothekenbetriebsordnung 2005 - Bundesrecht konsolidiert, Fassung vom 08.12.2022.  
391 <https://www.ris.bka.gv.at/GeltendeFassung.wxe?Abfrage=Bundesnormen&Gesetzesnumm>  
392 [er=20003947](https://www.ris.bka.gv.at/GeltendeFassung.wxe?Abfrage=Bundesnormen&Gesetzesnummer=20003947) (accessed 8 Dec 2022).
- 393 7 Urbanczyk K, Wnek P, Roleder T, *et al.* Optimized and cost-effective pharmacotherapy of  
394 vascular surgery patients: evaluation of clinical pharmacy service. *Italian Journal of Vascular and*  
395 *Endovascular Surgery* 2022.
- 396 8 Urbańczyk K, Guntschnig S, Antoniadis V, *et al.* Recommendations for wider adoption of  
397 clinical pharmacy in Central and Eastern Europe in order to optimise pharmacotherapy and  
398 improve patient outcomes. *Frontiers in Pharmacology*  
399 2023;**14**.<https://www.frontiersin.org/articles/10.3389/fphar.2023.1244151> (accessed 10 Aug  
400 2023).
- 401 9 Hellström L, Bondesson A, Höglund P, *et al.* Impact of the Lund Integrated Medicines  
402 Management (LIMM) model on medication appropriateness and drug-related hospital  
403 revisits. *European journal of clinical pharmacology* 2011;**67**:741–52. doi:10.1007/s00228-010-0982-  
404 3
- 405 10 Scott M, Scullin C, Hogg A, *et al.* Integrated medicines management to medicines  
406 optimisation in Northern Ireland (2000-2014): A review. *European Journal of Hospital Pharmacy:*  
407 *Science and Practice* 2015;**22**. doi:10.1136/ejhpharm-2014-000512
- 408 11 Scullin C, Scott MG, Hogg A, *et al.* An innovative approach to integrated medicines  
409 management. *J Eval Clin Pract* 2007;**13**:781–8. doi:10.1111/j.1365-2753.2006.00753.x
- 410 12 Burnett KM, Scott MG, Fleming GF, *et al.* Effects of an integrated medicines management  
411 program on medication appropriateness in hospitalized patients. *Am J Health Syst Pharm*  
412 2009;**66**:854–9. doi:10.2146/ajhp080176
- 413 13 Griese-Mammen N, Hersberger KE, Messerli M, *et al.* PCNE definition of medication  
414 review: reaching agreement. *Int J Clin Pharm* 2018;**40**:1199–208. doi:10.1007/s11096-018-  
415 0696-7
- 416 14 Bedouch P, Charpiat B, Conort O, *et al.* Assessment of Clinical Pharmacists' Interventions in  
417 French Hospitals: Results of a Multicenter Study. *The Annals of pharmacotherapy* 2008;**42**:1095–  
418 103. doi:10.1345/aph.1L045
- 419 15 Arbeitsgemeinschaft österreichischer Krankenhausapotheker. <https://www.aahp.at/>  
420 (accessed 10 May 2021).
- 421 16 Hatoum HT, Catizone C, Hutchinson RA, *et al.* An Eleven-Year Review of the Pharmacy  
422 Literature: Documentation of the Value and Acceptance of Clinical Pharmacy. *Drug*  
423 *Intelligence & Clinical Pharmacy* 1986;**20**:33–48. doi:10.1177/106002808602000105
- 424 17 Wolfgang Fischbach<sup>1</sup>, Jan Bornschein<sup>2</sup>, Jörg C. Hoffmann<sup>3</sup>, Sibylle Koletzko<sup>4</sup>, 5, Alexander  
425 Link<sup>6</sup>, Lukas Macke<sup>7,13</sup>, Peter Malfertheiner<sup>6,7</sup>, Kerstin Schütte<sup>8</sup>, Dieter-Michael  
426 Selgrad<sup>9,10</sup>, Sebastian Suerbaum<sup>11,12,13</sup> Christian Schulz. Aktualisierte S2k-Leitlinie  
427 *Helicobacter pylori* und gastroduodenale Ulkuskrankheit der Deutschen Gesellschaft für

- 428 Gastroenterologie, Verdauungs- und Stoffwechselkrankheiten (DGVS).  
 429 [https://register.awmf.org/assets/guidelines/021-0011\\_S2k\\_Helicobacter-pylori-](https://register.awmf.org/assets/guidelines/021-0011_S2k_Helicobacter-pylori-gastroduodenale-Ulkuskrankheit_2022-07_1.pdf)  
 430 [gastroduodenale-Ulkuskrankheit\\_2022-07\\_1.pdf](https://register.awmf.org/assets/guidelines/021-0011_S2k_Helicobacter-pylori-gastroduodenale-Ulkuskrankheit_2022-07_1.pdf)
- 431 18 Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment  
 432 instruments in psychology. - PsycNET. 1994./doiLanding?doi=10.1037%2F1040-  
 433 3590.6.4.284 (accessed 18 Mar 2021).
- 434 19 Bahrani L, Eriksson T, Höglund P, *et al.* The rate and nature of medication errors among  
 435 elderly upon admission to hospital after implementation of clinical pharmacist-led  
 436 medication reconciliation. *European Journal of Hospital Pharmacy: Science and Practice*  
 437 2014;**21**:156–60. doi:10.1136/ejhpharm-2013-000403
- 438 20 O’Sullivan D, O’Mahony D, O’Connor M, *et al.* The Impact of a Structured Pharmacist  
 439 Intervention on the Appropriateness of Prescribing in Older Hospitalized Patients. *Drugs &*  
 440 *aging* 2014;**31**. doi:10.1007/s40266-014-0172-6
- 441 21 Lozano-Montoya I, Vélez-Díaz-Pallarés M, Delgado-Silveira E, *et al.* Potentially  
 442 inappropriate prescribing detected by STOPP-START criteria: are they really inappropriate?  
 443 *Age Ageing* 2015;**44**:861–6. doi:10.1093/ageing/afv079
- 444 22 Bondesson Å, Eriksson T, Kragh A, *et al.* In-hospital medication reviews reduce unidentified  
 445 drug-related problems. *Eur J Clin Pharmacol* 2013;**69**:647–55. doi:10.1007/s00228-012-1368-5
- 446 23 Somers A, Robays H, De Paepe P, *et al.* Evaluation of clinical pharmacist recommendations  
 447 in the geriatric ward of a Belgian university hospital. *Clin Interv Aging* 2013;**8**:703–9.  
 448 doi:10.2147/CIA.S42162
- 449 24 Pandraud-Riguet I, Bonnet-Zamponi D, Bourcier E, *et al.* Monitoring of Potentially  
 450 Inappropriate Prescriptions in Older Inpatients: A French Multicenter Study. *J Am Geriatr*  
 451 *Soc* 2017;**65**:2713–9. doi:10.1111/jgs.15081
- 452 25 Pharmacist to patient ratios matter: SHPA. AJP. 2018.[https://ajp.com.au/news/pharmacist-](https://ajp.com.au/news/pharmacist-to-patient-ratios-matter-shpa/)  
 453 [to-patient-ratios-matter-shpa/](https://ajp.com.au/news/pharmacist-to-patient-ratios-matter-shpa/) (accessed 6 May 2021).
- 454 26 Stuhec M, Tement V. Positive evidence for clinical pharmacist interventions during  
 455 interdisciplinary rounding at a psychiatric hospital. *Sci Rep* 2021;**11**:13641.  
 456 doi:10.1038/s41598-021-92909-2
- 457 27 Leape LL, Cullen DJ, Clapp MD, *et al.* Pharmacist participation on physician rounds and  
 458 adverse drug events in the intensive care unit. *JAMA* 1999;**282**:267–70.  
 459 doi:10.1001/jama.282.3.267
- 460 28 Kucukarslan SN, Peters M, Mlynarek M, *et al.* Pharmacists on rounding teams reduce  
 461 preventable adverse drug events in hospital general medicine units. *Arch Intern Med*  
 462 2003;**163**:2014–8. doi:10.1001/archinte.163.17.2014
- 463 29 Chisholm-Burns MA, Lee JK, Spivey CA, *et al.* *US pharmacists’ effect as team members on patient*  
 464 *care: systematic review and meta-analyses.* Centre for Reviews and Dissemination (UK) 2010.  
 465 <https://www.ncbi.nlm.nih.gov/books/NBK80549/> (accessed 17 Apr 2023).
- 466 30 Scottish Government Polypharmacy Model of Care Group. Polypharmacy Guidance,  
 467 Realistic Prescribing, 3rd Edition, 2018.

- 468 31 Grimes T, Deasy E, Allen A, *et al.* Collaborative pharmaceutical care in an Irish hospital:  
 469 Uncontrolled before-after study. *BMJ quality & safety* 2014;**23**. doi:10.1136/bmjqs-2013-  
 470 002188
- 471 32 Gillespie U, Alassaad A, Henrohn D, *et al.* A comprehensive pharmacist intervention to  
 472 reduce morbidity in patients 80 years or older: a randomized controlled trial. *Arch Intern Med*  
 473 2009;**169**:894–900. doi:10.1001/archinternmed.2009.71
- 474 33 Spinewine A, Swine C, Dhillon S, *et al.* Effect of a collaborative approach on the quality of  
 475 prescribing for geriatric inpatients: a randomized, controlled trial. *J Am Geriatr Soc*  
 476 2007;**55**:658–65. doi:10.1111/j.1532-5415.2007.01132.x
- 477 34 Schumock GT, Butler MG, Meek PD, *et al.* Evidence of the Economic Benefit of Clinical  
 478 Pharmacy Services: 1996–2000. *Pharmacotherapy: The Journal of Human Pharmacology and Drug*  
 479 *Therapy* 2003;**23**:113–32. doi:https://doi.org/10.1592/phco.23.1.113.31910
- 480 35 Rose O, Derendorf H, Erzkamp S, *et al.* Development of clinical pharmacy services  
 481 in Australia, Austria, Belgium, Bosnia-Herzegovina, Canada, Germany, Japan, Kosovo,  
 482 Switzerland, the Netherlands, Thailand, USA and correlation with educational standards,  
 483 level of research, and implemen. *Int J Clin Pharmacol Ther* 2018;**56**:518–30.  
 484 doi:10.5414/CP203264
- 485 36 Stemer MG. Renal Clinical Pharmacy Services. ;:191.

486

## 487 Figures

- 488 Figure 1 Bar chart showing distribution of reported clinical pharmacy interventions ..... 8  
 489 Figure 2 Bar chart showing significance rating distribution for all 1064 interventions ..... 10  
 490 Figure 3 Significance rating for 25 interventions rated by four MDs vs one pharmacist ..... 11  
 491 Figure 4 Cost reduction potential ..... 12  
 492

## 493 Tables

- 494 Table 1 Types of clinical pharmacy interventions undertaken ..... **Error! Bookmark not defined.**  
 495