



## Is education level, as a proxy for socio-economic position, related to device-measured and self-reported sedentary behavior in European older adults? A cross-sectional study from the SITLESS project

Rodriguez Roca, B., Tully, M. A., Sansano-Nadal, O., Caserotti, P., Coll-Planas, L., Roqué, M., Brønd, J., Blackburn, N. E., Wilson, J. J., Rothenbacher, D., McIntosh, E., Deidda, M., Andrade-Gómez, E., & Giné-Garriga, M. (2023). Is education level, as a proxy for socio-economic position, related to device-measured and self-reported sedentary behavior in European older adults? A cross-sectional study from the SITLESS project. *Frontiers in public health*, 11, 1-9. Advance online publication. <https://doi.org/10.3389/fpubh.2023.1296821>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Frontiers in public health

**Publication Status:**  
Published online: 19/12/2023

**DOI:**  
[10.3389/fpubh.2023.1296821](https://doi.org/10.3389/fpubh.2023.1296821)

**Document Version**  
Publisher's PDF, also known as Version of record

**General rights**  
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk).



## OPEN ACCESS

## EDITED BY

Kathleen Potempa,  
University of Michigan, United States

## REVIEWED BY

Janet Larson,  
University of Michigan, United States  
Pawel Adam Piepiora,  
Wroclaw University of Health and Sport  
Sciences, Poland

## \*CORRESPONDENCE

Elena Andrade-Gómez  
✉ elena.andrade@unirioja.es

RECEIVED 19 September 2023

ACCEPTED 20 November 2023

PUBLISHED 19 December 2023

## CITATION

Rodríguez Roca B, Tully MA, Sansano-Nadal O, Caserotti P, Coll-Planas L, Roqué M, Brønd J, Blackburn NE, Wilson JJ, Rothenbacher D, McIntosh E, Deidda M, Andrade-Gómez E and Giné-Garriga M (2023) Is education level, as a proxy for socio-economic position, related to device-measured and self-reported sedentary behavior in European older adults? A cross-sectional study from the SITLESS project. *Front. Public Health* 11:1296821. doi: 10.3389/fpubh.2023.1296821

## COPYRIGHT

© 2023 Rodríguez Roca, Tully, Sansano-Nadal, Caserotti, Coll-Planas, Roqué, Brønd, Blackburn, Wilson, Rothenbacher, McIntosh, Deidda, Andrade-Gómez and Giné-Garriga. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

# Is education level, as a proxy for socio-economic position, related to device-measured and self-reported sedentary behavior in European older adults? A cross-sectional study from the SITLESS project

Beatriz Rodríguez Roca<sup>1</sup>, Mark A. Tully<sup>2</sup>, Oriol Sansano-Nadal<sup>3,4</sup>, Paolo Caserotti<sup>5</sup>, Laura Coll-Planas<sup>6</sup>, Marta Roqué<sup>7</sup>, Jan Brønd<sup>5</sup>, Nicole E. Blackburn<sup>8</sup>, Jason J. Wilson<sup>9</sup>, Dietrich Rothenbacher<sup>10</sup>, Emma McIntosh<sup>11</sup>, Manuela Deidda<sup>11</sup>, Elena Andrade-Gómez<sup>12\*</sup> and Maria Giné-Garriga<sup>3,13</sup>

<sup>1</sup>Department of Physiatry and Nursing, Faculty of Health Sciences, University of Zaragoza, Zaragoza, Spain, <sup>2</sup>Institute of Mental Health Sciences, School of Health Sciences, Ulster University, Newtownabbey, United Kingdom, <sup>3</sup>Department of Physical Activity and Sport Sciences, Faculty of Psychology, Education and Sport Sciences (FPCEE) Blanquerna, Ramon Llull University, Barcelona, Spain, <sup>4</sup>School of Health and Sport Sciences (EUSES), Rovira i Virgili University, Amposta, Spain, <sup>5</sup>Department of Sports Science and Clinical Biomechanics, Center for Active and Healthy Ageing (CAHA), Syddansk Universitet, Odense, Denmark, <sup>6</sup>Fundació Salut I Envelliment (Foundation on Health and Ageing) – UAB, Universitat Autònoma de Barcelona, Barcelona, Spain, <sup>7</sup>Iberoamerican Cochrane Centre – Biomedical Research Institute Sant Pau (IIB Sant Pau), CIBER Epidemiologia y Salud Pública (CIBERESP), Barcelona, Spain, <sup>8</sup>Institute of Nursing and Health Research, School of Health Sciences, Ulster University, Newtownabbey, United Kingdom, <sup>9</sup>Sport and Exercise Sciences Research Institute, School of Sport, Ulster University, Newtownabbey, United Kingdom, <sup>10</sup>Institute of Epidemiology and Medical Biometry, Ulm University, Ulm, Germany, <sup>11</sup>Health Economics and Health Technology Assessment (HEHTA), Institute of Health and Wellbeing (IHW), University of Glasgow, Glasgow, United Kingdom, <sup>12</sup>Department of Nursing, Faculty of Health Sciences, University of La Rioja, Logroño, Spain, <sup>13</sup>Department of Physical Therapy, Faculty of Health Sciences (FCS) Blanquerna, Ramon Llull University, Barcelona, Spain

**Background:** Sedentary behavior (SB) is a determinant of health in older adult people. Educational level is a primary driver of health disparities and is demonstrated to be a reliable measure of socioeconomic position. We aimed to examine the associations between educational level and self-reported along with device-measured SB in older adults living in Europe and the association of mentally active and passive SB domains with the educational level and gender in these associations.

**Methods:** The design is cross-sectional. One thousand three hundred and sixty participants aged 65 and over (75.3±6.3 years old, 61.8% women) participated. Inclusion criteria were scored with the Short Physical Performance Battery. Variables that describe the sample were assessed with an interview, and device-measured SB was assessed with an accelerometer. SB was assessed with the Sedentary Behavior Questionnaire and an accelerometer. Multiple linear regression models were used to study the association between the level of education and SB.

**Results:** Participants self-reported an average of 7.82 (SD: 3.02) daily waking hours of SB during weekend days, and the average of device-measured SB was 11.39 (1.23) h. Total mentally active SB (weekdays and weekends) was associated with the education level ( $p < 0.000$ ). Participants were more sedentary during the week

than during weekends, regardless of level of education ( $p < 0.000$ ). Education level was significantly associated with self-reported mean hours per day in 46SB ( $p = 0.000$ ;  $R=0.026$ ; 95%CI).

**Conclusion:** Low education level in older adults is associated with self-reported SB but not with objective SB measures.

#### KEYWORDS

older adults, sedentary behavior, education level, mentally active sedentary behavior, mentally passive sedentary behavior older adult, sedentary, mentally passive sedentary behavior, accelerometer

## 1 Introduction

Sedentary behavior (SB) has emerged as a new risk factor for health (1) and is defined as a behavior with low energy expenditure (1.5 metabolic equivalents [METs]), while in a sitting or reclining posture during waking hours, distinct from the simple absence of physical activity (PA) (2). SB has been positively associated with obesity and chronic diseases such as cardiovascular disease (3), decreased cognitive function (4), risk of depression (5), diabetes (4), some types of cancer and mortality (6), and increased vulnerability to falls and fractures (7). Importantly, replacing SB with the same amount of moderate-to-vigorous physical activity (MVPA) may contribute to better quality of life in older adults (8). Older adults are the most sedentary of all age groups (9, 10). SB tends to be more frequent after retirement from full-time work (11), when the older adult is alone at home (12) as well as after lunch and evening meals. Recent studies concluded that older adults spend an average of 9.4 h/day in SB according to objective measures (13), which represents between 65 and 79% of their awake time (14).

Several accelerometer devices have proven to be reliable and valid (e.g., ActiGraph (15, 16), activPAL (17), and Axivity (18)). Self-reported questionnaires, such as the Sedentary Behavior Questionnaire (SBQ), have also been used to subjectively measure SB in large samples due to their low administration cost, ease of administration, and ability to distinguish between different domains where SB takes place (19). Nevertheless, data from self-reported instruments should be taken with caution due to their weak-to-moderate correlation with device-measured SB in the older adult population (20, 21). However, self-reported questionnaires can collect the domains where SB takes place during a usual day.

Time spent in different domains of SB (such as screen time or transport) is likely to differ between older adults. Most studies have focused on only one SB domain or on overall SB time (22), making it difficult to assess the correlates of various domains (10). Recent studies have classified SB into two categories according to its cognitive stimulation: mentally active and passive SB. Mentally active SB, such as reading a book or playing a musical instrument, could be protective and promote healthy cognition, whereas mentally passive SB such as watching television has been associated with increased levels of depressive symptoms (23). For example, Hallgren et al. (24) concluded that not all sedentary activities have equal negative impacts on health; mentally passive SB was more associated with declines in physical function and mental health outcomes than mentally active SB (24).

Educational level seems to be associated with SB in adults according to European-based studies (25–27). The highest education level achieved has shown to be a good indicator of socioeconomic position (SEP) (26) and has been described as a fundamental cause of inequalities on health (28). It is considered a protective factor for the health of the older adult population and establishes the basis for a healthy lifestyle, and it is an indicator of stability throughout life (29). The highest prevalence of unhealthy behaviors happens among the population of lower SEP (30–32).

There is evidence that older persons from socioeconomically poor backgrounds are more likely to be sedentary (10). These factors include a complex interplay between social, physical, and cultural settings and health-related characteristics. On the one hand, prior research from middle-class and high-income nations ( $n=101$ ) found an inverse relationship between SB and a high SEP that was linked to a higher education level (33). Additionally, it was associated with higher occupation-based SB (34) and higher PA levels in subjects. Conversely, older persons with lower SEP and lower education levels are likely to engage in SB activities such as watching television (35) more frequently than older adults with higher SEPs and higher education levels. On the other hand, some research revealed the contrary, showing that the greater the adults' level of education, the greater their overall SB time (36).

To further prioritize future investments, it is necessary to identify and understand modifiable factors such as SB and its domains (mentally active and passive), as well as the extent to which it is related to education levels, in order to develop appropriate policies and effective interventions to reduce SB time and modify its patterns.

The activities carried out in each age group are associated with gender. In general, in the older adult population, men are more sedentary than women, analyzing them from the objective and subjective perspective (37, 38). Additionally, disruptions to sedentary behaviors also vary by gender. For example, women perform more SB interruptions than men. In women, they take place in the morning and, in men, throughout the day (37, 39).

Given that the SB levels in the older adult population are increasing, and the current literature on the socioeconomic determinants of SB in European older adults is still limited (40), we aimed to examine the associations between the educational level and self-reported as well as device-measured SB in a large sample of community-dwelling older adults living in four European countries. Furthermore, given that there is not enough bibliography that analyzes the relationship between gender and mentally active or passive sedentary behaviors, we have decided that our secondary aims were

to assess the association of several SB domains including mentally active and passive SB with older adults' educational level and to assess the role of gender in the associations.

## 2 Materials and methods

### 2.1 Design and study population

The SITLESS study was a cross-sectional study. It was a multicenter pragmatic three-armed parallel randomized controlled trial. Community-dwelling older adults aged 65 years or older, with a score on the Short Physical Performance Battery (SPPB) of four or above (41), who were insufficiently active and/or reported high levels of SB (42), were recruited in study centers in Denmark, Spain, United Kingdom, and Germany according to their existing primary prevention pathways. The present cross-sectional study uses data from the baseline assessments. A total of 1,360 community-dwelling older adults (75.3 ± 6.3 years old, 61.8% women) were analyzed at baseline. The Ethics and Research Committee of each intervention site approved the study design. Participation was voluntary, and all participants signed informed consent before the start of the study.

### 2.2 Procedure

In each of the health centers participating in the study, potential patients will be asked about their participation. The members of the research team will report on the objectives of the study. In addition, there will be a strategy to attract future participants through audiovisual means. Once patients begin to show interest in participating, the scientific team will be in charge of verifying that they meet all the inclusion criteria to be able to comment on the study. The study protocol can be found elsewhere (43).

### 2.3 Measures

Personal information regarding age, gender, country, number of current medical conditions, and body mass index (BMI) was collected by means of a structured interview in the study centers. Weight and height were objectively measured by a trained researcher using a TANITA BC 420 and a SECA 213 portable stadiometer, respectively, to derive the participants' BMI.

The highest education level achieved was collected at the interview with the following answer options: (a) I cannot read or write; (b) primary education (primary education, preindustrial learning, elementary school); (c) secondary education (high school, professional learning, industrial official, industrial expert, commercial sector); (d) university; (e) unwilling to answer.

According to Hallgren et al., we operationalized mentally active SB as the first five domains of the Sedentary Behavior Questionnaire, which included computer work, paperwork, reading, playing an instrument, and artwork. We operationalized mentally passive SB as the other four domains: television, listening to music, using the phone, and driving in a car (24). Participants self-reported the number of hours spent sitting on a weekday and on a weekend day in different domains using the Sedentary Behavior Questionnaire (SBQ).

All participants wore an ActiGraph wGT3X-BT triaxial accelerometer (ActiGraph, LLC, Pensacola, FL) on their dominant hip during waking hours for 7 consecutive days, removing it only for water-based activities such as bathing or swimming and to sleep during the night. Participants recorded the wear time in an activity diary. The devices were initialized to collect data at 30 Hz. To be included in the analysis, an accelerometer record needed to contain at least 4 valid days (including at least 1 weekend day), with a valid day defined as containing at least 600 min (10 h/d) of wear time as in previous studies (44). Non-wear time was defined using a two-window system: a 90-min window for checking for consecutive zero counts and another 30-min upstream and window for checking for more than 2 min of non-zero counts (45). Due to some participants wearing the ActiGraph during nighttime sleeping, a maximum daily wear time was set at 19 h using a pragmatic choice based on participants' diaries and sleep time duration recommendations for older adults (46). For relevant participants, a log diary was used to determine daily wear time when awake. SB was defined as <100 counts per minute (CPM) (47) on the vertical axis. Values were normalized to the total wear time. Raw accelerometer data were analyzed using ActiLife v6.13.3 software with the normal filter and summarized into 10-s epochs, as have been recommended for the estimation of SB in clinical older adult populations (48). Total daily SB time including time spent sitting and lying was used for data analysis.

### 2.4 Statistical analyses

Baseline cross-sectional characteristics were presented descriptively as mean and standard deviation (SD) for continuous variables and percentage for categorical variables. The normality of the variables was explored with the Kolmogorov–Smirnov test, and owing to its positive result, the non-parametric Kruskal–Wallis test was utilized. To compare the qualitative variables, the  $\chi^2$  test was used.

Analysis of variance (Kruskal–Wallis test) was used to examine differences across levels of education and continuous variables: age, number of medical conditions, self-reported hours per week of SB during weekdays and weekend days, and accelerometry derived daily hours of SB.

The associations between educational level and total daily SB time, time in each SB domain of the SBQ, and overall, mentally active and passive SB, were analyzed through multiple linear regression for each outcome with covariates: age, gender, country, number of medical conditions, and category of BMI (obese when BMI was  $\geq 30$  kg/m<sup>2</sup>, overweight when BMI was 25–29.9 kg/m<sup>2</sup>, normal weight when BMI was 18.5–24.9 kg/m<sup>2</sup>, underweight when BMI was less than 18.4 kg/m<sup>2</sup>). The results represented for the whole week, during week, and weekend days separately. Models with device-measured SB were also adjusted for accelerometer wear time. The results were reported as unstandardized regression coefficients with 95% confidence intervals. All analyses were conducted with IBM SPSS Statistics 26.0. The significance level was set at  $p < 0.05$ .

## 3 Results

Of the overall SITLESS participants ( $n = 1,360$ ), 137 participants were excluded as they did not meet the pre-specified ActiGraph wear

time criteria or did not complete the self-reported questionnaire and interview. A final subsample of 1,223 participants (75.20 ± 6.19 years old, 61.4% women) returned valid accelerometer data and completed the SBQ and the interview. N = 283 reported having primary education (23.14%, 77.06 ± 5.95 years old), 662 secondary education (54.13%, 75.16 ± 6.10 years old), and 278 university education (22.73%, 73.38 ± 6.11). In the primary education level group, 41.9% were obese whereas in the secondary and university education level groups. The overall self-reported mean hours per day in SB were 7.83 ± 3.04 on weekdays and 7.54 ± 2.91 at weekends (Table 1).

Participants with higher education levels spent more time in SB (both self-reported and objectively measured) and showed a higher number of daily steps than their counterparts with lower education levels ( $p < 0.000$ ). Regardless of their education level, participants were more sedentary during the week than at weekends ( $p < 0.000$ ) (Table 1).

Table 2 shows the results of the Kruskal–Wallis test that was used to examine differences between level of education and mentally active or passive SB during weekdays and weekend days. The distribution of mentally active and passive SB was significantly different between levels of education. During weekdays and weekend days, total mentally active SB was associated with the education level ( $p < 0.000$ ).

During weekdays, the average of daily hours spent in mentally active SB was 2.23 ± 1.96 in the primary education group, 2.82 ± 1.85 in the secondary education group, and 3.17 ± 2.01 in the university education group. Doing paperwork, reading or playing an instrument was the most common mentally active activities. During weekdays, mentally passive SB (watching television, listening to music, talking on the phone while sitting, or sitting during transport) was higher in participants with primary and secondary education than university education ( $p < 0.001$ ). The mean daily hours spent in mentally passive SB during weekdays were 5.13 ± 2.38 in participants with primary education, 5.23 ± 2.22 in participants with secondary education, and 4.59 ± 2.04 in participants with university-level education ( $p < 0.001$ ) (Table 2).

Table 3 shows the multiple linear regression models with daily SB (measured with SBQ questionnaire and accelerometry) and covariates (level of education, age, gender, comorbidity, and BMI categories). The models explained a relatively small amount of the variability. Education level was associated with self-reported daily SB time ( $R^2 = 0.026$ ,  $p < 0.000$ ), mentally active SB ( $R^2 = 0.028$ ,  $p < 0.000$ ), and mentally passive SB ( $R^2 = 0.034$ ,  $p < 0.000$ ), with higher amounts of self-reported SB time related to lower education levels. However, it was not associated with accelerometer-derived daily SB ( $R^2 = 0.052$ ,  $p = 0.149$ ). BMI categories were associated with self-reported SB ( $R^2 = 0.026$ ,  $p < 0.013$ ) and SB accelerometer-derived ( $R^2 = 0.034$ ,  $p < 0.000$ ), with higher amounts of SB time related to higher BMI.

### 4 Discussion

The aim of this cross-sectional study was to examine the relationship between the educational level and self-reported as well as device-measured SB. Our results have shown that education level was significantly associated with self-reported SB but not with total daily SB time assessed with accelerometry. In addition, significant differences were found among mentally active and passive SB. Mentally active SB was higher in highly educated participants, and mentally

TABLE 1 Characteristics of the overall sample according to the education level and divided by gender.

	Primary education			Secondary education			University	p
	Male (n = 66)	Female (n = 217)	Total (n = 283)	Male (n = 266)	Female (n = 396)	Total (n = 662)	Total (n = 278)	
Age (years), mean (SD)	77.56 (5.58)	76.91 (6.06)	77.06 (5.95)	75.99 (6.33)	74.61 (5.88)	75.16 (6.10)	73.38 (6.11)	0.000*
SBQ hours/day, mean (SD)	6.89 (2.39)	7.42 (3.2)	7.30 (3.04)	7.84 (2.71)	8.08 (2.92)	7.99 (2.84)	7.64 (2.61)	<0.001*
SBQ WD hours/day, mean (SD)	6.94 (2.59)	7.47 (3.37)	7.35 (3.21)	7.92 (2.86)	8.14 (3.11)	8.05 (3.01)	7.77 (2.86)	
SBQ WE hours/day, mean (SD)	6.74 (2.36)	7.27 (3.28)	7.15 (3.09)	7.63 (2.81)	7.92 (2.94)	7.80 (2.89)	7.31 (2.70)	
Accelerometry:								<0.000*
Hours daily sedentary time, Mean (SD)	11.74 (1.19)	11.23 (1.30)	11.35 (1.29)	11.48 (1.28)	11.10 (1.22)	11.25 (1.26)	11.39 (1.23)	
Number of daily steps, number (SD)	4575.60 (2247.87)	4839.22 (2168.86)	4777.74 (2186.36)	4701.12 (2774.45)	4996.22 (2342.53)	4877.65 (2527.09)	5571.63 (2844.46)	

\*Kruskal–Wallis test.

TABLE 2 Description of mentally active and passive SB time on weekdays and weekend days according to education levels.

	Low primary education	Secondary education	University	<i>p</i>
	Mean (SD)	Mean (SD)	Mean (SD)	
Mentally active SB during weekdays (hours/day)				0.000
Computer	0.34 (0.87)	0.42 (0.81)	0.31 (0.70)	
Paperwork	0.6 (0.85)	0.71 (1.01)	1.16 (1.36)	
Reading	0.95 (0.96)	1.26 (1.05)	1.35 (1.12)	
Play instrument	0.02 (0.27)	0.04 (0.25)	0.07 (0.22)	
Artwork	0.54 (1.14)	0.37 (0.89)	0.26 (0.74)	
Total	2.23 (1.96)	2.82 (1.85)	3.17 (2.01)	
Mentally passive SB during weekdays (hours/day)				0.001
TV	3.46 (1.60)	3.33 (1.46)	2.85 (1.53)	
Listening music	0.72 (1.24)	0.63 (1.01)	0.55 (0.84)	
Phone	0.34 (0.36)	0.44 (0.60)	0.42 (0.56)	
Car (co-driver)	0.60 (0.78)	0.81 (0.90)	0.76 (0.68)	
Total	5.13 (2.38)	5.23 (2.22)	4.59 (2.04)	
Mentally active SB during weekends (hours/day)				0.000
Computer	0.36 (0.83)	0.44 (0.91)	0.24 (0.55)	
Paperwork	0.5 (0.84)	0.62 (1.01)	0.82 (0.95)	
Reading	0.98 (1.06)	1.27 (1.06)	1.41 (1.12)	
Play instrument	0.02 (0.18)	0.03 (0.22)	0.07 (0.32)	
Artwork	0.48 (1.11)	0.31 (0.79)	0.20 (0.64)	
Total	2.10 (1.83)	2.55 (1.81)	2.71 (1.75)	
Mentally passive SB during weekends (hours/day)				0.001
TV	3.47 (1.68)	3.37 (1.50)	2.80(1.53)	
Listening music	0.63 (1.12)	0.66 (1.01)	0.59 (0.87)	
Phone	0.33 (0.39)	0.42 (0.53)	0.37 (0.47)	
Car (co-driver)	0.59 (0.66)	0.79 (0.80)	0.82 (0.83)	
Total	5.04 (2.26)	5.25 (2.15)	4.60 (2.02)	

passive SB was higher in participants with primary and secondary education levels, both during weekdays and weekend days. Gender was associated with overall SB but did not differ between mentally active or passive SB.

To determine the context of sedentary behavior, it is necessary to consider the differences observed in our study between the hours of sedentary behavior measured objectively or subjectively. The subjects who participated in the study were more sedentary than they thought. This fact could be justified through the theory of Parker and Bates in which they conclude that “any difference is information, regardless of whether it makes a difference to someone or something.” On the one hand, the objective part, which in our study would coincide with the sedentary lifestyle measured through the accelerometer, would be considered independent of the observer and the situation. On the other hand, subjective data are considered by the person themselves, and this information makes a difference from someone’s point of view (49, 50). It is easier for participants to be able to remember the time they dedicate to a specific sedentary activity than to remember the total sedentary time (51), although it is important to keep in mind that the questionnaire used is composed of multiple items and tends to

underestimate total sedentary time (52). It is important to keep in mind that very active people were not included in the study, and therefore, this could reduce the magnitude of the correlations; this has happened in other studies in which when correlating the objective measures with the subjective ones it is found that these, and they do so weakly (53).

Our results have shown that education level was significantly associated with self-reported SB and was not significantly associated with device-measured total daily SB, similar to the results obtained in other studies (54–58). Our findings suggest that the higher the education level, the lower the self-reported SB time. This could be explained by the fact that people with low levels of education are not as conscious of the negative health impact of prolonged maintenance of SB than the more educated counterparts, such as a higher risk of obesity (59), or increased cardiovascular disease (60) or they may live in environments which encourage sitting and place less value on the benefits of sitting less. The fact that people with lower education levels tend to spend more time in SB should be considered when addressing public health needs (25–27).

In our study, a higher educational level was associated with mentally active SB including activities such as reading, doing

TABLE 3 Multiple linear regression model adjusted for level of education, age, gender, comorbidities, and BMI categories.

	R <sup>2</sup>	Level of education			Age			Gender			Comorbidity			BMI categories		
		Root mean square	F	p	Root mean square	F	p	Root mean square	F	p	Root mean square	F	p	Root mean square	F	p
Total daily SB time with ActiGraph	0.052	2.913	1.909	0.149	42.683	27.973	0.000	38.172	25.017	0.000	1.399	0.917	0.338	5.544	3.633	0.057
SBQ total	0.026	3238.701	8.374	0.000	97.62	0.252	0.615	1898.98	4.910	0.027	152.242	0.394	0.531	4359.11	11.27	0.001
SBQ weekend	0.026	69.290	8.314	0.000	12.451	1.494	0.222	38.647	4.637	0.031	26.092	3.131	0.077	51.317	6.157	0.013
SBQ weekday	0.024	65.069	7.222	0.001	0.319	0.035	0.851	38.798	4.306	0.038	0.180	0.020	0.888	106.901	11.865	0.001
SBQ mentally active	0.028	4.098	16.440	0.000	0.185	0.744	0.389	0.185	0.744	0.389	0.056	0.223	0.637	0.024	0.096	0.757
SBQ mentally passive	0.034	2.755	8.101	0.000	0.039	0.115	0.735	0.583	1.713	0.191	0.271	0.798	0.372	5.553	16.329	0.000

paperwork, or playing instruments. In contrast, mentally passive SB activities such as watching television, listening to music, talking on the phone, or traveling in a car as a co-driver were associated with low educational levels. These differences may be due to different causes. For example, people who have attained higher levels of education tend to spend more time in cognitively demanding activities which they find more stimulating than passive activities (61). People with higher levels of education are likely to have more disposable income to spend on leisure time activities, which can positively impact health (62). In addition, they spend less time on mentally passive sedentary activities (58). Our results are like other studies; Van Cauwenberg and Prince concluded that older adults with higher levels of education had a decreased television time, and those with lower education levels had an increased overall mentally passive sedentary time (56, 58). In addition, similar studies reported that higher education level was linked to mentally active SB such as reading (63, 64). In terms of the amount of time adults spend in mentally active and passive SB, Hallgren et al. reported that the average times in mentally active and passive SB time were 193 and 258 min per week, respectively (24).

The current world situation is complex, especially if we focus on the way in which people from previous generations tried to maintain an intellectual and emotional balance. If this imbalance is reached, organic diseases caused by mental illnesses (65) can develop. Therefore, it is important to know the characteristics of older people who tend to be sedentary and be able to influence the reduction of sitting. In our study, knowing that the educational level is related to the level of sitting is an interesting fact that can help develop new health policies focused on adulthood. This future line of work would focus on the responsibility we have as human beings to provide equitable care for the next generations, related to scientific discoveries on health and life protection (65). In this case, the health and educational systems must join forces to reinforce the personal and collective security of individuals (65).

There is an association between gender and sedentary behaviors, and it varies throughout life. In each of the age groups, there is an association between certain sedentary activities and gender. In our study, gender was associated with device-measured and subjective measures of SB. Device-measured SB was higher in men with primary education levels than with higher levels of education. In contrast, self-reported SB was higher in women with all levels of education and increased with the level of education. To date, there are studies that present results similar to ours in which it is concluded that gender was associated with SB (66, 67). However, there are other scientific studies carried out in the older adult population in which men are more sedentary than women, analyzing it from the objective and subjective perspective (68, 69).

One of the strengths of this study was to analyze the association between educational level and sedentary time measured objectively and subjectively in a cohort of community-dwelling older adults.

from four European countries. In addition, it also studies the difference between sedentary time, depending on whether the activity performed is mentally active or passive. This study also has limitations. First, the accelerometer had some well-known limitations for assessing posture that could be overcome using an inclinometer (e.g., time spent standing is likely to be classified as sedentary using an accelerometer). Second, by excluding active people from this study, it is possible that the correlation presented is affected, and furthermore, the data obtained would only be extrapolated to insufficiently active older people.

## 5 Conclusion

Findings from this study reveal that there is a significant association between the education level with subjective SB in European older adults but not for device-measured SB. In the decade of healthy aging (2020–2030) (WHO), a worldwide situation in which life expectancy is increasing, but living longer does not go hand in hand with living healthier, identifying people at higher risk of poorer health outcomes is relevant and compelling. Having low education levels has shown to be an indicator of health inequalities such as higher sedentary time, which, in turn, may be related to decreased functional performance and risk of dependency in the longer term. There is a need to identify the most vulnerable groups and target them as a priority for health-based interventions. There is a need to enhance higher education levels among the population, focus on those with lower education levels, and design preventive strategies to decrease overall SB (and enhance movement), and more specifically mentally passive sedentary activities.

## Scope statement

This study is a step forward toward the knowledge of the association between the education level of older adult people and sedentary behavior. In addition, we would like to study the association between the mentally active or passive activities and level of education. These results are significantly important since low education level in older adults is associated with self-reported sedentary but not with an objective sedentary measure. These results are important to identify the groups of older adults who need specifically interventions to decrease sedentary behavior. This study is based on the major problem of older adult people, sedentary behavior.

That means the readers of this journal will definitely be interested in reading this study as it is a novel topic, and it can be used for improving the knowledge about it.

## Data availability statement

The original contributions presented in the study are included in the article/[Supplementary material](#), further inquiries can be directed to the corresponding author.

## References

- Owen N, Sugiyama T, Eakin EE, Gardiner PA, Tremblay MS, Sallis JF. Adults' sedentary behavior: determinants and interventions. *Am J Prev Med.* (2011) 19:189–96. doi: 10.1016/j.amepre.2011.05.013
- Tremblay MS, Aubert S, Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE, et al. Sedentary behavior research network (SBRN)-terminology consensus project process and outcome. *Int J Behav Nutr Phys Act.* (2017) 14:75. doi: 10.1186/s12966-017-0525-8
- Dunstan DW, Barr ELM, Healy GN, Salmon J, Shaw JE, Balkau B, et al. Television viewing time and mortality the Australian diabetes, obesity and lifestyle study (AusDiab). *Circulation.* (2010, 2010) 26:384–91. doi: 10.1161/CIRCULATIONAHA.109.894824
- Ku PW, Steptoe A, Liao Y, Hsueh MC, Chen LJ. A cut-off of daily sedentary time and all-cause mortality in adults: a meta-regression analysis involving more than 1 million participants. *BMC Med.* (2018) 16:1–9. doi: 10.1186/s12916-018-1062-2
- Eriksson M, Näs S, Berginström N, Nordström P, Hansson P, Nordström A. Sedentary behavior as a potential risk factor for depression among 70-year-olds. *J Affect Disord.* (2020) 15:605–8. doi: 10.1016/j.jad.2019.11.035
- Biswas A, Oh PI, Faulkner GE, Bajaj RR, Silver MA, Mitchell MS, et al. Sedentary time and its association with risk for disease incidence, mortality, and hospitalization in adults: a systematic review and meta-analysis. *Ann Intern Med.* (2015) 162:123–32. doi: 10.7326/M14-1651
- Gomes dos Santos R, Medeiros JC, Dittrich Schmitt B, Meneguci J, Teles Santos DA, Damião R, et al. Comportamento Sedentário em Idosos: Uma Revisão Sistemática. / Sedentary behavior in elderly: a systematic review. *Motricidade.* (2015) 11:171–86. doi: 10.6063/motricidade.3184
- Yasunaga A, Shibata A, Ishii K, Inoue S, Sugiyama T, Owen N, et al. Replacing sedentary time with physical activity: effects on health-related quality of life in older Japanese adults. *Health Qual Life Outcomes.* (2018) 16:1–5. doi: 10.1186/s12955-018-1067-8

## Author contributions

BR-R: Writing – original draft, Writing – review & editing. MT: Writing – original draft, Writing – review & editing. OS-N: Writing – original draft, Writing – review & editing. PC: Writing – original draft, Writing – review & editing. LC-P: Writing – original draft, Writing – review & editing. MR: Writing – original draft, Writing – review & editing. JB: Writing – original draft, Writing – review & editing. NB: Writing – original draft, Writing – review & editing. JW: Writing – original draft, Writing – review & editing. DR: Writing – original draft, Writing – review & editing. EM: Writing – original draft, Writing – review & editing. MD: Writing – original draft, Writing – review & editing. EA-G: Writing – original draft, Writing – review & editing. MG-G: Writing – original draft, Writing – review & editing.

## Funding

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

## Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

## Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

## Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpubh.2023.1296821/full#supplementary-material>



9. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, et al. Original contribution amount of time spent in sedentary Behaviors in the United States, 2003–2004. *Am J Epidemiol.* (2008) 167:875–81. doi: 10.1093/aje/kwm390
10. Chastin SFM, Buck C, Freiburger E, Murphy M, Brug J, Cardon G. Systematic literature review of determinants of sedentary behaviour in older adults: a DEDIPAC study. *Int J Behav Nutr Phys Act.* (2015) 12:127. doi: 10.1186/s12966-015-0292-3
11. Sprod J, Ferrar K, Olds T, Maher C. Changes in sedentary behaviours across the retirement transition: a systematic review. *Age Ageing.* (2015) 44:918–25. doi: 10.1093/ageing/afv140
12. Leask CF, Harvey JA, Skelton DA, Chastin SFM. Exploring the context of sedentary behaviour in older adults (what, where, why, when and with whom). *Eur Rev Aging Phys Act.* (2015) 12:1–8. doi: 10.1186/s11556-015-0146-7
13. Semanik PA, Lee J, Song J, Chang RW, Sohn MW, Ehrlich-Jones LS, et al. Accelerometer-monitored sedentary behavior and observed physical function loss. *Am J Public Health.* (2015) 105:560–6. doi: 10.2105/AJPH.2014.302270
14. Giné-Garriga M, Sansano-Nadal O, Tully MA, Caserotti P, Coll-Planas L, Rothenbacher D, et al. Accelerometer-measured sedentary and physical activity time and their correlates in European older adults: the SITLESS study. *J Gerontol A Biol Sci Med Sci.* (2020) 75:1754–62. doi: 10.1093/gerona/glaa016
15. Sasaki JE, John D, Freedson PS. Validation and comparison of ActiGraph activity monitors. *J Sci Med Sport.* (2011) 14:411–6. doi: 10.1016/j.jsams.2011.04.003
16. Steeves JA, Bowles HR, McClain JJ, Dodd KW, Brychta RJ, Wang J, et al. Ability of thigh-worn ActiGraph and activPAL monitors to classify posture and motion HHS public access. *Med Sci Sports Exerc.* (2015) 47:952–9. doi: 10.1249/MSS.0000000000000497
17. Chastin SFM, Dontje ML, Skelton DA, Čukić I, Shaw RJ, Gill JMR, et al. Systematic comparative validation of self-report measures of sedentary time against an objective measure of postural sitting (activPAL). *Int J Behav Nutr Phys Act.* (2018) 15:1–12. doi: 10.1186/s12966-018-0652-x
18. Hedayatrad L, Stewart T, Duncan S. Concurrent validity of ActiGraph GT3X+ and Axivity AX3 accelerometers for estimating physical activity and sedentary behavior. *J Meas Phys Behav.* (2020) 4:1–8.
19. Ainsworth B, Cahalin L, Buman M, Ross R. The current state of physical activity assessment tools. *Prog Cardiovasc Dis.* (2015) 57:387–95. doi: 10.1016/j.pcad.2014.10.005
20. Urda JL, Larouere B, Verba SD, Lynn JS. Comparison of subjective and objective measures of office workers' sedentary time. *Prev Med Rep.* (2017) 8:163–8. doi: 10.1016/j.pmedr.2017.10.004
21. Sansano-Nadal O, Wilson JJ, Martín-Borrás C, Brønd JC, Skjødt M, Caserotti P, et al. Validity of the sedentary behavior questionnaire in European older adults using English, Spanish. *German Danish Versions Meas Phys Educ Exerc.* (2021) 26:1–14. doi: 10.1080/1091367X.2021.1922910
22. Wilmot EG, Edwardson CL, Achana FA, Davies MJ, Gorely T, Gray LJ, et al. Sedentary time in adults and the association with diabetes, cardiovascular disease and death: systematic review and meta-analysis. *Diabetologia.* (2012) 55:2895–05. doi: 10.1007/s00125-012-2677-z
23. Werneck AO, Oyeyemi AL, Szwarcwald CL, Vancampfort D, Silva DR. Associations between TV viewing and depressive symptoms among 60,202 Brazilian adults: the Brazilian national health survey. *J Affect Disord.* (2018) 236:23–30. doi: 10.1016/j.jad.2018.04.083
24. Hallgren M, Dunstan DW, Owen N. Passive versus mentally active sedentary Behaviors and depression. *Exerc Sport Sci Rev.* (2020) 48:20–7. doi: 10.1249/JES.0000000000000211
25. Nicolson G, Hayes C, Darker C. Examining total and domain-specific sedentary behaviour using the socio-ecological model - a cross-sectional study of Irish adults. *BMC Public Health.* (2019) 19:1–13. doi: 10.1186/s12889-019-7447-0
26. Shaw RJ, Cukic I, Deary IJ, Gale CR, Chastin SFM, Dall PM, et al. Relationships between socioeconomic position and objectively measured sedentary behaviour in older adults in three prospective cohorts. *BMJ Open.* (2017) 7:1–10. doi: 10.1136/bmjopen-2017-016436
27. Van Der Berg JD, Bosma H, Caserotti P, Eiriksdottir G, Arnardottir NY, Martin KR, et al. Midlife determinants associated with sedentary behavior in old age. *Med Sci Sport Exerc.* (2014) 46:1359–65. doi: 10.1249/MSS.0000000000000246
28. Phelan JC, Link BG, Tehranifar P. Social conditions as fundamental causes of health inequalities: theory, evidence, and policy implications. *J Health Soc Behav.* (2010) 51:S28–40. doi: 10.1177/0022146510383498
29. Barro RJ, Lee JW. A new data set of educational attainment in the world, 1950–2010. *J Dev Econ.* (2013) 104:184–98. doi: 10.1016/j.jdeveco.2012.10.001
30. Lynch JW, Kaplan GA, Salonen JT. Why do poor people behave poorly? Variation in adult health behaviours and psychosocial characteristics by stages of the socioeconomic lifecourse. *Soc Sci Med.* (1997) 44:809–19. doi: 10.1016/S0277-9536(96)00191-8
31. Martikainen P, Brunner E, Marmot M. Socioeconomic differences in dietary patterns among middle-aged men and women. *Soc Sci Med.* (2003) 56:1397–10. doi: 10.1016/S0277-9536(02)00137-5
32. De la Cruz-Sánchez E, Feu S, Vizuete-Carrizosa M. El nivel educativo como factor asociado al bienestar subjetivo y la salud mental en la población española. *Univ Psychol.* (2012) 12:31–40. doi: 10.11144/javeriana.upsy12-1.efab
33. Lakerveld J, Luyen A, Schotman N, Peeters CFW, Cardon G, van der Ploeg HP, et al. Sitting too much: a hierarchy of socio-demographic correlates. *Prev Med.* (2017) 101:77–83. doi: 10.1016/j.ypmed.2017.05.015
34. Prince SA, Reed JL, McFetridge C, Tremblay MS, Reid RD. Correlates of sedentary behaviour in adults: a systematic review. *Obes Rev.* (2017) 18:915–35. doi: 10.1111/obr.12529
35. Palmer VJ, Gray CM, Fitzsimons CF, Mutrie N, Wyke S, Deary IJ, et al. What do older people do when sitting and why? Implications for decreasing sedentary behavior. *Gerontologist.* (2019) 59:686–97. doi: 10.1093/geront/gny020
36. Dunlop DD, Song J, Arntson EK, Semanik PA, Lee J, Chang RW, et al. Sedentary time in US older adults associated with disability in activities of daily living independent of physical activity. *J Phys Act Health.* (2015) 12:93–01. doi: 10.1123/jpah.2013-0311
37. Belletiere J, Carlson JA, Rosenberg D, Singhanian A, Natarajan L, Berardi V, et al. Gender and age differences in hourly and daily patterns of sedentary time in older adults living in retirement communities. *PLoS One.* (2015) 10:e0136161. doi: 10.1371/journal.pone.0136161
38. Loginov SI, Malkov MN, Nikolayev AY. Gender-specific manifestations of daily physical activity and sedentary behavior in elderly residents of Surgut. *Adv Gerontol.* (2017) 30:573–8.
39. U.S. Department of Labor Bureau of Labor. American time use survey—2013 results (2014).
40. Brug J, Chinapaw M. Determinants of engaging in sedentary behavior across the lifespan; lessons learned from two systematic reviews conducted within DEDIPAC. *Int J Behav Nutr Phys Act.* (2015) 12:10–3. doi: 10.1186/s12966-015-0293-2
41. Guralnik JM, Ferrucci L, Pieper CF, Leveille SG, Markides KS, Ostir GV, et al. Lower extremity function and subsequent disability: consistency across studies, predictive models, and value of gait speed alone compared with the short physical performance battery. *J Gerontol A Biol Sci Med Sci.* (2000) 55:M221–31. doi: 10.1093/gerona/55.4.m221
42. Ku PW, Steptoe A, Liao Y, Hsueh MC, Chen LJ. Clinical medicine a threshold of objectively-assessed daily sedentary time for all-cause mortality in older adults: a meta-regression of prospective cohort studies. *J Clin Med.* (2019) 8:564. doi: 10.3390/jcm8040564
43. Giné-Garriga M, Coll-Planas L, Guerra M, Domingo À, Roqué M, Caserotti P, et al. The SITLESS project: exercise referral schemes enhanced by self-management strategies to battle sedentary behaviour in older adults: study protocol for a randomised controlled trial. *Trials.* (2017) 18:1–14. doi: 10.1186/s13063-017-1956-x
44. Migueles JH, Cadenas-Sanchez C, Ekelund U, Nyström CD, Mora-Gonzalez J, Lóf M, et al. Accelerometer data collection and processing criteria to assess physical activity and other outcomes: a systematic review and practical considerations Europe PMC funders group. *Sports Med.* (2017) 47:1821–45. doi: 10.1007/s40279-017-0716-0
45. Choi L, Liu Z, Matthews CE, Buchowski MS. Validation of accelerometer wear and nonwear time classification algorithm. *Med Sci in Sports Exerc.* (2011) 43:357–64. doi: 10.1249/MSS.0b013e3181ed61a3
46. Hirshkowitz M, Whitton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National sleep foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health.* (2015) 1:40–3. doi: 10.1016/j.sleh.2014.12.010
47. Pate RR, Stevens J, Webber LS, Dowda M, Murray DM, Young DR, et al. Age-related change in physical activity in adolescent girls. *J Adolesc Health.* (2009) 44:275–82. doi: 10.1016/j.jadohealth.2008.07.003
48. Byrom B, David R. Measuring free-living physicalactivity in COPD patients: deriving methodology standards for clinical trials through a review of research studies. *Contemp Clin Trials.* (2016) 2016:172–84. doi: 10.1016/j.cct.2016.01.006
49. Parker EB. Information and society In: CA Cuadra and MJ Bates, editors. *Library and information service needs of the nation: Proceedings of a conference on the needs of occupational, ethnic and other groups in the United States.* Washington, DC: U.S.G.P.O (1974)
50. Bates MJ. An introduction to metatheories, theories, and models In: KE Fisher, S Erdelez and L McKechnie, editors. *Theories of information behavior.* Medford, NJ: Information Today (2005a)
51. Healy GN, Clark BK, Winkler EA, Gardiner PA, Brown WK, Matthews CE. (2011). Measurement of adults' sedentary time in population-based studies. *Am J Prev Med.* (2011) 41:216–27. doi: 10.1016/j.amepre.2011.05.005
52. Prince SA, Cardilli L, Reed JL, Saunders TJ, Kite C, Douillette K, et al. A comparison of self-reported and device measured sedentary behaviour in adults: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* (2020) 17:31. doi: 10.1186/s12966-020-00938-3
53. Bakker EA, Hartman YAW, Hopman MTE, Hopkins ND, Graves LEF, Dunstan DW, et al. Validity and reliability of subjective methods to assess sedentary behaviour in adults: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* (2020) 17:75. doi: 10.1186/s12966-020-00972-1

54. Burton NW, Haynes M, Van Uffelen JGZ, Brown WJ, Turrell G. Mid-aged adults' sitting time in three contexts. *Am J Prev Med.* (2012) 42:363–73. doi: 10.1016/j.amepre.2011.11.012
55. Stamatakis E, Hillsdon M, Mishra G, Hamer M, Marmot M. Television viewing and other screen-based entertainment in relation to multiple socioeconomic status indicators and area deprivation: the Scottish health survey 2003. *J Epidemiol Community Health.* (2009) 63:734–40. doi: 10.1136/jech.2008.085902
56. Van Cauwenberg J, Van Holle V, De Bourdeaudhuij I, Owen N, Deforche B. Diurnal patterns and correlates of older adults' sedentary behavior. *PLoS One.* (2015) 10:e0133175. doi: 10.1371/journal.pone.0133175
57. Van Dyck D, Cardon G, De Bourdeaudhuij I. Longitudinal changes in physical activity and sedentary time in adults around retirement age: what is the moderating role of retirement status, gender and educational level? *BMC Public Health.* (2016) 16:1–11. doi: 10.1186/s12889-016-3792-4
58. Prince SA, Roberts KC, Melvin A, Butler GP, Thompson W. Gender and education differences in sedentary behaviour in Canada: an analysis of national cross-sectional surveys. *BMC Public Health.* (2020) 20:1–17. doi: 10.1186/s12889-020-09234-y
59. Jakes RW, Day NE, Khaw KT, Luben R, Oakes S, Welch A, et al. Television viewing and low participation in vigorous recreation are independently associated with obesity and markers of cardiovascular disease risk: EPIC-Norfolk population-based study. *Eur J Clin Nutr.* (2003) 57:1089–96. doi: 10.1038/sj.ejcn.1601648
60. Kronenberg F, Pereira MA, Schmitz MK, Arnett DK, Evenson KR, Crapo RO, et al. Influence of leisure time physical activity and television watching on atherosclerosis risk factors in the NHLBI family heart study. *Atherosclerosis.* (2000) 153:433–43. doi: 10.1016/S0021-9150(00)00426-3
61. Johnson W, Kyvik KO, Skytthe A, Deary IJ, Sørensen T. Education modifies genetic and environmental influences on BMI. *PLoS One.* (2011) 6:16290. doi: 10.1371/journal.pone.0016290
62. Herzog AR, Franks MM, Markus HR, Holmberg D. Activities and well-being in older age: effects of self-concept and educational attainment. *Psychol Aging.* (1998) 13:179–85. doi: 10.1037/0882-7974.13.2.179
63. Molina JA, Campaña JC, Ortega R. What do you prefer for a relaxing time at home: reading, watching TV or listening to the radio? *Appl Econ Lett.* (2016) 23:1278–84. doi: 10.1080/13504851.2016.1150943
64. U.S. Department of Labor. American time use - 2018 results. (2019). Available at: <https://www.bls.gov/news.release/pdf/atus.pdf> (accessed 17 September 2019).
65. Piepiora PA, Maciej R. *Hypothesis on the supreme value criteria of the global civilization. Healthcare and medical devices.* San Francisco (USA): AHFE International (2023). 280–89.
66. Darmon N, Drewnowski A. Does social class predict diet quality? *In Am J Clin Nutr.* (2008) 87:1107–17. doi: 10.1093/ajcn/87.5.1107
67. Aguiar-Bloemer AC, Palazzo CC, Diez-Garcia RW. Relationship of negative emotion with leptin and food intake among overweight women. *Physiol Behav.* (2021) 237:113457. doi: 10.1016/j.physbeh.2021.113457
68. Shields M, Tremblay MS. Sedentary behavior and obesity. *Health Rep.* (2008) 19:19–30.
69. Dogra S, Ashe MC, Biddle SJH, Brown WJ, Buman MP, Chastin S, et al. Sedentary time in older men and women: an international consensus statement and research priorities. *Br J Sports Med.* (2017) 51:1526–32. doi: 10.1136/bjsports-2016-097209