



Predictors of Posttraumatic Stress Disorder following the 2023 Earthquakes in Turkey.

Han Alpay , E., Redican, E., Hyland, P., Karatzias, T., & Shevlin, M. (in press). Predictors of Posttraumatic Stress Disorder following the 2023 Earthquakes in Turkey. *Journal of Loss and Trauma*.

[Link to publication record in Ulster University Research Portal](#)

Published in:

Journal of Loss and Trauma

Publication Status:

Accepted/In press: 23/01/2024

Document Version

Author Accepted version

General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

Earthquake-related traumatic events and PTSD
Predictors of Posttraumatic Stress Disorder following the 2023 Earthquakes in Turkey.

Authors:

Emre Han Alpay¹

Enya Redican²

Philip Hyland³

Thanos Karatzias⁴

Mark Shevlin²

Affiliations

¹ Department of Psychology, Mersin University, Mersin, Türkiye

² Ulster University, School of Psychology, Coleraine, UK

³ Department of Psychology, Maynooth University, Ireland.

⁴ Edinburgh Napier University, School of Health and Social Care, Edinburgh, UK.

Corresponding Author

Name: Mark Shevlin.

Postal address: Ulster University, School of Psychology, Cromore Road, Coleraine, Co.

Londonderry, UK, BT52 1SA

Email: m.shevlin@ulster.ac.uk, Telephone: +44 28 7012 3141

Data availability statement: The data are not available.

Declaration of interest: The authors report no financial or other relationship relevant to the subject of this article.

Competing interests' statement: Authors have no competing interests to declare.

Acknowledgements: We would like to thank the NGO *Kilikya- Nehir Sosyal Dayanışma ve Kültür Derneđ*.

Abstract

On February 6, 2023, an earthquake registering 7.8 on the Richter scale struck southeast Turkey, and this was followed by a second earthquake of magnitude 7.5. The earthquake resulted in widespread loss, devastation, and destruction, and is anticipated to bear a significant psychological toll on survivors. This study aimed to investigate the prevalence and correlates of ICD-11 posttraumatic stress disorder (PTSD) in those living in earthquake-affected areas of Turkey. The sample was a convenience sample of 527 adults living in earthquake-affected areas of Turkey. Prevalence rates and gender differences in exposure to earthquake-related traumatic events were investigated. Moreover, prevalence rates of ICD-11 PTSD were computed, and demographic and trauma-related correlates of ICD-11 PTSD were assessed using logistic regression analyses. Most of the sample (99.2%; n= 523) experienced at least one earthquake-related traumatic event, with the majority experience multiple events. Findings demonstrated high endorsement of the ICD-11 PTSD symptom clusters, with 54.1% meeting criteria for diagnosis of probable ICD-11 PTSD. Correlates associated with increased risk of PTSD include female gender, physical injury in the earthquake, physical injury of family or loved ones in the earthquake and being in the earthquake-related traumatic events exposure quartile. These findings indicate high levels of trauma exposure and PTSD among survivors of the recent earthquakes which occurred in Turkey. The identification of correlates associated with increased risk of PTSD is useful for identifying Turkish earthquake survivors who are most likely to need psychological support.

Keywords: PTSD; natural disaster; earthquake; Turkey; trauma

A devastating 7.8 magnitude earthquake struck southeast Turkey and the northwest of Syria on February 6, 2023. A number of aftershocks, including one with a magnitude similar to the first earthquake, also occurred. According to estimates, the earthquakes have resulted in more than 50,000 fatalities, tens of thousands of injuries, and the internal displacement of more than 216,000 people (United Nations Population Fund (UNFPA, 2023)). It has also rendered nearly 1.5 million people temporarily homeless due to severe infrastructural damage (United Nations (UN), 2023). Accordingly, the earthquakes are projected to have significant long-term effects (Duzova et al., 2023), including having a detrimental impact on the mental health of the Turkish people (Garfin & Silver, 2023).

It is widely established that natural disasters commonly result in psychological distress in addition to the onset or exacerbation of psychiatric disorders among exposed populations including PTSD, depression, anxiety, substance abuse, grief reactions, and sleep disturbances (Agyapong et al., 2022; Beaglehole et al., 2018; Golitaleb et al., 2022; Neria et al., 2016; Morganstein & Ursano, 2020; Saeed et al., 2022; Zhang et al., 2022). Given that disasters possess all the defining characteristics of large-scale traumatic events (Goldmann & Galea, 2014), it is unsurprising that one of the most prevalent psychological disorders observed among individuals exposed to natural disasters is posttraumatic stress disorder (PTSD) (Beaglehole et al., 2018; Goldmann & Galea, 2014; Hosseinnejad et al., 2021; North, 2016; Neria et al., 2016; Rafiey et al., 2019). A meta-analytic study reported prevalence rates ranging from 1.20 to 82.64% for PTSD among earthquake survivors, with the combined incidence of PTSD within nine months of an earthquake being 28.76% and after nine months being 19.48% (Dai et al., 2016). Explanations for the significant heterogeneity in rates of PTSD across studies have included variations in earthquake intensities, the degree to which

victims were exposed to the earthquake, time of assessment, demographic and cultural backgrounds of samples, type of property loss, and whether a bereavement occurred (Dai et al., 2016; Tang et al., 2017). Another meta-analytic study by Hosseinnejad et al. (2022) that examined the prevalence of PTSD following earthquakes in Iran and Pakistan found that the rates of PTSD for Iranian and Pakistani survivors were 60.2% and 49.2%, respectively. A shorter time lapse, sex, and method of assessment were found to explain variation in prevalence estimates. Hence, given the widespread variation in post-earthquake PTSD prevalence rates, extrapolating findings from other disaster-exposed populations is unlikely to provide an accurate depiction of the extent of PTSD among the Turkish population.

Given the high rates of PTSD in the aftermath of an earthquake, it is important to determine the factors that may contribute to greater risk. One such factor which has shown to play an important role in elevating the risk of developing PTSD is exposure to earthquake-related traumatic events (e.g., Baral et al., 2019; Tang et al., 2017; Tarikulu-Tanrikulu et al., 2023; Liang et al., 2019; Pino et al., 2021). For instance, research has shown significant correlates of PTSD for earthquake victims to include being trapped, experiencing fear, injury, or bereavement (Baral et al., 2019; Dai et al., 2016; Tang et al., 2017), as well as experiencing property damage or witnessing death (Dai et al., 2016; Pino et al., 2021). Additionally, research has shown a dose-response association between exposure to earthquake-related traumatic events and symptoms of PTSD (Naeem et al., 2011). Overall, the extant evidence base indicates that exposure to earthquake-related traumatic events is a robust predictor of PTSD in earthquake survivors and that specific types of experiences are more likely to infer greater risk of PTSD.

Research pertaining to the epidemiology of PTSD among earthquake-exposed populations have relied on DSM-based definitions of the condition. However, in the most recent version of the *International Classification of Diseases* (ICD-11; World Health

Organization [WHO], 2018), a revised formulation of PTSD was included. The ICD-11 describes PTSD as being comprised of three symptom clusters including (1) re-experiencing of the trauma in the here and now (Re), (2) avoidance of traumatic reminders (Av), and (3) sense of current threat (Th). Additionally, symptoms must be present several weeks after the traumatic event and cause significant impairment across multiple functional domains (WHO, 2018). ICD-11 PTSD symptoms have been explored in a range of populations such as those who have experienced war (Karatzias et al., 2023), clinical samples (Karatzias et al., 2019), and refugees (Vallières et al., 2018), but not yet in a natural disaster exposed population. This is particularly important given that studies have shown how significantly fewer individuals meet requirements for diagnosis of ICD-11 PTSD as compared to DSM-5 PTSD (e.g., Boelen, 2023; Heeke et al., 2020; Hyland et al., 2018; Shevlin et al., 2018).

Consequently, the current study sought to explore rates and correlates of ICD-11 PTSD symptoms in individuals who were exposed to the recent earthquakes in Turkey. It is critical to identify the prevalence and correlates of ICD-11 PTSD symptoms in order to develop interventions and mental health preparedness procedures that may alleviate the symptoms of PTSD (Lee et al., 2020). Based on existing literature (e.g., Dai et al., 2016; Hosseinnejad et al., 2016; Pino et al., 2021; Tarikulu et al., 2023), it was anticipated that rates and overall levels of PTSD symptoms would be high. Given that this was the first study to assess for ICD-11 PTSD, and prior research has shown substantial differences in prevalence rates according to the ICD and DSM-5 definitions of PTSD (e.g., Shevlin et al., 2018), it is important to determine the prevalence of ICD-11 PTSD in a post natural disaster context. It was expected that earthquake-related traumatic events relating to experiences of entrapment, fear, injury, bereavement, property damage, and witnessing death would be linked to higher PTSD symptom levels (Baral et al., 2019; Dai et al., 2016; Tang et al., 2017;). Additionally, it

was anticipated that rates of PTSD symptoms would be highest among those who had experienced higher levels of earthquake-related traumatic events.

Methods

Participants and procedure

The sample was a convenience sample of adults living in earthquake-affected areas of Turkey. Specifically, following the earthquake, more than 500,000 Turkish citizens moved to the city of Mersin where the government provided housing in student dormitories, empty flats, guesthouses, and hotels. Data was collected during the voluntary aid process and most data was collected from Kilikya-Nehir Social Solidarity and Culture Association; a non-governmental organisation (NGO) based in Mersin. Most of the data was collected from the accommodation center of this NGO which hosted more than 400 earthquake victims.

Inclusion criteria for participation in the study included that participants were literate, did not have any psychotic illness, and had been affected by the earthquake. On March 21, 2023 (about 45 days after the earthquakes), data collecting started and concluded on April 22, 2023. Because most of the earthquake victims had no smartphone or computer, data was collected using pen and paper. The final sample comprised of 527 participants, of which the average age was 31.69 years (SD = 13.80, Median = 26.00, Range = 18 to 75 years). Females comprised more than two-thirds of the sample (65.3%; n = 344), while more than half of the sample were not in a romantic relationship (63.0%; n = 338). Almost half of the sample reported university as their highest educational attainment (46.7%; n = 246). Ethical approval for the current study was provided by the ethics committee of *blinded for review*.

Measures

Demographic variables: Demographic variables include gender (Female = 0, Male = 1) and age (measured in years).

Earthquake-related traumatic events: The Traumatic Exposure Severity Scale (TESS; Elal & Slade, 2005) is a self-report measure of exposure to traumatic events among disaster survivors. The original Traumatic Exposure Severity Scale is comprised of 24-items, however, for the purposes of the present study only ten of those items were included. Ten items which comprehensively captured the various types of experiences which can occur following exposure to an earthquake were included to avoid burdening participants. Hence, the current study used an abbreviated version of the Traumatic Exposure Severity Scale (TESS-A; Elal & Slade, 2005). A list of those items is presented in Table 1. For the current study, individual items were investigated as opposed to total scores.

ICD-11 PTSD: The International Trauma Questionnaire (ITQ; Cloitre et al., 2018) is a self-report measure of ICD-11 PTSD symptoms. The ITQ first screens for an individual's 'index' traumatic event and participants are instructed to answer all questions in reference to this event. In the present study, participants were instructed to answer all questions in reference to the earthquake they had experienced. The ITQ consists of 12 items, of which six evaluate the three PTSD symptom clusters (Re-experiencing, Avoidance, Threat) and six evaluate the extra disturbances in self-organisation (DSO) symptom clusters that are a component of the Complex PTSD (CPTSD) diagnosis. For the present study, only the PTSD module of the ITQ items were used because of the assessment of symptoms in relation to the recent earthquake experience. Three additional items are included which assess for functional impairment across numerous domains including social, occupation, and other important areas of life. Participants indicate how bothered they have been by the symptoms over the past month. All ITQ items are rated on a 5-point Likert scale ranging from 0 ('Not at all') to 4 ('Extremely'). The ITQ can be used to capture symptom severity and to determine if diagnostic requirements are met. The severity score method involves summing responses to all PTSD items, generating possible scores ranging from 0 to 24. Diagnostic requirements for

Earthquake-related traumatic events and PTSD

PTSD include the presence of one of two items from each of the PTSD symptom clusters and at least one functional impairment item must be positively endorsed (endorsement is based on a Likert score of ≥ 2 (i.e., 'Moderately')). The internal reliability of the PTSD ($\alpha = .83$) scores in this sample was excellent.

Data Analysis

First, descriptive statistics were calculated to determine endorsement rates for each earthquake-related traumatic events, and gender differences were tested using the Pearson chi-square (χ^2) test. The strength of the association was quantified using Cramer's V ($\leq 0.2 =$ weak, 0.2 to $0.6 =$ moderate, > 0.6 strong). Second, rates of endorsement for the PTSD items and symptom clusters, as well as the prevalence rates of ICD-11 PTSD and differences in prevalence rates according to gender and age were examined. Moreover, descriptive statistics were calculated for the PTSD items as well as overall PTSD score. Third, both a bivariate and multivariate binary logistic regression analysis were conducted to determine the unadjusted and adjusted associations between 12 correlates (i.e., gender, age, and the ten earthquake-related traumatic events) and meeting diagnostic requirements for ICD-11 PTSD. To examine the cumulative effect of exposure to multiple earthquake-related traumatic events controlling for gender and age, the Traumatic Exposure Severity Scale -A scores were summed and transformed into a quartile variable with the lowest quartile used as the reference category. Transforming the summed score into a quartile variable allowed for an examination of how different quantities of traumatic exposure inferred greater risk of PTSD. Fourth, a bivariate and multivariate regression analysis was conducted to examine the association between correlates and total PTSD scores. These analyses provided both unstandardized (B) regression coefficients and standardized regression coefficients (β). All analyses were conducted using SPSS Version 28.

Results

Earthquake-related traumatic events and PTSD

Most of the sample (99.2%; $n = 523$) experienced at least one earthquake-related traumatic event. Of those trauma-exposed participants, 4.6% ($n = 24$) reported experiencing one earthquake-related traumatic event, 6.3% ($n = 33$) reported exposure to two, 8.4% ($n = 44$) reported exposure to three, and 80.7% ($n = 422$) reported exposure to four or more. The average number of earthquake-related traumatic events was 5.43 ($SD = 2.30$, Range = 0 – 10). The most commonly endorsed earthquake-related traumatic events were *'In the days following the earthquake did you have to spend the night somewhere other than in your home'* (93.4%; $n = 492$), *'Did you need food and water aid after the earthquake'* (84.3%; $n = 444$), and *'Was your home damaged in the earthquake'* (82.4%; $n = 434$).

Females were significantly more likely to endorse the items *'Did you need food and water aid after the earthquake'* (Male = 30.9%, Female = 60.9%; $\chi^2(1) = 4.91$, $p = .03$, $V = .097$) while males were significantly more likely to endorse the items *'Were you involved in the rescue work?'* (Male = 13.1%, Female = 6.8%; $\chi^2(1) = 55.55$, $p < .001$, $V = .33$) and *'Did you see any dead bodies or body parts during the rescue and clearing up work period?'* (Male = 15.7%, Female = 15.4%; $\chi^2(1) = 26.51$, $p < .001$, $V = .22$).

Item-level descriptives for the PTSD items are included in Table 2. The mean PTSD score for the sample was 13.13 ($SD = 5.76$), with females having significantly higher scores ($M = 14.01$, $SD = 5.50$) than males ($M = 11.47$, $SD = 5.88$), $t(525) = -4.94$, $p < .001$, Cohen's $d = 0.45$. The ITQ items with the highest rates of endorsement were *'Being "super-alert", watchful, or on guard'* (76.4%, $n = 403$), *'Having powerful images or memories that sometimes come into your mind in which you feel the experience is happening again in the here and now'* (77.0%, $n = 406$), and *'Feeling jumpy or easily startled'* (75.0%, $n = 395$). The ITQ symptom cluster with the highest endorsement rate was sense of current threat (83.5%, $n = 440$), followed by re-experiencing (82.5%, $n = 435$), and avoidance (74.0%, $n = 390$). Overall, 54.1% (95% CI = 49.8%, 58.4%) met diagnostic requirements for ICD-11 PTSD.

Earthquake-related traumatic events and PTSD

There was a significant gender difference in rates of PTSD (Males = 14.4%, Females = 39.7%; $\chi^2(1) = 17.78, p < .001, V = .184$). There was no age difference in likelihood of meeting criteria for PTSD (Criteria met: $M = 32.12 [SD = 14.14]$ vs. Criteria not met: $M = 31.18 [SD = 13.39]; t(525) = -.77, p = .439, d = .07$).

The unadjusted and adjusted associations between the correlates and meeting criteria for ICD-11 PTSD are presented in Table 3. In the unadjusted models, meeting criteria for ICD-11 PTSD was associated with being physically injured in the earthquake ($OR = 2.54$), family or loved ones being physically injured in the earthquake ($OR = 2.23$), losing family in the earthquake ($OR = 1.62$) and female gender ($OR = 2.65$). The adjusted logistic regression model was statistically significant ($\chi^2(12) = 59.28, p < .001$). In the adjusted model, female gender ($AOR = 2.65$), being physically injured in the earthquake ($AOR = 2.26$) and having a family member or loved ones being physically injured in the earthquake ($AOR = 1.72$) remained significantly associated with meeting criteria for PTSD.

Table 4 reports the association between the earthquake-related traumatic events quartile and meeting criteria for ICD-11 PTSD, controlling for gender and age. The adjusted logistic regression model was statistically significant ($\chi^2(5) = 52.07, p < .001$). Those in the highest quartile were significantly more likely to meet criteria for PTSD ($AOR = 3.54$).

Finally, the unadjusted and adjusted associations between the correlates and total PTSD scores are presented in Table 5. In the unadjusted models, significant positive correlates of PTSD symptom levels included damage to home in the earthquakes, being physically injured in the earthquakes, family or loved ones being physically injured in the earthquakes, losing family in the earthquakes, losing relatives in the earthquakes, seeing dead bodies or body parts during the rescue, and clearing up work period and female gender. In the adjusted models, significant positive correlates included being physically injured in the

earthquakes and family or loved ones being physically injured in the earthquakes and female gender. Table 6 reports the association between the earthquake-related traumatic events quartile and total PTSD symptom levels, controlling for gender and age. The overall adjusted model was statistically significant, $R^2 = .14$, $F(12, 513) = 6.79$, $p < .001$. Those in the third quartile and those in the highest quartile had significantly higher levels of PTSD symptoms.

Discussion

The primary goal of this study was to examine the occurrence and correlates of ICD-11 PTSD in a convenience sample of people living in the earthquake-exposed regions of Turkey shortly after the disaster. Almost everyone surveyed reported exposure to at least one earthquake-related traumatic events, with the majority reporting exposure to four or more events. The most commonly endorsed earthquake-related traumatic events related to both loss of resources and damage to one's home. Significant gender differences were observed across the different types of earthquake-related traumatic events with females being more likely to report requiring food and water aid after the earthquake and males being more likely to report involvement in rescue work and exposure to dead bodies during the clearing up process. These findings align with research demonstrating how the experience of catastrophic events is markedly different for males and females such that males typically play a significant role in the rescue phase while females often assume a caring role (Moreno et al., 2022). Overall, these findings offer a novel understanding of the severity of posttraumatic stress symptoms in the immediate wake of the earthquakes, as well as an overview of the characteristics of the survivors and the earthquakes that increased susceptibility to maladaptive posttraumatic stress reactions.

Earthquake-related traumatic events and PTSD

Consistent with the extant evidence base (e.g., Dai et al., 2016; Hosseinejad et al., 2022; Tang et al., 2017), rates and overall levels of PTSD symptoms were extremely elevated in the current sample. The PTSD symptom cluster that was most pertinent to participants related to hyperarousal. This is unsurprising given that increased occurrence and frequency of earthquakes are expected in the affected regions for many years to come (Dal Zilio & Ampuero, 2023). The endorsement rates of all PTSD symptom clusters in the present study were substantially higher than those observed in studies conducted on other earthquake-exposed populations. For instance, Lai et al. (2004) found that in their sample of survivors of the 1999 Taiwanese earthquake (7.3 on Richter scale) that rates of endorsement of intrusion symptoms was 61.5%, avoidance/numbing symptoms was 12.3%, and hyperarousal symptoms was 48.4%. Another study conducted by Livanou et al. (2002) on survivors of the previous Turkish earthquake (7.4 on Richter scale) also reported lower endorsement rates of the symptom clusters. The higher endorsement of these symptom clusters in the present study is likely attributable to the greater magnitude of the recent Turkish earthquakes, with research demonstrating magnitude of exposure to a natural disaster to represent a key risk factor of PTSD (Neria et al., 2008). Hence, this study provides initial evidence for a high burden of PTSD symptoms among survivors of the recent Turkish earthquakes, indicating that strategies for the assessment and treatment of PTSD should be implemented.

Overall, findings demonstrated that just over half of the sample met criteria for diagnosis of probable ICD-11 PTSD. The prevalence of PTSD in the present study is notably higher than the pooled estimates reported in a meta-analytic study where 28.76% of earthquake survivors met criteria for diagnosis of probable PTSD (based on DSM definitions) (Dai et al., 2016), yet comparable to rates observed following the Iranian and Pakistan. The prevalence of PTSD observed in the present study is also notably higher than those reported for the previous Turkish earthquake where prevalence rates have been found to range from

Earthquake-related traumatic events and PTSD

19.2% to 23% (Önder et al., 2006; Başoğlu et al., 2004). Again, it is possible that the greater magnitude of the most recent earthquake accounts for the higher prevalence of PTSD, supporting predictions that the Turkish earthquakes are likely to result in increased levels of trauma and PTSD as compared to the previous earthquake (Iqbal & Sheikh, 2023). Alternatively, assessment of PTSD in the current study was conducted soon after the exposure to the earthquakes whereas studies of prior Turkish earthquakes were conducted fourteen months (Başoğlu et al., 2004) and thirty-six months (Önder et al., 2006) following the earthquake. Research has shown how PTSD symptoms decrease during the first two years post-earthquake (Hosseinnejad et al., 2022; Karamustafalıoğlu et al., 2023), and hence, further research is required to capture changes in the prevalence of PTSD among Turkish earthquake survivors over time.

Consistent with prior research conducted on earthquake-exposed populations (Tang et al., 2017) in addition to the broader PTSD literature (e.g., McGinty et al., 2021), rates of PTSD were highest among female survivors of the earthquakes. Numerous explanations have been proposed for the greater incidence of PTSD among females including that females are often exposed to traumas that are more likely to evoke a posttraumatic stress response (Tolin & Foa, 2006), as well as genetic predisposition and hormonal influences and fluctuations (Christiansen & Berke, 2020). However, when discussing these gender effects, it is important to mention that over two-thirds of the sample were female. Although (as highlighted) females have been frequently identified as being at greatest risk of PTSD, the poor representativity of males within the current study may preclude an accurate representation of the extent of PTSD among males in post-earthquake Turkey. Moreover, it may be a contributing factor to the high rates of PTSD in the current sample.

The final aim of the present study was to explore correlates of ICD-11 PTSD among the earthquake-exposed sample. First, associations with demographic (i.e., gender, age) and

earthquake-related traumatic events were examined. Findings from the bivariate and multivariate analyses indicated that female gender was a significant correlate of PTSD, with possible explanations for this previously highlighted. Consistent with our initial hypothesis and previous research (Baral et al., 2019; Dai et al., 2016; Tang et al., 2017), physical injury of the self as well as family or loved ones were identified as significant correlates of PTSD in both the bivariate and multivariate analyses. Notably, although the loss of family members and relatives in the earthquakes were significant correlates of PTSD in the bivariate analyses, these associations reached a level of statistical non-significance in the multivariate analyses. It is likely that the shared variance between these earthquake-related traumatic events and PTSD is fully accounted for by the other traumatic events included in the analysis. There are numerous ways to examine the impact of earthquake-related trauma exposure on risk of PTSD; with many studies investigating exposure in an aggregated form (e.g., Jin et al., 2014; Wang et al., 2020) and the current study investigated individual trauma types. There are advantages associated with both approaches such that the investigation of individual earthquake-related events in the present study enabled the identification of those experiences most highly associated with risk of PTSD (Kessler et al., 2017), whereas examining cumulative exposure to multiple types of earthquake-related experiences can also be a highly robust correlate of subsequent PTSD (e.g., Karam et al., 2014). To ensure that both approaches were addressed, the association between demographics (i.e., gender, age) and earthquake-related traumatic events quartiles were also examined in the current study. Findings from the bivariate and multivariate analyses demonstrated how those in earthquake-related traumatic events quartile four were almost four times more likely to meet criteria for diagnosis of probable PTSD. This is an expected finding given the widely established dose-response association between trauma exposure and risk of PTSD (e.g., Cloitre et al., 2019; Kvedaraite et al., 2022). The analyses were all replicated for total PTSD symptom levels,

with findings being mostly consistent with those for diagnostic status. Hence, this demonstrates that even if an individual fails to meet the diagnostic criteria for PTSD, the correlates of interest are still associated with higher levels of symptom severity.

Notably, the current study was unable to account for prior traumatic experiences when examining the association between earthquake-related traumatic events and risk of PTSD. Research has demonstrated prior trauma exposure and pre-trauma psychopathology to represent significant risk factors for subsequent trauma-related psychopathology (Sayed et al., 2015). This has also been evidenced in studies looking at risk factors of PTSD among earthquake survivors, where it has been shown that prior trauma exposure played a significant role in the development and onset of PTSD (e.g., Ehring et al., 2011; Tang et al., 2017). Hence, it is not possible to discount the possibility that the increased risk of PTSD among particular participants may be driven by pre-existing traumas and psychopathologies. Moreover, Complex PTSD (CPTSD) typically develops in response to exposure to multiple, repeated, and prolonged interpersonal traumas from which escape is often difficult (Hyland et al., 2017). Hence, future research may wish to examine the association between poly-traumatisation (co-occurring traumatic events that are both earthquake and non-earthquake related) and risk of CPTSD.

Despite being the first study to examine the prevalence and correlates of PTSD among survivors of the recent Turkish earthquakes, findings from this study should be considered in light of several limitations. First, participants were recruited using convenience sampling methods and hence, the generalisability to the wider earthquake affected Turkish population is uncertain. Second, and as previously mentioned, the gender composition of the present sample comprised of females for the majority, and thus research is now needed using more representative samples. Third, and also as previously mentioned, only earthquake-related traumatic events were included in this study while PTSD was examined only with respect to

the earthquake. Fourth, given the recency of the earthquake, it was deemed that the examination of Complex PTSD (CPTSD) may be inappropriate. Research is necessary to determine the prevalence of CPTSD among survivors in the future. Finally, because the data collection took place shortly after the earthquakes, the questionnaires were designed to be quick and simple to complete. As a result, many potentially significant factors, including relationships, education, and mental history, were not examined.

Overall, the current study indicates high levels of PTSD among survivors of the recent Turkish earthquake. The identification of correlates associated with increased risk of PTSD symptom levels and diagnostic status such as female gender, physical injury in the earthquake, physical injury of family or loved ones in the earthquake and cumulative trauma exposure is useful for identifying Turkish earthquake survivors who are most likely to need psychological support following the earthquake.

References

- Adhikari Baral, I., & KC, B. (2019). Post traumatic stress disorder and coping strategies among adult survivors of earthquake, Nepal. *BMC Psychiatry, 19*(1), 1-8.
<https://doi.org/10.1186/s12888-019-2090-y>
- Agyapong, B., Shalaby, R., Eboreime, E., Obuobi-Donkor, G., Owusu, E., Adu, M. K., ... & Agyapong, V. I. (2022). Cumulative trauma from multiple natural disasters increases mental health burden on residents of Fort McMurray. *European Journal of Psychotraumatology, 13*(1), 2059999.

- Baçoğlu, M., Şalcioğlu, E., & Livanou, M. (2002). Traumatic stress responses in earthquake survivors in Turkey. *Journal of Traumatic Stress: Official Publication of The International Society for Traumatic Stress Studies*, 15(4), 269-276.
<https://doi.org/10.1023/A:1016241826589>
- Beaglehole, B., Mulder, R. T., Frampton, C. M., Boden, J. M., Newton-Howes, G., & Bell, C. J. (2018). Psychological distress and psychiatric disorder after natural disasters: systematic review and meta-analysis. *The British Journal of Psychiatry*, 213(6), 716-722. doi: 10.1192/bjp.2018.210.
- Boelen, P. A. (2023). Nature and prevalence of PTSD as per DSM-5 and ICD-11 after natural and unnatural deaths. *Journal of Affective Disorders Reports*, 14, 100664.
- Cénat, J. M., McIntee, S. E., & Blais-Rochette, C. (2020). Symptoms of posttraumatic stress disorder, depression, anxiety, and other mental health problems following the 2010 earthquake in Haiti: A systematic review and meta-analysis. *Journal of affective disorders*, 273, 55-85. <https://doi.org/10.1016/j.jad.2020.04.046>
- Christiansen, D. M., & Berke, E. T. (2020). Gender-and sex-based contributors to sex differences in PTSD. *Current Psychiatry Reports*, 22, 1-9.
<https://doi.org/10.1007/s11920-020-1140-y>
- Cloitre, M., Hyland, P., Bisson, J. I., Brewin, C. R., Roberts, N. P., Karatzias, T., & Shevlin, M. (2019). ICD-11 posttraumatic stress disorder and complex posttraumatic stress disorder in the United States: A population-based study. *Journal of Traumatic Stress*, 32(6), 833-842. doi: 10.1002/jts.22454.
- Cloitre, M., Shevlin, M., Brewin, C. R., Bisson, J. I., Roberts, N. P., Maercker, A., ... & Hyland, P. (2018). The International Trauma Questionnaire: Development of a self-

report measure of ICD-11 PTSD and complex PTSD. *Acta Psychiatrica*

Scandinavica, 138(6), 536-546. <https://doi.org/10.1111/acps.12956>

Cohen, J. (1988). The effect size (2nd ed., pp. 77–83). *Statistical Power Analysis for the Behavioral Sciences*.

Dai, W., Chen, L., Lai, Z. *et al.* The incidence of post-traumatic stress disorder among survivors after earthquakes :a systematic review and meta-analysis. *BMC Psychiatry* 16, 188 (2016). <https://doi.org/10.1186/s12888-016-0891-9>

Dal Zilio, L., & Ampuero, J. P. (2023). Earthquake doublet in Turkey and Syria. *Communications Earth & Environment*, 4(1), 71. <https://doi.org/10.1038/s43247-023-00747-z>

Düzova, A., Akgül, S., ÜTİNE, G., & Yildiz, Y. (2023). The Türkiye-Syria Earthquake: a response from the editors of the Turkish Journal of Pediatrics. *Turkish Journal of Paediatrics*, 65(1). <https://doi.org/10.24953/turkjped.2023.E001>

Ehring, T., Razik, S., & Emmelkamp, P. M. (2011). Prevalence and predictors of posttraumatic stress disorder, anxiety, depression, and burnout in Pakistani earthquake recovery workers. *Psychiatry Research*, 185(1-2), 161-166. <https://doi.org/10.1016/j.psychres.2009.10.018>

Facer-Irwin E, Karatzias T, Bird A, Blackwood N, MacManus D (2022). PTSD and complex PTSD in sentenced male prisoners in the UK: prevalence, trauma antecedents, and psychiatric comorbidities. *Psychological Medicine* 52, 2794–2804. <https://doi.org/10.1017/S0033291720004936>

Folke, S., Nielsen, A. B., Andersen, S. B., Karatzias, T., & Karstoft, K. I. (2019). ICD-11 PTSD and complex PTSD in treatment-seeking Danish veterans: A latent profile

analysis. *European Journal of Psychotraumatology*, 10(1), 1686806.

doi: [10.1080/20008198.2021.1930703](https://doi.org/10.1080/20008198.2021.1930703)

Garfin, D. R., & Silver, R. C. (2023). Addressing mental health aftershocks from the Turkey–Syria earthquake: a call to action. *Nature Mental Health*, 1(4), 238-239.

Gökalp, P. G. (2002). Disaster mental health care: the experience of Turkey. *World Psychiatry*, 1(3), 159.

Goldmann, E., & Galea, S. (2014). Mental health consequences of disasters. *Annual Review of Public Health*, 35, 169-183. <https://doi.org/10.1146/annurev-publhealth-032013-182435>

Golitaleb, M., Mazaheri, E., Bonyadi, M., & Sahebi, A. (2022). Prevalence of post-traumatic stress disorder after flood: a systematic review and meta-analysis. *Frontiers in Psychiatry*, 13, 890671. doi: 10.3389/fpsy.2022.890671.

Heeke, C., O'Donald, A., Stammel, N., & Böttche, M. (2020). Same same but different? DSM-5 versus ICD-11 PTSD among traumatized refugees in Germany. *Journal of Psychosomatic Research*, 134, 110129. <https://doi.org/10.1016/j.jpsychores.2020.110129>

Hosseinnejad M, Yazdi-Feyzabadi V, Hajebi A, Bahramnejad A, Baneshi R, Ershad Sarabi R, Okhovati M, Zahedi R, Saberi H, Zolala F (2022) Prevalence of posttraumatic stress disorder following the earthquake in Iran and Pakistan: a systematic review and meta-Analysis. *Disaster Med Public Health Prep* 16: 801–808 doi: <https://doi.org/10.1017/dmp.2020.411>.

Hyland, P., Murphy, J., Shevlin, M., Vallières, F., McElroy, E., Elklit, A., ... & Cloitre, M. (2017). Variation in post-traumatic response: The role of trauma type in predicting

ICD-11 PTSD and CPTSD symptoms. *Social Psychiatry and Psychiatric Epidemiology*, 52, 727-736. <https://doi.org/10.1007/s00127-017-1350-8>

Hyland, P., Shevlin, M., Fyvie, C., & Karatzias, T. (2018). Posttraumatic stress disorder and complex posttraumatic stress disorder in DSM-5 and ICD-11: Clinical and behavioral correlates. *Journal of Traumatic Stress*, 31(2), 174-180. <https://doi.org/10.1002/jts.22272>

Hyland, P., Shevlin, M., McNally, S., Murphy, J., Hansen, M., & Elklit, A. (2016). Exploring differences between the ICD-11 and DSM-5 models of PTSD: Does it matter which model is used?. *Journal of Anxiety Disorders*, 37, 48-53. <https://doi.org/10.1016/j.janxdis.2015.11.002>

Hyland, P., Vallières, F., Cloitre, M., Ben-Ezra, M., Karatzias, T., Olf, M., ... & Shevlin, M. (2021). Trauma, PTSD, and complex PTSD in the Republic of Ireland: prevalence, service use, comorbidity, and risk factors. *Social Psychiatry and Psychiatric Epidemiology*, 56, 649-658. doi: 10.1007/s00127-020-01912-x.

Hyland, P., Murphy, J., Shevlin, M., Vallières, F., McElroy, E., Elklit, A., ... & Cloitre, M. (2017). Variation in post-traumatic response: The role of trauma type in predicting ICD-11 PTSD and CPTSD symptoms. *Social Psychiatry and Psychiatric Epidemiology*, 52, 727-736. <https://doi.org/10.1007/s00127-017-1350-8>

Iqbal, A., & Sheikh, A. (2023). The devastating earthquake that struck Turkey and Syria: Post-traumatic stress disorder and mental health issues among the victims. *The International Journal of Health Planning and Management*. doi: 10.1002/hpm.3644.

Jin, Y., Xu, J., Liu, H., & Liu, D. (2014). Posttraumatic stress disorder and posttraumatic growth among adult survivors of Wenchuan earthquake after 1 year: Prevalence and

correlates. *Archives of Psychiatric Nursing*, 28(1), 67-73.

<https://doi.org/10.1016/j.apnu.2013.10.010>

Karam, E. G., Friedman, M. J., Hill, E. D., Kessler, R. C., McLaughlin, K. A., Petukhova, M., ... & Koenen, K. C. (2014). Cumulative traumas and risk thresholds: 12-month PTSD in the World Mental Health (WMH) surveys. *Depression and anxiety*, 31(2), 130-142. <https://doi.org/10.1002/da.22169>

Karamustafalıoğlu, O. K., Fostick, L., Çevik, M., Zukerman, G., Tankaya, O., Güveli, M., ... & Zohar, J. (2023). Ten-year follow-up of earthquake survivors: long-term study on the course of PTSD following a natural disaster. *The Journal of Clinical Psychiatry*, 84(2), 45763.

Karatzias, T., Hyland, P., Bradley, A., Cloitre, M., Roberts, N. P., Bisson, J. I., & Shevlin, M. (2019). Risk factors and comorbidity of ICD-11 PTSD and complex PTSD: Findings from a trauma-exposed population-based sample of adults in the United Kingdom. *Depression and anxiety*, 36(9), 887-894. <https://doi.org/10.1002/da.22934>

Karatzias, T., Shevlin, M., Ben-Ezra, M., McElroy, E., Redican, E., Vang, M. L., ... & Hyland, P. (2023). War exposure, posttraumatic stress disorder, and complex posttraumatic stress disorder among parents living in Ukraine during the Russian war. *Acta Psychiatrica Scandinavica*. doi: 10.1111/acps.13529.

Kessler, R. C., Aguilar-Gaxiola, S., Alonso, J., Benjet, C., Bromet, E. J., Cardoso, G., ... & Koenen, K. C. (2017). Trauma and PTSD in the WHO world mental health surveys. *European journal of Psychotraumatology*, 8(sup5), 1353383.

Knefel, M., Lueger-Schuster, B., Karatzias, T., Shevlin, M., & Hyland, P. (2019). From child maltreatment to ICD-11 complex post-traumatic stress symptoms: The role of

emotion regulation and re-victimisation. *Journal of Clinical Psychology*, 75(3), 392-403. doi: [10.1002/jclp.22655](https://doi.org/10.1002/jclp.22655)

Kvedaraite, M., Gelezelyte, O., Kairyte, A., Roberts, N. P., & Kazlauskas, E. (2022). Trauma exposure and factors associated with ICD-11 PTSD and complex PTSD in the Lithuanian general population. *International Journal of Social Psychiatry*, 68(8), 1727-1736. <https://doi.org/10.1177/00207640211057720>

Lai, T. J., Chang, C. M., Connor, K. M., Lee, L. C., & Davidson, J. R. (2004). Full and partial PTSD among earthquake survivors in rural Taiwan. *Journal of Psychiatric Research*, 38(3), 313-322. DOI: 10.1016/j.jpsychires.2003.08.005

Lee, J. Y., Kim, S. W., & Kim, J. M. (2020). The impact of community disaster trauma: a focus on emerging research of PTSD and other mental health outcomes. *Chonnam Medical Journal*, 56(2), 99. doi: [10.4068/cmj.2020.56.2.99](https://doi.org/10.4068/cmj.2020.56.2.99)

Liang, Y., Cheng, J., Ruzek, J. I., & Liu, Z. (2019). Posttraumatic stress disorder following the 2008 Wenchuan earthquake: A 10-year systematic review among highly exposed populations in China. *Journal of Affective Disorders*, 243, 327-339. <https://doi.org/10.1016/j.jad.2018.09.047>

Livanou, M., Bassoglu, M., Ssalcioglu, E., & Kalendar, D. (2002). Traumatic stress responses in treatment-seeking earthquake survivors in Turkey. *The Journal of Nervous and Mental Disease*, 190(12), 816-823. DOI: 10.1097/01.NMD.0000041958.58852.1D

McGinty G, Fox R, Ben-Ezra M, Cloitre M, Karatzias T, Shevlin M, Hyland P (2021). Sex and age differences in ICD-11 PTSD and complex PTSD: An analysis of four general population samples. *European Psychiatry*, 64(1), e66, 1–9
<https://doi.org/10.1192/j.eurpsy.2021.2239>

- Morganstein, J. C., & Ursano, R. J. (2020). Ecological disasters and mental health: causes, consequences, and interventions. *Frontiers in psychiatry*, 11, 1.
<https://doi.org/10.3389/fpsy.2020.00001>
- Murphy, D., Karatzias, T., Busuttill, W., Greenberg, N., & Shevlin, M. (2021). ICD-11 posttraumatic stress disorder (PTSD) and complex PTSD (CPTSD) in treatment seeking veterans: risk factors and comorbidity. *Social Psychiatry and Psychiatric Epidemiology*, 56, 1289-1298. doi: 10.1007/s00127-021-02028-6.
- Naeem, F., Ayub, M., Masood, K., Gul, H., Khalid, M., Farrukh, A., ... & Chaudhry, H. R. (2011). Prevalence and psychosocial risk factors of PTSD: 18 months after Kashmir earthquake in Pakistan. *Journal of Affective Disorders*, 130(1-2), 268-274.
<https://doi.org/10.1016/j.jad.2010.10.035>
- Neria, Y., Nandi, A., & Galea, S. (2008). Post-traumatic stress disorder following disasters: a systematic review. *Psychological medicine*, 38(4), 467-480.
<https://doi.org/10.1017/S0033291707001353>
- North, C. S. (2016). Disaster mental health epidemiology: Methodological review and interpretation of research findings. *Psychiatry*, 79(2), 130-146. doi: 10.1080/00332747.2016.1155926.
- Önder, E., Tural, Ü., Aker, T., Kılıç, C., & Erdoğan, S. (2006). Prevalence of psychiatric disorders three years after the 1999 earthquake in Turkey: Marmara Earthquake Survey (MES). *Social Psychiatry and Psychiatric Epidemiology*, 41, 868-874.
<https://doi.org/10.1007/s00127-006-0107-6>
- Pino, O., Pelosi, A., Artoni, V., & Mari, M. (2021). Post-traumatic outcomes among survivors of the earthquake in central Italy of August 24, 2016. a study on PTSD risk

Earthquake-related traumatic events and PTSD and vulnerability factors. *Psychiatric quarterly*, 92(4), 1489-1511. doi: 10.1007/s11126-021-09908-9.

Rafiey, H., Alipour, F., LeBeau, R., & Salimi, Y. (2019). Prevalence and determinants of PTSD 3 years after an earthquake in Iran. *Community Mental Health Journal*, 55, 542-547. doi: 10.1007/s10597-019-00384-x.

Saeed, S. A., & Gargano, S. P. (2022). Natural disasters and mental health. *International Review of Psychiatry*, 34(1), 16-25. <https://psycnet.apa.org/doi/10.1080/09540261.2022.2037524>

Sayed, S., Iacoviello, B. M., & Charney, D. S. (2015). Risk factors for the development of psychopathology following trauma. *Current Psychiatry Reports*, 17, 1-7. <https://doi.org/10.1007/s11920-015-0612-y>

Shevlin, M., Hyland, P., Vallières, F., Bisson, J., Makhshvili, N., Javakhishvili, J., ... & Roberts, B. (2018). A comparison of DSM-5 and ICD-11 PTSD prevalence, comorbidity, and disability: An analysis of the Ukrainian Internally Displaced Person's Mental Health Survey. *Acta Psychiatrica Scandinavica*, 137(2), 138-147. doi: 10.1111/acps.12840.

Tang, B., Deng, Q., Glik, D., Dong, J., & Zhang, L. (2017). A meta-analysis of risk factors for post-traumatic stress disorder (PTSD) in adults and children after earthquakes. *International Journal of Environmental Research and Public Health*, 14(12), 1537. <https://doi.org/10.3390/ijerph14121537>

Tanrikulu, A. B., Kaya, H., Örum, M. H., & Akyıldırım, S. (2023). Prevalence and determinants of post-traumatic stress disorder in patients with schizophrenia 2 years after an earthquake in Turkey. *The International Journal of Psychiatry in Medicine*, 00912174231180467.

Earthquake-related traumatic events and PTSD

The World Bank IBRD IDA (2023, February 27). Earthquake Damage in Türkiye Estimated to Exceed \$34 billion: World Bank Disaster Assessment Report.

<https://www.worldbank.org/en/news/press-release/2023/02/27/earthquake-damage-in-turkiye-estimated-to-exceed-34-billion-world-bank-disaster-assessment-report>.

Tolin, D. F., & Foa, E. B. (2006). Sex differences in trauma and posttraumatic stress disorder: a quantitative review of 25 years of research. *Psychological Bulletin*, 132(6), 959.

<https://doi.org/10.1037/0033-2909.132.6.959>

Türkiye earthquake situation report #6. UNFPA Türkiye. (2023, March 22). Retrieved April 21, 2023, from <https://turkiye.unfpa.org/en/turkiye-earthquake-situation-report-no-6>

United Nations. (2023, February 22). *1.5 million now homeless in Türkiye after quake disaster, warn UN development experts*. United Nations Türkiye. <https://turkiye.un.org/en/2023-15-million-now-homeless-t%C3%BCrkiye-after-quake-disaster-warn-un-development-experts>

Wang, Y., Xu, J., & Lu, Y. (2020). Associations among trauma exposure, post-traumatic stress disorder, and depression symptoms in adolescent survivors of the 2013 Lushan earthquake. *Journal of Affective Disorders*, 264, 407-413. doi: 10.1016/j.jad.2019.11.067.

Zhang, R., Zhang, Y., & Dai, Z. (2022). Impact of natural disasters on mental health: a cross-sectional study based on the 2014 China family panel survey. *International Journal of Environmental Research and Public Health*, 19(5), doi: 10.3390/ijerph19052511.

Table 1: Endorsement Rates for Earthquake-related Experiences.

	Total Sample (N=527) (%)	Males (n= 183) (%)	Females (n= 344) (%)
1. In the days following the earthquake did you have to spend the night somewhere other than in your home?	492 (93.4%)	171 (32.4%)	321 (60.9%)
2. Did you need food and water aid after the earthquake?	444 (84.3%)	163 (30.9%)	281 (53.3%)*
3. Was your home damaged in the earthquake?	434 (82.4%)	158 (30.0%)	276 (52.4%)
4. Did you have to relocate because your house became structurally unsafe to live in?	346 (65.8%)	124 (23.6%)	222 (42.2%)
5. Were you physically injured in the earthquake?	141 (26.8%)	55 (10.4%)	86 (16.3%)
6. Were any members of your family or your loved ones physically injured in the earthquake?	287 (54.5%)	102 (19.4%)	185 (35.1%)
7. Did you lose any members of your family in the earthquake?	216 (41.0%)	75 (14.2%)	141 (26.8%)
8. Did you lose any relatives (e.g., aunts, uncles, cousins, grandparents) in the earthquake?	233 (44.2%)	89 (16.9%)	144 (27.3%)
9. Were you involved in the rescue work?	105 (19.9%)	69 (13.1%)**	36 (6.8%)
10. Did you see any dead bodies or body parts during the rescue and clearing up work period?	164 (31.1%)	81 (15.4%)**	83 (15.4%)

Table 2: Descriptive statistics for ITQ PTSD items.

PTSD items	Endorsement	Mean	SD	Mdn	Range
Re-experiencing					
Having upsetting dreams that replay part of the experience or are clearly related to the experience?	59.2 % (n = 312)	1.87	1.25	2.00	0-4
Having powerful images or memories that sometimes come into your mind in which you feel the experience is happening again in the here and now?	77.0% (n = 406)	2.46	1.18	2.00	0-4
Re cluster	82.5% (n = 435)	4.33	2.11	4.00	0-8
Avoidance					
Avoiding internal reminders of the experience (for example, thoughts, feelings, or physical sensations)?	65.6% (n = 350)	2.07	1.34	2.00	0-4
Avoiding external reminders of the experience (for example, people, places, conversations, objects, activities, or situations)?	58.9% (n = 310)	1.82	1.38	2.00	0-4
Av cluster	74.0% (n = 390)	3.90	2.46	4.00	0-8
Threat					
Being “super-alert”, watchful, or on guard?	76.4% (n = 403)	2.45	1.29	3.00	0-4
Feeling jumpy or easily startled?	75.0% (n = 395)	2.46	1.36	3.00	0-4
Th cluster	83.5% (n = 440)	4.91	2.44	5.00	0-8
Total PTSD	54.1% (n = 285)	13.13	5.76	14.00	0-24

Table 3: Logistic Regression Analysis of Age, Gender, and Earthquake-Related Experiences predicting PTSD.

	Bivariate			Multivariate		
	<i>OR</i>	<i>CI</i>	<i>p</i>	<i>OR</i>	<i>CI</i>	<i>p</i>
Age	1.01	(0.99, 1.02)	.417	1.00	(0.99, 1.02)	.709
Gender (Female)	2.18	(0.51, 3.14)	<.001	2.65	(1.75, 4.02)	<.001
In the days following the earthquake did you have to spend the night somewhere other than in your home?	1.62	(0.81, 3.24)	.171	1.48	(0.71, 3.13)	.298
Did you need food and water aid after the earthquake?	1.32	(0.83, 2.12)	.242	1.08	(0.62, 1.89)	.783
Was your home damaged in the earthquake?	1.39	(0.89, 2.18)	.150	1.17	(0.65, 2.11)	.593
Did you have to relocate because your house became structurally unsafe to live in?	1.21	(0.84, 1.73)	.308	0.73	(0.44, 1.17)	.178
Were you physically injured in the earthquake?	2.54	(1.68, 3.83)	<.001	2.26	(1.41, 3.60)	<.001
Were any members of your family or your loved ones physically injured in the earthquake?	2.23	(1.57, 3.17)	<.001	1.72	(1.13, 2.62)	.010
Did you lose any members of your family in the earthquake?	1.62	(1.14, 2.31)	.007	1.20	(0.76, 1.90)	.430
Did you lose any relatives (e.g., aunts, uncles, cousins, grandparents) in the earthquake?	1.41	(1.00, 1.99)	.053	0.95	(0.60, 1.50)	.824
Were you involved in the rescue work?	0.92	(0.60, 1.41)	.696	0.95	(0.60, 1.50)	.648
Did you see any dead bodies or body parts during the rescue and clearing up work period?	1.35	(0.93, 1.96)	.117	1.50	(0.91, 2.46)	.110

Abbreviations: OR = Odds Ratio, CI = 95% Confidence Intervals, p = significance level

Table 4: Logistic Regression Analysis of Cumulative Earthquake-Related Experiences predicting PTSD.

	PTSD n (%)	Multivariate		
		<i>OR</i>	<i>CI</i>	<i>p</i>
Age		1.00	(0.99, 1.02)	.736
Gender (Female)	39.7% (n = 209)	2.76	(1.86, 4.09)	<.001
Earthquake-related traumatic events quartile 1	9.9% (n = 53)	-	-	-
Earthquake-related traumatic events quartile 2	12.9% (n = 68)	0.78	(0.47, 1.29)	.329
Earthquake-related traumatic events quartile 3	16.1% (n = 85)	1.38	(0.81, 2.33)	.234
Earthquake-related traumatic events quartile 4	15.2% (n = 80)	3.54	(1.92, 6.51)	<.001

Abbreviations: OR = Odds Ratio, CI = 95% Confidence Intervals, p = significance level

Note: Earthquake-related traumatic events quartile 1 = those scoring 25% on total number of traumas ; Earthquake-related traumatic events quartile 2 = those scoring between 25.1% and 50% on total number of traumas ; Earthquake-related traumatic events quartile 3 = those scoring 51% to 75% on total number of traumas; Earthquake-related traumatic events quartile 4 = those scoring 100% on total number of traumas.

Table 5: Regression Analysis on PTSD symptom levels.

	Bivariate					Multivariate				
	B	SE	β	t	p	B	SE	β	t	p
Age	.009	(.018)	.022	.494	.622	.001	(.019)	.002	.051	.959
Gender (Female)	2.545	(.516)	.211	4.936	<.001	3.08	(.533)	.255	5.777	<.001
In the days following the earthquake did you have to spend the night somewhere other than in your home?	1.334	(1.007)	.058	1.325	.186	.764	(.972)	.033	.786	.432
Did you need food and water aid after the earthquake?	.885	(.688)	.056	1.286	.199	.000	(.736)	.000	.000	1.000
Was your home damaged in the earthquake?	1.321	(.656)	.088	2.014	.045	.710	(.776)	.047	.915	.361
Did you have to relocate because your house became structurally unsafe to live in?	.847	(.528)	.070	1.603	.109	-.678	(.640)	-.056	-1.058	.290
Were you physically injured in the earthquake?	2.135	(.559)	.164	3.817	<.001	1.200	(.589)	.092	2.039	.042

Earthquake-related traumatic events and PTSD

Were any members of your family or your loved ones physically injured in the earthquake?	2.872	(.488)	.249	5.882	<.001	2.026	(.558)	.175	3.632	<.001
Did you lose any members of your family in the earthquake?	1.653	(.505)	.141	3.271	.001	.176	.591	.015	.297	.766
Did you lose any relatives (e.g., aunts, uncles, cousins, grandparents) in the earthquake?	1.696	(.500)	.146	3.392	<.001	.665	.593	.057	1.121	.263
Were you involved in the rescue work?	.431	(.628)	.030	.686	.493	.329	.729	.023	.451	.652
Did you see any dead bodies or body parts during the rescue and clearing up work period?	1.386	(.539)	.112	2.571	.010	1.156	.623	.093	1.856	.064
R-squared								.137		

Abbreviations: B, unstandardized beta value; p, significance level; SE, standard error of B; t, t-statistic; β , standardised beta value.

Table 6: Regression Analysis of Cumulative Earthquake-Related Experiences on PTSD symptom levels.

	PTSD \bar{x} (SD)	Multivariate				
		B	SE	β	t	p
Age		-.004	(.018)	-.054	-.984	.325
Gender (Female)	14.01 (5.50)	3.007	(.503)	.249	5.981	< .001
Earthquake-related traumatic events quartile 1	12.27 (5.88)	-	-	-	-	-
Earthquake-related traumatic events quartile 2	11.44 (5.76)	-.675	(.686)	-.054	-.984	.325
Earthquake-related traumatic events quartile 3	14.05 (5.54)	2.065	(.703)	.163	2.938	.003
Earthquake-related traumatic events quartile 4	15.13 (5.76)	3.550	(.762)	.251	4.657	<.001

Abbreviations: B, unstandardized beta value; p, significance level; SE, standard error of B; t, t-statistic; β , standardised beta value.

Note: Earthquake-related traumatic events quartile 1 = those scoring 25% on total number of traumas ; Earthquake-related traumatic events quartile 2 = those scoring between 25.1% and 50% on total number of traumas ; Earthquake-related traumatic events quartile 3 = those scoring 51% to 75% on total number of traumas; Earthquake-related traumatic events quartile 4 = those scoring 100% on total number of traumas