

Artificial intelligence and radiographer preliminary image evaluation: What might the future hold for radiographers providing x-ray interpretation in the acute setting?

Rainey, C. (2024). Artificial intelligence and radiographer preliminary image evaluation: What might the future hold for radiographers providing x-ray interpretation in the acute setting? *Journal of Medical Radiation Sciences*, 1-4. Advance online publication. https://doi.org/10.1002/jmrs.821

Link to publication record in Ulster University Research Portal

Published in:

Journal of Medical Radiation Sciences

Publication Status:

Published online: 20/09/2024

10.1002/jmrs.821

Document Version

Publisher's PDF, also known as Version of record

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s)

If the document is licenced under Creative Commons, the rights of users of the documents can be found at https://creativecommons.org/share-your-work/cclicenses/.

Take down policy
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk

Download date: 14/11/2024

Journal of Medical Radiation Sciences

Open Access

EDITORIAL

Artificial intelligence and radiographer preliminary image evaluation: What might the future hold for radiographers providing x-ray interpretation in the acute setting?

Clare Rainey, PhD, FHEA 🕞

School of Health Sciences, Ulster University, Belfast, UK

J Med Radiat Sci 00 (2024) 1-4

doi: 10.1002/jmrs.821

Introduction

Health care services around the world are becoming increasingly stretched. This may be due to many factors, such as increasing population age, increasing burden of sickness and, importantly, limited human resources. 1-3 Indeed, the most recent Royal College of Radiologists (RCR) census in the United Kingdom (UK) noted that there is a 30% shortfall of radiologists, predicted to increase to 40% by 2028 if no action is taken.4 Additionally, the increase in the rate of technological development in radiology, suggests that imaging is being increasingly utilised.⁵ This increase in data volume adds additional pressure on an already stretched workforce, and production of suitably qualified radiology staff does not hold pace with demand, with delays in the provision of radiology reports.⁶ Radiographers are ideally placed to help in both the formal and informal review and interpretation of radiographic images. This has been supported in the Getting it Right First Time (GIRFT) radiology report⁷ in the UK, where it is noted that, in some healthcare trusts in England, radiographers are providing final, formal 'reports' on 50% of plain projection radiographic images. However, the Report also recognises that significant staffing shortages in the radiography workforce mean that this may not be possible and that the practice of formal radiographer reporting in the UK varies widely. There are, however, means of using the expertise of the radiographer other than provision of an 'official report' which can reduce patient waiting times, improve accuracy and safeguard the patient. Furthermore, new advances in computer vision and Artificial Intelligence (AI) have led to developments in the use of these technologies for assistance in image interpretation and diagnostic decision-making. These, also, have been proposed to reduce reporting turnaround times and increasing diagnostic accuracy. ^{7,8}

The Unique Advantages of Radiographer Image Evaluation

The clinical radiographer has the advantage of both production of the image, based on the clinical information provided by the referrer, and having the patient with them in the x-ray room to gain further understanding of the clinical presentation, should the need arise. They are, therefore, ideally placed to provide an initial assessment of any pathology present on the images produced. In the UK and elsewhere, some means of radiographer identification of abnormality/pathology on radiographs ('red dotting') has been in place since the 1980s; however, problems with the lack of required information to explain what the radiographer was referring to meant that this system had limited usefulness.

In 2013, The Society and College of Radiographers in the UK published guidance for the provision of Preliminary Clinical Evaluation (PCE) or Preliminary Image Evaluation (PIE) as it is known in many other parts of the world. 10 PIE describes a brief statement which provides additional information, in written format, to support the 'red dot'. Harcus and Stephens (2021)¹¹ report that emergency department clinicians radiology reporters (reporting radiographers radiologists) find a brief 'bullet-point' comment of 'What, Where and How' in relation to the pathology is desirable. Despite the simplicity of the information desired, radiographers remain reticent to adopt PIE in their clinical practice with myriad reasons cited for this, such as accountability, lack of confidence, feelings of being ill-prepared from an education and training perspective¹² and time pressures. 13 Specifically, a large Australian study Al and radiographer PIE C. Rainey

by Brown et al., ¹³ published in this edition of the Journal of Medical Radiation Sciences (JMRS), investigates the impact of 'workload' on the accuracy of PIE and reports that whilst radiographers perceive that a lack of time impacts the accuracy of their PIE, this was in fact not the case in their setting.

Why Is Initial Image Evaluation Important?

Alexander-Bates et al., 14 noted that although a formal, definitive report remains the optimal standard, this may not always be provided in a timely manner for the efficient movement of the patient through the required pathway, especially in the emergency setting. Emergency departments are feeling the impact of staffing shortages also and the provision of an initial interpretation can increase diagnostic accuracy and help the various clinicians in the emergency department. A relatively recent study by Bachmann et al., 15 found that reporting radiographers (with 1 to 4 years' experience) in a clinical centre in Denmark were significantly less likely to make errors which would be detrimental to the patient compared to radiology and orthopaedic trainees, each working for some part of their routine clinical work in the emergency department. Whilst this study focussed on reporting radiographers, it indicates that with suitable training for frontline radiographers, improvements can be made in error reduction for patients. This training is usually part of pre-registration training for radiographers in many countries. For instance, there is a requirement in the Standard of Proficiency for diagnostic radiographers for both the Health and Care Professionals Council (HCPC) in the UK and the Medical Radiation Practice Board of Australia (MRPBA) which states that practicing radiographers should be able to identify and communicate any urgent or unexpected findings to the referrer. 16,17

Radiographer PIE Capabilities

Data gathered in the UK suggests that, despite impressive PIE performances of up to 92% accuracy, there may be a higher instance of false negative interpretations (sensitivity 80%, specificity 97%).¹⁸ This, of course, requires careful scrutiny due to the impact this may have on the patient. However, the incidence and implications of a false negative report are less investigated, despite this also having impact on the patient in terms of 'overtreatment' and the impact on the service provision. This is investigated in the Australian setting by Takapautolo et al.,¹⁹ available in this edition of JMRS. Indeed, the issue of 'overservicing' is explicitly noted by The Royal Australian and New Zealand College of

Radiologists (RANZCR) and presented as a reason for radiographers to not 'report' or indeed provide PIE. Takapautolo et al. 19 propose that the findings from their large study may support the content focus for education to further improve radiographer PIE accuracy, therefore, potentially reducing the false positive rate. However, based on the findings from this study, it could be argued that in the context of PIE, some degree of 'overcalling' in cases of uncertainty may be beneficial as there were an additional 10 patients where pathology was missed by the traditional radiologist report route. The impact of educational intervention for radiographers is further supported in a smaller study by Lewis et al., 20 based in New Zealand and published in this edition, where, with regular training and feedback on their image evaluation, radiographer PIE accuracy rose over the course of the study (6 months) from 91.9% to an impressive 96.5%.

The Use of Artificial Intelligence in Radiology and the Impact on PIE and Reporting

In 2019, the National Health Service (NHS) in the UK introduced their long-term plan. This recognised the current plight of the NHS and proposed some strategies to help alleviate the pressure long term and increase the quality of the service. The Plan noted that one key area for development was in the incorporation of technology in patient treatment and care.

The articles referenced in this editorial demonstrate that radiographers are not yet confident on the provision of written PIE, and accuracy rates, whilst impressive, are not perfect. One wonders what the impact of AI will be for these clinicians? With an AI model provided to give a 'second opinion', might confidence increase, and accuracy follow? A recent study by the author of this editorial and a team in the UK found that reporting radiographers would be more certain of their diagnosis if they had agreement from an AI tool and that disagreement would drive them to seek a second opinion.²² The provision of these technologies in the initial stages of image review are not new, indeed some technology companies have developed apps which allow for the bedside assessment of projection chest radiographs for plain interpretation before a definitive report is generated.²³ With studies indicating that confidence decision-making may be a particular issue with radiographers in provision of PIE, and indications that AI assistance may help alleviate this, perhaps the use of technology and radiographer together may shift the paradigm and have impact on how radiographer PIE and by extension, reporting, is viewed by radiologists. However, there should be cognisance of the interaction

C. Rainey Al and radiographer PIE

and development of the relationship between human and machine. Studies suggest that where those who may feel less confident in their decision, such as less experienced clinicians, may relay on the feedback provided by the AI that they are using, even if this contradicts their own judgement. ^{22,24} This is known as Automation Bias and may impact users adversely in the initial stages of implementation; however, engagement with adequate education and training may alleviate this somewhat. ^{22,24–26}

Conclusions

As a profession we need to recognise the unique position clinical radiographers are placed – between technology and patient. Health care systems are struggling around the world and skilled radiographers can be utilised to relieve some of the burden. Whilst the role and development of the radiographer varies internationally, PIE is one area which all radiographers should be in a position to step in; however, this is not being fully implemented, even in the UK. The studies published in this edition of JMRS, in Australia and New Zealand by Brown, Takapautolo and Lewis 13,19,20 demonstrate that radiographers are competent and that reasons for reticence, such as workload, do not seem to have an impact on performance. However, regular education and feedback may be needed to support this. AI may also have a role in the immediate provision of this feedback but should be approached with caution in an inexperienced group. With time and development of the human-AI relationship, this may provide a useful support for radiographers working 'at the front line', allowing radiographers' confidence in their interpretation, and potentially accuracy, to increase.

Conflict of Interest

The author declares no conflict of interest.

Data Availability Statement

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

References

- 1. Holman HR. The relation of the chronic disease epidemic to the health care crisis. *ACR Open Rheumatol* 2020; **2**: 167–73.
- Gaget V, Inacio MC, Tivey DR, et al. Trends in utilisation of plain X-rays by older Australians (2010–2019). BMC Geriatr 2022; 22: 100.

3. England A. Rise in chronic diseases: will radiology survive? Eur Med J 2024; 5: 52–5.

- 4. The Royal College of Radiologists. 2023 Clinical Radiology And Clinical Oncology Workforce Census Reports. 2024 Available from: https://www.rcr.ac.uk/news-policy/latest-updates/2023-clinical-radiology-and-clinical-oncology-workforce-census-reports/#:~:text=There%20is%20a% 2030%25%20shortfall,MRI%20reporting%20surged%20by %2011%25. (accessed 19th July 2024).
- Kalidindi S, Gandhi S. Workforce crisis in radiology in the UK and the strategies to deal with it: is artificial intelligence the saviour? *Cureus* 2023; 15: e43866.
- Care Quality Commission. Radiology Review: A National Review of Radiology Reporting within the NHS in England. 2018. Available from: https://www.cqc.org.uk/ sites/default/files/20180718-radiology-reporting-reviewreport-final-for-web.pdf (accessed 24th June 2020).
- Halliday K, Maskell G, Beeley L, Quick E. Radiology: Programme National Specialty Report. 2020 Available from: https://www.gettingitrightfirsttime.co.uk/wp-content/ uploads/2020/11/GIRFT-radiology-report.pdf (accessed 20th July 2024).
- 8. Batra K, Xi Y, Bhagwat S, Espino A, Peshock RM. Radiologist worklist reprioritization using artificial intelligence: impact on report turnaround times for CTPA examinations positive for acute pulmonary embolism. *AJR Am J Roentgenol* 2023; 221: 324–33.
- Kranz R, Cosson P. Anatomical and/or pathological predictors for the "incorrect" classification of red dot markers on wrist radiographs taken following trauma. Br J Radiol 2015; 88(1046): 20140503.
- 10. The Society and College of Radiographers. Preliminary Clinical Evaluation and Clinical Reporting by Radiographers: Policy and Practice Guidance. 2013 Available from: https://www.sor.org/learning-advice/ professional-body-guidance-and-publications/documentsand-publications/policy-guidance-document-library/ preliminary-clinical-evaluation-and-clinical-repor (accessed 18th July 2024).
- 11. Harcus JW, Stevens BJ. What information is required in a preliminary clinical evaluation? A service evaluation. *Radiography* 2021; **27**: 1033–7.
- Doona D, Wright C, Harcus J. Preliminary clinical evaluation (PCE): Perceptions and barriers to implementation. 2016. In: United Kingdom Radiological Congress (UKRC) 2016, Liverpool, June 6-8 2016.
- 13. Brown C, Burck A, Neep MJ. Workload as a predictor of radiographer preliminary image evaluation accuracy. *J Med Radiat Sci* 2024. https://doi.org/10.1002/jmrs.803.
- 14. Alexander-Bates I, Neep MJ, Davis B, Starkey D. An analysis of radiographer preliminary image evaluation A focus on common false negatives. *J Med Radiat Sci* 2021; **68**: 237–44.

Al and radiographer PIE C. Rainey

- 15. Bachmann R, Ingebrigtsen RL, Holm O, et al. Comparison of reporting radiographers' and medical doctors' performance in reporting radiographs of the appendicular skeleton, referred by the emergency department. *Radiography* 2021; **27**: 1099–104.
- HCPC. Standards of Proficiency (Radiographers). 2023
 Available from: https://www.hcpc-uk.org/standards/ standards-of-proficiency/radiographers/ (accessed 18th July 2024).
- MRPBA. Professional Capabilities for Medical Radiation Practice. 2020 Available from: https://www. medicalradiationpracticeboard.gov.au/Registration-Standards/Professional-Capabilities.aspx (accessed 18th July 2024).
- 18. Verrier W, Pittock LJ, Bodoceanu M, Piper K. Accuracy of radiographer preliminary clinical evaluation of skeletal trauma radiographs, in clinical practice at a district general hospital. *Radiography* 2022; **21**: 312–8.
- 19. Takapautolo J, Neep M, Starkey D. Analysing false-positive errors when Australian radiographers use preliminary image evaluation. *J Med Radiat Sci* 2024. https://doi.org/10.1002/jmrs.809.
- Lewis K, Mdletshe S, Doubleday A, Pieterse T. Preliminary image evaluation performance of radiographers in one New Zealand District: a 6-month prospective review. *J Med Radiat Sci* 2024. https://doi.org/10.1002/jmrs.810.
- 21. NHS. The NHS Long Term Plan. 2019 Available from: https://www.longtermplan.nhs.uk/publication/nhs-long-term-plan/ (accessed 20th July 2024).

- 22. Rainey C, O'Regan T, Matthew J, et al. UK reporting radiographers' perceptions of AI in radiographic image interpretation: current perspectives and future developments. *Radiography* 2022; **28**: 881–8.
- 23. Qure.ai. AI for Blazing Fast Reporting on Chest X-Rays. n.d. Available from: https://www.qure.ai/product/qxr (accessed 20th July 2024).
- 24. Goddard K, Roudsari A, Wyatt JC. Automation bias: empirical results assessing influencing factors. *Int J Med Inform* 2014; **83**: 368–75.
- 25. Bond RR, Novotny T, Andrsova I, et al. Automation bias in medicine: The influence of automated diagnoses on interpreter accuracy and uncertainty when reading electrocardiograms. *J Electrocardiol* 2018; **51**: S6–S11.
- Rainey C, Bond R, McConnell J, Hughes C, Kumar D, McFadden S. Reporting radiographers' interaction with Artificial Intelligence how do different forms of AI feedback impact trust and decision switching? *PLOS Digit Health* 2024; 3: e0000560. https://doi.org/10.1371/journal.pdig.0000560

Correspondence

Clare Rainey, School of Health Sciences, Ulster University, York Street, Belfast BT15 1AP, UK. Tel: +44 2871 675980; E-mail: c.rainey@ulster.ac.uk