



**Danish Translation and Adaptation of the Context Assessment Index (CAI) with Implications for Evidence-based Practice**

Journal:	<i>Worldviews on Evidence-Based Nursing</i>
Manuscript ID	WVN-18-125.R2
Manuscript Type:	Original Article
Keywords:	Research Methods, Instrument development/Tool development/Test construction/Psychom, Nursing Practice, Cultural issues, Education/Curriculum/learning, Evidence-based practice, Professional issues/Professional ethics/Professional standards, Survey

SCHOLARONE™  
Manuscripts

Review

1  
2  
3  
4 11-Sep-2018

5  
6 Dear Reviewers.

7  
8 Thank you for the minor suggestions regarding our manuscript ID WVN-18-125.R1 titled "Danish  
9 Translation and Adaptation of the Context Assessment Index (CAI) with Implications for Evidence-based  
10 Practice".

11  
12 We have addressed all the comments from the reviewer and included in the attached reply to  
13 reviewers/cover letter.

14  
15 All changes in the revised article are highlighted in yellow, including one line that has been removed.

16  
17 We hope you find the paper suitable for publication and thank you for this opportunity.  
18  
19  
20  
21

Reviewer: 1	Authors responses and actions:
<p>22 23 24 Page 3, lines 39-42, what does "For both 25 models, factor loadings and fit statistics 26 were factor model RMSEA 0.07, CFI=0.923" 27 mean? I am confused.</p>	<p>This has now been corrected to the following in the abstract, Results, page 1, line 14: <i>For both models, factor loadings and fit statistics were acceptable, appropriate and statistically significant and the measurement models were confirmed (5-factor model RMSEA 0.07, CFI=0.923; 3-factor model RMSEA 0.07, CFI=0.924).</i></p>
<p>28 29 30 From the results, I cannot find that the 31 three-factor model is more advantageous 32 than the five-factor model.</p>	<p>We are sorry that we did not make this clearer. It is written in the text that the three factor model is no more advantageous, but rather the three factor model aligned closer to the PARIHS framework, which we use in our program of research. (p 1, Results and Linking Evidence to Action: <i>Cronbach alpha scores showed the models to have broadly acceptable scores (5-factor 0.64 – 0.89; 3-factor model 0.72 – 0.89) and: The three-factor model can advantageously be used when the PARIHS framework is part of the project.</i>)(S. 9 line 5 – 9: <i>There are two competing measurement models to explain the factor structure of the CAI. McCormack et al (2009) reported strong statistical evidence of a five-factor model, however Kajermo et al (2013) could not confirm the five-factor model and suggested the need to explore a theoretically derived three-factor model that aligned with elements relating to the construct 'Context' in the PARIHS framework.</i>)</p>
<p>33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 Also, it is better to provide the specific 51 values of Cronbach alpha scores in the 52 Results.</p>	<p>Thank you for this comment. We have now provided those in Results in the abstract p 1, line 15: <i>Cronbach alpha scores showed the models to have broadly acceptable scores (5-factor 0.64 – 0.89; 3-factor model 0.72 – 0.89).</i></p>

<p>I recommend to introduce the original version of CAI in the Methods including its construct and measurement properties.</p>	<p>We have now introduced the original five-factor CAI model in the Methods section including referring to the original for its construct and measurement properties, p. 4, line. 7 – 10: <i>the original model measures a 5-factor model of Collaborative Practice, Evidence informed Practice, Respect for Persons, Practice Boundaries, Evaluation. The 5-factor model has been reported to have acceptable psychometric properties (McCormack et al., 2009).</i></p>
<p>Page 10, lines 24, what is the whole questionnaire? Which sections are included in the whole questionnaire? Demographic data and CAI? I am not clear. Lines 41-42, how to define incomplete questionnaires?</p>	<p>This is now mentioned in Phase 2: Distribution of the survey, p 6. line.9-16: <i>The survey consisted of three elements: demographic data, the translation of the CAI, and a number of self-developed questions concerning the nurses attitudes and experiences regarding research in general and the implementation of new knowledge in practice in their local context. The three elements were gathered in an on-line survey developed in SurveyXact ("SurveyXact by Ramboll," 2018), a secure data management application that has certified access and encrypted communication. A personal link combined the email with the name of the individual nurse and the specific hyperlink to the survey. The link could be activated whenever wanted. In this paper, we only report the results from the CAI.</i></p>
<p>Lines 41-42, how to define incomplete questionnaires?</p>	<p>Thank you for identifying our error, this should read: Incomplete responses. It has been changed in the text, p 8. line 16: <i>Incomplete responses (n=164) were excluded from the sample.</i></p>
<p>Page 11, line 3-5, I recommend to put "(KMO &gt;0.9 is excellent, Bartlett's measure significant (Hutcheson &amp; Sofroniou, 1999))" into statistical analysis section.</p>	<p>Thank you for this suggestion. We have moved it to the start of section: Phase 3: Data analysis, p 7. line. 9-10: <i>The 37-item instrument was tested for appropriateness for factor analysis using Kaiser-Meyer-Olkin measures of sampling adequacy and the Bartlett's test for Sphericity (KMO &gt;0.9 is excellent, Bartlett's measure significant (Hutcheson &amp; Sofroniou, 1999).</i></p>
<p>From the results of confirmatory factor analysis, how to decide the three-factor model is better than the five-factor model, only based on the findings of Cronbach alpha scores?</p>	<p>As mentioned above, neither of the models were better than the other as both were strong psychometrically (with the exception of a slightly rogue Cronbach's alpha measure in 'evaluation' of the 5-factor model). We mentioned that the three-factor model was chosen because it aligned to the PARIHS framework, we hope that this is clear now.</p> <p>P 1, Results and Linking Evidence to Action: <i>Cronbach alpha scores showed the models to have broadly acceptable scores (5-factor 0.64 – 0.89; 3-factor model 0.72 – 0.89) and: The three-factor model can advantageously be used when the PARIHS framework is</i></p>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

	<p><i>part of the project.)(P 9, line 5 – 9: There are two competing measurement models to explain the factor structure of the CAI. McCormack et al (2009) reported strong statistical evidence of a five-factor model, however Kajermo et al (2013) could not confirm the five-factor model and suggested the need to explore a theoretically derived three-factor model that aligned with elements relating to the construct 'Context' in the PARIHS framework.)</i></p>
--	--

For Peer Review

## Abstract

**Background:** Healthcare contexts are rapidly changing with growing demand for health services to accommodate an ageing population and financial pressures. Assessment of context in healthcare settings has been the subject of increasing debate. The Context Assessment Index (CAI) examines three interconnected contextual elements derived from the PARIHS-Framework with the purpose of providing practitioners with an understanding of the context in which they work.

**Aims:** 1: To describe the translation of the CAI into Danish and adapt the instrument for use in Danish hospitals. 2: To evaluate the psychometric properties of the Danish version of the CAI.

**Methods:** Translation and adaption included an expert panel and a translation/back-translation process. The CAI was then sent to 4416 nurses in the Region Zealand, Denmark.

There are two alternative measurement models to explain the factor structure of the CAI, the five-factor model and the three-factor model. In order to provide the best explanation for the data both measurement models were examined using confirmatory factor analysis.

**Results:** The CAI was translated and modified based on expert review and usability testing. 2261 nurses completed the CAI. For both models, factor loadings and fit statistics were acceptable, appropriate and statistically significant, and the measurement models were confirmed (5-factor model RMSEA 0.07, CFI=0.923; 3-factor model RMSEA 0.07, CFI=0.924). Cronbach alpha scores showed the models to have broadly acceptable scores (5-factor 0.64 – 0.89; 3-factor model 0.72 – 0.89).

**Linking Evidence to Action:** The three-factor model can advantageously be used when the PARIHS framework is part of the project. In a translation process, differences in cultural specificity, language, and working environment have to be considered. By understanding the context of practice, nurses may enable person-centered care and improve patient outcomes.

## Introduction

This article describes the Danish translation and validation of the Context Assessment Index (CAI) (McCormack, McCarthy, Wright, Slater, & Coffey, 2009). The CAI was originally developed to assess readiness for use of evidence in a practice context providing care to older people and showed evidence of acceptable reliability and validity as well as practical utility (McCormack et al., 2009).

Both the original study and the later Swedish translation (Nilsson Kajermo et al., 2013) indicated that the CAI might be a suitable instrument to assess a health care organization's readiness for use of evidence, but also, that further development and evaluation was needed.

## Background

The current study is part of a newly established five-year research program: CAPAcity building in clinical Nursing (CAPAN), at Zealand University Hospital (ZUH), Denmark. CAPAN is concerned with developing clinical nursing towards person-centered practice (McCormack & McCance, 2006) and to create and implement a meaningful, accessible and flexible infrastructure for translating and integrating nursing evidence across the departments and specialties at the hospital.

Nurses' perceptions of evidence is a significant predictor of research translation into clinical practice (Lizarondo, Grimmer-Somers, & Kumar, 2011) but most nurses find that much research evidence is not easily available, applicable or adaptable to clinical practice (Saunders & Vehviläinen-Julkunen, 2016).

CAPAN is inspired by the Promoting Action on Research Implementation in Health Services (PARIHS) framework (Kitson, Harvey, & McCormack, 1998). According to PARIHS, organizational context is considered highly influential for the integration of evidence in healthcare settings. Assessment of context in healthcare settings has been the subject of increasing debate, particularly as the linkage

1  
2  
3 between context and implementation are intertwined (Pfadenhauer et al., 2015, 2017). However,  
4  
5 context is dynamic with variable transparency. As a result, it is difficult to measure the individual  
6  
7 components and how these components impact on the provision of person-centred care (Duranti &  
8  
9 Goodwin, 1992; Thomsen, Soelver, & Hølge-Hazelton, 2017). Consequently, the development of  
10  
11 measurement tools for assessing context may offer greater insight in developing strategies for im-  
12  
13 plementation (Nilsson Kajermo et al., 2013). One example of context assessment tools, that are  
14  
15 based on PARIHS and have been translated and tested in international settings, is the Alberta Con-  
16  
17 text Tool (ACT) (Eldh, Ehrenberg, Squires, Estabrooks, & Wallin, 2013; Estabrooks, Squires, Cum-  
18  
19 mings, Birdell, & Norton, 2009; Hoben et al., 2013), another is the CAI (McCormack et al., 2009;  
20  
21 Nilsson Kajermo et al., 2013).

22  
23  
24  
25  
26 The ACT was designed to measure the context of research utilisation (Estabrooks et al., 2009) and  
27  
28 the CAI examines the readiness of a practice context for establishing person-centered care with the  
29  
30 specific purpose of providing practitioners with an understanding of the context in which they  
31  
32 work. The CAI measurements are closely related to the purpose of CAPAN and was therefore cho-  
33  
34 sen as a tool.  
35  
36

37  
38 Even though testing of the CAI has shown reliability (McCormack et al., 2009) Kajermo et al have  
39  
40 suggested, that further evaluation of its psychometric properties is required (Nilsson Kajermo et al.,  
41  
42 2013).  
43  
44

## 45 46 Aims of the study

47  
48  
49 The aims of the current study were to:

- 50  
51  
52 1) Describe the translation and adaptation of the CAI for use in Danish hospitals.
- 53  
54  
55 2) Evaluate the psychometric properties of the Danish version of the CAI.

## Methods

This study consisted of three phases: 1: Instrument translation and modifications. 2: Distribution of the survey. 3: Data analysis.

### Phase 1: Instrument translation and modifications

Before entering the translation process, consent was obtained from the instrument copyright holder.

The CAI consists of 37 items with four response alternatives on a Likert scale: *Strongly agree*, *Agree*, *Disagree* and *Strongly disagree* and the original model measures a 5-factor model of Collaborative Practice, Evidence informed Practice, Respect for Persons, Practice Boundaries, Evaluation. The 5-factor model has been reported to have acceptable psychometric properties (McCormack et al., 2009).

Two bilingual translators whose first language is Danish produced two independent translations. The translators had different profiles, one with a clinical background and one with no in-depth knowledge of the health care system. This combination contributed to a translation reflecting both a more reliable equivalence from a measurement perspective and a translation that could highlight ambiguous meanings in the original questionnaire (Beaton, Bombardier, Guillemin, & Ferraz, 2000).

#### *Expert review*

The two translations were reviewed and discussed by an expert panel consisting of five bilingual nurse researchers, all holding at least a PhD-degree, one member having English as a first language.

The deliberation process consisted of cultural differences in the health care systems, terminology in daily clinical settings, structure of the organization and disparity in the languages. For instance, staff appointments in a UK context differ from the Danish context. Therefore, the distinction of the clini-



1  
2  
3 cal and non-clinical staff was discussed thoroughly before choosing staff designations. Moreover,  
4  
5 the CAI was originally developed for an older people context where multidisciplinary teams are  
6  
7 much more integrated than the teams are in general departments at Danish hospitals.  
8  
9

10 The panel agreed on a version for back-translation.  
11

### 12 *Back-translation*

13  
14  
15 A professional translator with English as first language made a back-translation to English. The  
16  
17 translator had no clinical background and was not aware of the intent and concepts in the ques-  
18  
19 tionnaire which could reveal unexpected meanings in the pre-final version (Beaton et al., 2000;  
20  
21 Guillemin, Bombardier, & Beaton, 1993; Wild et al., 2005).  
22  
23

24  
25 The copyright holder was presented with the back-translation to revise and discuss. Item 13, "*Staff*  
26  
27 *have explicit understanding of their own attitudes and beliefs toward the provision of care*", raised  
28  
29 some discussion as *attitudes* directly translated into Danish has a different meaning in every-day  
30  
31 language than it does in English. The translation was subsequently accepted.  
32  
33

### 34 *Usability testing*

35  
36  
37 To further emphasize the comprehensibility and thereby validate the usability in the target group  
38  
39 the questionnaire underwent a cognitive debriefing in a group of nurses similar to the target group  
40  
41 (Wild et al., 2005). The pilot respondents were native Danish staff from another similar hospital.  
42  
43

44  
45 Both the introductory letter, explanatory texts and the items in the CAI were displayed in the same  
46  
47 look and digital questionnaire form as the final version would be.  
48  
49

50  
51 Additionally, in several places of the pilot questionnaire, the respondents could write their opinions  
52  
53 on matters such as clarity of language and comprehensibility. These answers identified confusing  
54  
55 and unclear language and confirmed cultural relevance. Three items were highlighted as difficult to  
56  
57

1  
2  
3 interpret. Therefore, these were modified in consultation with the expert panel and subsequently  
4  
5 retested in the pilot group.  
6

7  
8 An explanation of the term *Evidence* was added to the text due to comments from the pilot group  
9  
10 regarding insufficient understanding of the term.  
11

12  
13 The expert panel reached consensus on the final version.  
14  
15

## 16 Phase 2: Distribution of the survey

### 17 *Sample and Setting - Data Collection*

18  
19  
20  
21 The survey consisted of three sections: 1) demographic data, 2) the translation of the CAI, and 3) a  
22  
23 number of self-developed questions concerning the nurses attitudes and experiences regarding  
24  
25 research in general and the implementation of new knowledge in practice in their local context. The  
26  
27 three sections were gathered in an on-line survey developed in SurveyXact ("SurveyXact by Ram-  
28  
29 boll," 2018), a secure data management application that has certified access and encrypted com-  
30  
31 munication. An email was sent to the individual nurse with a specific hyperlink to the survey. The  
32  
33 link could be activated whenever wanted. In this paper, we only report the results from the CAI.  
34  
35  
36  
37

38 In order to compare the context at ZUH with other hospitals in the region that are not in a trans-  
39  
40 formation process of becoming at university hospital, all 4416 hospital employed nurses from Re-  
41  
42 gion Zealand were invited to participate in the study. Of these, 1673 came from the ZUH, 2194 from  
43  
44 the non-university hospitals, and 549 from the psychiatric hospital. All nurses were included even if  
45  
46 on sick leave, parental leave or leave of absence due to other reasons, as it was possible to com-  
47  
48 plete the questionnaire online. Reminders and a link to the questionnaire were sent to those who  
49  
50 had not completed or started the questionnaire within 10 days. Further reminders were sent one  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 week and two weeks after the first reminder. In total, the questionnaire was open for completion  
4  
5 for three months after which it was automatically closed.  
6  
7

8 The ratio of respondent to item is important in factor analysis and Nunally (1978) recommend at  
9  
10 least a ratio of 10:1 while Osborne and Costello (2004) reported the that bigger the sample size the  
11  
12 better. In this study a 59:1 ratio of respondent to item was achieved.  
13  
14

### 15 Phase 3: Data analysis

16  
17  
18 SPSS 23.0 and Mplus were used in the statistical analysis of the data set.  
19  
20

21 The 37-item instrument was tested for appropriateness for factor analysis using Kaiser-Meyer-Olkin  
22  
23 measures of sampling adequacy and the Bartlett's test for Sphericity (KMO >0.9 is excellent, Bart-  
24  
25 letts measure significant (Hutcheson & Sofroniou, 1999)). The original model was established using  
26  
27 exploratory factor analysis, and the stability of the measurement model was tested in this study  
28  
29 using confirmatory factor analysis with an exploratory element. Model modifications were identi-  
30  
31 fied using statistical feedback from the modification indices, and based on a criterion of being theo-  
32  
33 retically relevant, introduced one at a time and selected on highest score first (exceeding scores of  
34  
35 3.98) (Byrne, 2013). Firstly, within factor correlated errors, secondly between factor correlated  
36  
37 errors. Only statistically significant relationships were maintained in the final model in order to  
38  
39 maintain as parsimonious a model as possible. Acceptable factor loadings were based on the sam-  
40  
41 ple size and were set at 0.35 (Hair, Anderson, Black, & Babin, 2010). Cronbach' alpha scores were  
42  
43 also generated for the final factors in the model. Acceptable fit statistics were set at Root Mean  
44  
45 Square Estimations of Approximation (RMSEA) of 0.05 or below; 90% RMSEA higher bracket below  
46  
47 0.08; and Confirmation Fit Indices (CFI) of >0.90 (Hu & Bentler, 1999).  
48  
49  
50  
51  
52  
53

54 Inferential statistics and post hoc analyses were performed in order to compare the three sites,  
55  
56 ZUH, non-university hospitals and the psychiatric hospital.  
57  
58  
59  
60

## Ethical Considerations

Ethical approval was obtained from the regional ethics committee in Denmark (J.nr. 17-000048).

Study participants were provided with written information concerning the purpose of the study, instructions for withdrawal and their anonymity of the survey.

Permission for completion of the survey was obtained from the hospital management at each hospital and subsequently at the Data Protection Agency in Denmark (j.nr. 2008-58-0020).

## Results

### Sample characteristics

Of the invited 4416 registered nurses, 2181 (49.4%) completed the whole questionnaire and 2261 (51.2%) completed the CAI. The response rate varied from 30% to 76.9% for the participating wards. 46 (2.0%) of the respondents held a master degree in nursing, 15 (0.7%) a PhD-degree and 888 (39.3%) reported no formal education other than their nursing degree (in Denmark there is only one nursing degree which is Bachelor of Nursing). Of the respondents, 96.0% were women and the mean age was 45.8 years (range 22 – 76 y). In all, 44.2% (n=999) were from ZUH, 44.0% (n=994) from non-University hospital and 11.9% from the psychiatric hospital (n=268). Incomplete responses (n=164) were excluded from the sample.

### Item Analysis

The mean scores for 34 of the 37 items scored at a high level of agreement (>2.5 with 22 items) to strongly agree (>3.0 with 12 items). Examination of the Kaiser-Meyer-Olkin measures of sampling adequacy (0.970) and the Bartlett's test for Sphericity (37372, df=630, p=0.001) indicated that the 37 items were appropriate to analysis using factor analysis (KMO >0.9 is excellent, Bartlett's measure significant (Hutcheson & Sofroniou, 1999)). A review of the correlations between items showed

1  
2  
3 that all items were moderate and positively correlated. Rho scores ranged from 0.248 to 0.774, in-  
4  
5 dicating no collinearity between items.  
6  
7

### 8 *Statistics of Fit of Model*

9

10 There are two competing measurement models to explain the factor structure of the CAI. McCor-  
11  
12 mack et al (2009) reported strong statistical evidence of a five-factor model, however Kajermo et al  
13  
14 (2013) could not confirm the five-factor model and suggested the need to explore a theoretically  
15  
16 derived three-factor model that aligned with elements relating to the construct 'Context' in the  
17  
18 PARIHS framework. In order to provide the best explanation for the data both measurement mod-  
19  
20 els were examined in this study. Examination of the normality of distribution demonstrated that the  
21  
22 items were non-normally distributed and Weighted Least Square Mean and Variance (WLSMV) is a  
23  
24 robust estimator which does not assume normally distributed variables and provides the best op-  
25  
26 tion for modelling categorical or ordered data (Brown, 2015).  
27  
28  
29

### 30 *Five-factor model*

31

32 Modifications were permitted, and guided by the suggestions identified in the modification matrix  
33  
34 and limited to correlated errors. The original model fit statistics were not acceptable: Root mean  
35  
36 square error of approximation (RMSEA) = 0.090; Df = 602; 90%; p = 0.001; RMSEA C.I. = 0.089 –  
37  
38 0.092; CFI = 0.870. Correlated errors both within-factor items and between-factor items were per-  
39  
40 mitted in the model. These included the correlated errors selected base on the modification indices  
41  
42 > 5, with highest scored modification introduced first. All correlated errors were statistically signifi-  
43  
44 cant (p≤0.05). Once introduced, the model produced acceptable fit statistics: Root mean square  
45  
46 error of approximation (RMSEA) = 0.070; Df = 602; 90%; p = 0.001; RMSEA C.I. = 0.069 – 0.072; CFI =  
47  
48 0.924. All factor loading, and fit statistics were acceptable, appropriate and statistically significant  
49  
50 (see **Error! Reference source not found.**) and the measurement model was confirmed.  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Cronbach alpha scores indicated that four of the five items were acceptable (Collaborative Practice:  
4 0.83, Evidence informed Practice: 0.89, Respect for Persons: 0.77, Practice Boundaries: 0.78, Evalu-  
5 ation: 0.64). One factor (Evaluation) scored 0.64, indicating a score that is slightly below acceptable.  
6  
7 However, all Cronbach alpha scores must be viewed with an element of caution especially when  
8 using factors with a large number (>6) items within them (Sijtsma, 2009).  
9

10  
11  
12 Statistically significant differences were noted across hospital sites on the constructs 'Evaluation',  
13 (f=18.30, p=0.001) and 'Collaborative Practice' (f=13.09, p=0.001). Post hoc analysis shows these  
14 differences to be between the psychiatric hospital (x=2.95) and the other two sites (ZUH (x=2.74)  
15 and non-university hospitals (x=2.70)) on 'Evaluation' (p=0.01 and p=0.01 respectively) and 'Collab-  
16 orative Practice' (psychiatric hospital x=2.83 vs ZUH x=2.80, p=0.01 and non-university hospitals  
17 x=2.96, p=0.01 respectively).  
18  
19

### 20 *Three-factor model*

21  
22 The three-factor model was tested and produced unacceptable fit statistics: Root mean square er-  
23 ror of approximation (RMSEA) = 0.090; Df = 602; 90% RMSEA C.I. = 0.089 – 0.092; p = 0.001; CFI =  
24 0.868. All correlated errors were statistically significant (p≤0.05). Correlated errors both within-  
25 factors and between-factors items were permitted in the model. Once introduced the model pro-  
26 duced acceptable fit statistics: Root mean square error of approximation (RMSEA) = 0.070; Df = 602;  
27 90%; RMSEA C.I. = 0.069 – 0.072; p = 0.001; CFI = 0.923. All factor loading, and fit statistics were  
28 acceptable, appropriate and statistically significant (see **Error! Reference source not found.**) and  
29 the measurement model was confirmed.  
30

31  
32 Examination of the Cronbach alpha scores showed the three-factor model too had acceptable  
33 scores (Culture 0.89, Leadership 0.72, and Evaluation 0.88).  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Statistically significant differences were noted across hospital sites on the three constructs 'Evaluation'  
4 'tion' ( $f=3.60$ ,  $p=0.03$ ), 'Leadership', ( $f=37.04$ ,  $p=0.001$ ), 'Culture' ( $f=5.18$ ,  $p=0.006$ ). Post hoc analy-  
5 sis shows these differences to be between psychiatric hospital ( $x=2.81$ ) and non-university hospitals  
6 ( $x=2.72$ ) on 'Evaluation', 'Leadership' with psychiatric hospital ( $x=2.85$ ,  $p=0.04$ ) and ZUH ( $x=2.77$ ,  
7  $p=0.001$ ); and 'Culture' with psychiatric hospital ( $x=3.11$ ) and non-university hospitals ( $x=3.03$ ,  
8  $p=0.007$ ) and ZUH ( $x=3.03$ ,  $p=0.007$ ).

9  
10  
11  
12  
13  
14  
15  
16  
17 **INSERT TABLE 1 HERE**

## 18 Discussion

19  
20  
21  
22  
23 The aims of this paper were to describe the translation of the CAI into Danish and to evaluate the  
24 psychometric properties of the Danish version. The main findings demonstrated factor loadings and  
25 fit statistics were acceptable, appropriate and statistically significant and the measurement models  
26 for both the five-factor and three-factor models were confirmed.

### 27 *Translation and adaptation*

28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
When adapting self-report measures to a setting that is different in both country, culture and lan-  
guage, unique methods are necessary. In the translation process, several aspects and differences in  
cultural specificity, language, and working environment had to be considered. For instance, the  
original CAI was developed for use in settings providing non-acute care to older people (McCormack  
et al., 2009). In the UK, multidisciplinary teams (MDT) in older people care settings are integrated  
differently than in Denmark, which is reflected in the original CAI (e.g. item 10: *HCPs in the MDT  
have equal authority in decision-making*). In the translation to Danish, the instrument had to be  
altered so that the Danish edition is neutral to the type of care provided.

1  
2  
3 In addition, the significantly greater number of participants could alter the results compared to  
4  
5 smaller studies.  
6

7  
8 The degree of interception in the current study is estimated acceptable according to the confirma-  
9  
10 tory factor analysis. Besides, a definition of evidence was included in the introduction to the ques-  
11  
12 tionnaire. Nilsson Kajermo et al. (2013) assessed such a definition as enhancing the validity of the  
13  
14 CAI.  
15

### 16 17 *Psychometric evaluation*

18  
19  
20 Two main findings emerged when testing the psychometric properties of the Danish version of the  
21  
22 instrument with the two models.  
23

24  
25 Firstly, the two models produced similar findings, and both required similar levels of modification to  
26  
27 achieve acceptable statistical fit. By analysing the dataset by both models, we found acceptable  
28  
29 factor loadings and produced acceptable Cronbach alpha scores of the Danish version of the in-  
30  
31 strument.  
32  
33

34  
35 Secondly, the Danish validation produced different results to the former validation studies of the  
36  
37 CAI (McCormack et al., 2009; Nilsson Kajermo et al., 2013). These studies were conducted in other  
38  
39 geographical regions and possibly highlight the sensitivity of the tool to variations in context as it  
40  
41 was designed to do.  
42  
43

44  
45 In addition, inferential statistics and post hoc analyses demonstrated that the CAI was able to iden-  
46  
47 tify differences across all three sites. All sites scored the constructs in both models positively, how-  
48  
49 ever there were statistically significant differences between hospital sites construct scores, with  
50  
51 psychiatric units scoring the constructs more positively. This difference was noted more on the  
52  
53 three construct version of the CAI.  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 The CAI focuses on elements of PARIHS, including culture, leadership and evaluation.  
4

5  
6 *Culture:* McCormack refers to Drennan's (1992) definition of organizational culture, as 'how things  
7 are done around here', it is what holds the organization together and even though it is strong, it can  
8 be transformed as responsive to a changing context. In a Danish context, recent sociological re-  
9 search describes that the public sector, due to new public management, has been transformed into  
10 a *culture of silence* where staff are expected to be hardened and robust, withholding critique or  
11 attempts to change practice (Willig, 2016).  
12  
13  
14  
15  
16  
17  
18

19  
20 *Leadership:* Within the PARIHS framework, leadership is identified as something everyone has po-  
21 tential to develop and release (Rycroft-Malone, 2004). However, it is also necessary that nurse  
22 leaders have transformative abilities and knowledge about different kinds of evidence (Holge-  
23 Hazelton, Kjerholt, Berthelsen, & Thomsen, 2015). In Denmark, evidence based nursing is still in its  
24 early years, and can be challenging for leaders to include, particularly in more rural regions such as  
25 the one where the present study took place.  
26  
27  
28  
29  
30  
31  
32  
33

34 *Evaluation:* Evaluation and documentation are key issues in nursing practice (Blair & Smith, 2012),  
35 and at Danish hospitals auditing and benchmarking with other health services have become key  
36 activities and tools (Ernst, 2016). The dominating positions in practice are promoting efficiency and  
37 standardization, which promote measurement and categorization in nursing (Holen, 2011). The  
38 current implementation of a new IT-system, EPIC, is one example of this.  
39  
40  
41  
42  
43  
44  
45

46 In other words, since the CAI assesses context, it is not surprising that different validation studies  
47 showed different results. A systematic review of the cross-cultural equivalence of participation  
48 instruments (Stevelinck & van Brakel, 2013) highlights the same issue and stresses the importance of  
49 being aware of prior testing of cultural validity in new contexts.  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Linking Evidence to Action

- If a project is using the PARIHS framework, the three-factor model is relevant as a context assessment tool.
- When adapting self-report measures to a setting that is different in both country, culture and language, unique methods are necessary.
- During the translation process differences in cultural specificity, language, and working environment have to be considered.
- When using CAI, sample size and the variation in setting should be considered as this may produce differences in results.
- By understanding the context of practice, nurses may enable person-centered care and improve patient outcomes.

### Strengths and limitations of the study

This study is larger than both the original study consisting of 460 nurses (McCormack et al., 2009) and the later study translating the CAI into Swedish consisting of 375 nurses (Nilsson Kajermo et al., 2013).

In 27 of the 37 items, more than 15% of the respondents chose the highest value that could mirror a ceiling effect. A ceiling effect could be a result from limited response alternatives and can reduce the variability in the gathered data. This bias is also mentioned by Nilsson Kajermo et al. (2013). Furthermore, a sample of the respondents expressed the need to have the opportunity to tick a “do not know” box.

The CAI was tested in several settings, not just one sub-speciality or working environment, neutral to the type of care provided and thereby applicable in many Danish settings.

### Conclusions

This study described and evaluated the CAI framework adapted for use in Danish hospitals. Over 2000 nurses covering a wide geographic area tested the instrument, offering a solid basis for evalu-

1  
2  
3 ating the context of practice, and nurses readiness to implement evidence into practice throughout  
4  
5 Region Zealand in Denmark.

6  
7  
8 The statistics demonstrated that the five- and three-factor model (with modifications) are equally  
9  
10 acceptable, however potentially more useful in practice. It also demonstrated that the psychiatric  
11  
12 hospital had a more positive workplace context as defined by culture, leadership and the use of  
13  
14 evaluation.

15  
16  
17 The CAI has the potential to provide practitioners with an understanding of context in which they  
18  
19 work. This is an important step, in order to enable nurses to integrate evidence and undertake per-  
20  
21 son-centred care, ultimately leading to improved patient outcomes.

22  
23  
24 The current study establishes the three-factor model, but further research is required.

## 25 26 27 28 Acknowledgements

29  
30  
31 We thank Zealand University Hospital, postdoc Jonathan David Comins and postdoc Elizabeth Emilie  
32  
33 Rosted, ZUH, and colleagues from Aarhus University Hospital for assistance and support.

## 34 35 36 37 38 39 References

- 40  
41 Beaton, D. E., Bombardier, C., Guillemin, F., & Ferraz, M. B. (2000). Guidelines for the process of  
42  
43 cross-cultural adaptation of self-report measures. *Spine*, 25(24), 3186–3191.
- 44  
45 Blair, W., & Smith, B. (2012). Nursing documentation: frameworks and barriers. *Contemporary*  
46  
47 *Nurse*, 41(2), 160–168. <https://doi.org/10.5172/conu.2012.41.2.160>
- 48  
49  
50 Brown, T. A. (2015). *Confirmatory Factor Analysis for Applied Research* (Second edition). New  
51  
52 York: Guilford Publications.
- 53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Byrne, B. M. (2013). *Structural Equation Modeling with Mplus: Basic Concepts, Applications, and*  
4  
5 *Programming*. New York: Routledge.  
6  
7 Drennan, D. (1992). *Transforming Company Culture: Getting Your Company from Where You Are*  
8  
9 *Now to Where You Want to Be*. London, UK: McGraw- Hill.  
10  
11 Duranti, A., & Goodwin, C. (1992). *Rethinking context: Language as an interactive phenomenon*.  
12  
13 Great Britain: Cambridge University Press.  
14  
15 Eldh, A. C., Ehrenberg, A., Squires, J. E., Estabrooks, C. A., & Wallin, L. (2013). Translating and  
16  
17 testing the Alberta context tool for use among nurses in Swedish elder care. *BMC Health Ser-*  
18  
19 *vices Research, 13*, 68. <https://doi.org/10.1186/1472-6963-13-68>  
20  
21 Ernst, J. (2016). Evidence-based Nursing in the IED: From Caring to Curing? *Nordic Journal of*  
22  
23 *Working Life Studies, 6*(1), 47–66. <https://doi.org/10.19154/njwls.v6i1.4885>  
24  
25 Estabrooks, C. A., Squires, J. E., Cummings, G. G., Birdell, J. M., & Norton, P. G. (2009). Devel-  
26  
27 opment and assessment of the Alberta Context Tool. *BioMed Central Health Services Re-*  
28  
29 *search, 9*, 234. <https://doi.org/10.1186/1472-6963-9-234>  
30  
31 Guillemin, F., Bombardier, C., & Beaton, D. (1993). Cross-cultural adaptation of health-related qual-  
32  
33 ity of life measures: Literature review and proposed guidelines. *Journal of Clinical Epidemi-*  
34  
35 *ology, 46*(12), 1417–1432. [https://doi.org/10.1016/0895-4356\(93\)90142-N](https://doi.org/10.1016/0895-4356(93)90142-N)  
36  
37 Hair, J., Anderson, R., Black, B., & Babin, B. (2010). *Multivariate Data Analysis* (7th ed.). Pearson  
38  
39 Education.  
40  
41 Hoben, M., Mahler, C., Bär, M., Berger, S., Squires, J. E., Estabrooks, C. A., & Behrens, J. (2013).  
42  
43 German translation of the Alberta context tool and two measures of research use: methods,  
44  
45 challenges and lessons learned. *BMC Health Services Research, 13*, 478.  
46  
47  
48  
49  
50 <https://doi.org/10.1186/1472-6963-13-478>  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Hølen, M. (2011). *Medinddragelse og lighed - en god idé? En analyse af patient-tilblivelser i det*  
4 *moderne hospital [Inclusiveness and equality - a good idea? An analysis of patient-*  
5 *becomings in the modern hospital]*. Roskilde Universitet, Forskerskolen i Livslang Læring.  
6  
7  
8  
9 Hølge-Hazelton, B., Kjerholt, M., Berthelsen, C. B., & Thomsen, T. G. (2015). Integrating nurse  
10 researchers in clinical practice - a challenging, but necessary task for nurse leaders. *Journal*  
11 *of Nursing Management*. <https://doi.org/10.1111/jonm.12345>  
12  
13  
14  
15 Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Con-  
16 ventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary*  
17 *Journal*, 6(1), 1–55. <https://doi.org/10.1080/10705519909540118>  
18  
19  
20  
21  
22 Hutcheson, G., & Sofroniou, N. (1999). *The Multivariate Social Scientist: Introductory Statistics*  
23 *Using Generalized Linear Models* (1 edition). London: SAGE Publications Ltd.  
24  
25  
26  
27 Kitson, A. L., Harvey, G., & McCormack, B. (1998). Enabling the implementation of evidence-based  
28 practice: a conceptual framework. *Qual Health Care*, 7.  
29  
30  
31 Lizarondo, L., Grimmer-Somers, K., & Kumar, S. (2011). A systematic review of the individual de-  
32 terminants of research evidence use in allied health. *Journal of Multidisciplinary Healthcare*,  
33 4, 261–272. <https://doi.org/10.2147/JMDH.S23144>  
34  
35  
36  
37 McCormack, B., & McCance, T. V. (2006). Development of a framework for person-centred nurs-  
38 ing. *Journal of Advanced Nursing*, 56(5), 472–479. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2648.2006.04042.x)  
39 [2648.2006.04042.x](https://doi.org/10.1111/j.1365-2648.2006.04042.x)  
40  
41  
42  
43 McCormack, B., McCarthy, G., Wright, J., Slater, P., & Coffey, A. (2009). Development and testing  
44 of the Context Assessment Index (CAI). *Worldviews on Evidence-Based Nursing*, 6(1), 27–  
45 35. <https://doi.org/10.1111/j.1741-6787.2008.00130.x>  
46  
47  
48  
49  
50 Nilsson Kajermo, K., Bøe, H., Johansson, E., Henriksen, E., McCormack, B., Gustavsson, J. P., &  
51 Wallin, L. (2013). Swedish translation, adaptation and psychometric evaluation of the Con-  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 text Assessment Index (CAI). *Worldviews on Evidence-Based Nursing*, 10(1), 41–50.

4  
5 <https://doi.org/10.1111/j.1741-6787.2012.00252.x>

6  
7 Nunnally, J. C. (1978). *Psychometric Theory* (2nd edition). New York: The McGraw-Hill Compa-  
8  
9 nies.

10  
11 Osborne, J., & Costello, A. B. (2004). Sample size and subject to item ratio in principal components  
12  
13 analysis. *Practical Assessment, Research & Evaluation*, 9(11). Retrieved from

14  
15 <http://PAREonline.net/getvn.asp?v=9&n=11>

16  
17 Pfadenhauer, L. M., Ansgar Gerhardus, Kati Mozygemba, Kristin Bakke Lysdahl, Andrew Booth,  
18  
19 Bjørn Hofmann, ... Eva Rehfuss. (2017). Making sense of complexity in context and im-  
20  
21 plementation: the Context and Implementation of Complex Interventions (CICI) framework.

22  
23 *Implementation Science*, 12(21). <https://doi.org/10.1186/s13012-017-0552-5>

24  
25 Pfadenhauer, L. M., Mozygemba, K., Gerhardus, A., Hofmann, B., Booth, A., Lysdahl, K. B., ...

26  
27 Rehfuss, E. A. (2015). Context and implementation: A concept analysis towards conceptual  
28  
29 maturity. *Zeitschrift Fur Evidenz, Fortbildung Und Qualitat Im Gesundheitswesen*, 109(2),

30  
31 103–114. <https://doi.org/10.1016/j.zefq.2015.01.004>

32  
33 Rycroft-Malone, J. (2004). The PARIHS framework - a framework for guiding the implementation  
34  
35 of evidence-based practice. *Journal of Nursing Care Quality*, 19(4), 297–304.

36  
37 Saunders, H., & Vehviläinen-Julkunen, K. (2016). The state of readiness for evidence-based practice  
38  
39 among nurses: An integrative review. *International Journal of Nursing Studies*, 56, 128–140.

40  
41 <https://doi.org/10.1016/j.ijnurstu.2015.10.018>

42  
43 Sijtsma, K. (2009). On the Use, the Misuse, and the Very Limited Usefulness of Cronbach's Alpha.

44  
45 *Psychometrika*, 74(1), 107–120. <https://doi.org/10.1007/s11336-008-9101-0>

46  
47 Stevelink, S. a. M., & van Brakel, W. H. (2013). The cross-cultural equivalence of participation in-  
48  
49 struments: a systematic review. *Disability and Rehabilitation*, 35(15), 1256–1268.

50  
51 <https://doi.org/10.3109/09638288.2012.731132>

- 1  
2  
3 SurveyXact by Ramboll. (2018). Retrieved November 10, 2017, from <https://www.surveymxact.com/>  
4  
5 Thomsen, T. G., Soelver, L., & Hølge-Hazelton, B. (2017). The influence of contextual factors on  
6  
7 patient involvement during follow-up consultations after colorectal cancer surgery: a case  
8  
9 study. *Journal of Clinical Nursing*, 26, 3688–3698. <https://doi.org/10.1111/jocn.13741>  
10  
11 Wild, D., Grove, A., Martin, M., Eremenco, S., McElroy, S., Verjee-Lorenz, A., ... ISPOR Task  
12  
13 Force for Translation and Cultural Adaptation. (2005). Principles of Good Practice for the  
14  
15 Translation and Cultural Adaptation Process for Patient-Reported Outcomes (PRO)  
16  
17 Measures: report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value in*  
18  
19 *Health: The Journal of the International Society for Pharmacoeconomics and Outcomes Re-*  
20  
21 *search*, 8(2), 94–104. <https://doi.org/10.1111/j.1524-4733.2005.04054.x>  
22  
23  
24 Willig, R. (2016). *Afvæbnet kritik - om flexismens små doser af arsenik [Disarmed criticism - about*  
25  
26 *small doses of arsenic in flexism]* (1.). Latvia: Hans Reitzels Forlag. Retrieved from  
27  
28 <http://ibog3.gyldendal.dk/Afvaebnetkritik#/double/4/>  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Table 1: Standardized factor loadings scores of the CAI shown for both five-factor and the three-factor models. The brackets refers to position in the respective models. Five-factor model constructs: Collaborative Practice (CP), Evidence informed Practice (EIP), Respect for Persons (RP), Practice Boundaries (PB) Evaluation (E3). Three-factor model constructs: Culture (C), Leadership (L), Evaluation (E3)

ITEMS	Three-Factor loading	Five-Factor Loading
1. Personal and professional boundaries between healthcare staff are respected (PB)(C)	0.555	0.543
2. Decisions regarding care and treatment are clearly documented by all employees (RP)(L)	0.541	0.616
3. Healthcare staff work proactively and prospectively (CP)(C)	0.617	0.619
4. All aspects of care and treatment are based on best available knowledge/evidence (EIP)(E3)	0.638	0.565
5. Nurse managers are role models for good clinical practice (RP)(E)	0.626	0.567
6. Healthcare staff allow patients to participate in decisions regarding their own care and treatment (CP) (L)	0.545	0.607
7. The further education of healthcare staff is prioritized	0.627	0.671
8. There is a high level of cooperation between healthcare staff and other professional groups (e.g. secretaries, porters, cleaning) (RP)(E3)	0.541	0.539
9. Healthcare staff receive feedback on the outcome of complaints (E)(C)	0.564	0.714
10. Healthcare staff in an interdisciplinary team have equal authority in clinical decision making (CP) (L)	0.626	0.729
11. Results from audits and/or research are used to develop clinical practice (EIP)(E3)	0.698	0.672
12. Performance and development reviews (PDRs) are conducted on a regular basis, which allows healthcare staff to set goals and reflect on their own practice (E)(C)	0.567	0.625
13. The staff have a clear attitude how to practice of care and treatment (PB)(E3)	0.617	0.730
14. Patients are encouraged to participate actively in their own care and treatment (CP)(E3)	0.588	0.575
15. There is great respect for the patients' privacy and dignity (RP)(C)	0.614	0.745
16. Healthcare staff and other professionals have an understanding of each other's roles (PB)(C)	0.670	0.702
17. The management structure is democratic and inclusive (EIP)(L)	0.723	0.544
18. Relevant information materials are available to patients (e.g. other languages, large print, apps) (E)(C)	0.540	0.532
19. Healthcare staff and patients collaborate on organising personalised care and treatment (CP)(E3)	0.644	0.622
20. Care and treatment are based on thorough assessments (RP)(E3)	0.722	0.725
21. Nurse managers support and encourage staff in being critical of clinical practice (PB)(C)	0.736	0.512
22. Feedback meetings are planned between healthcare staff and patients (CP) (L)	0.521	0.677
23. Nurse managers prioritise staff competence development (EIP)(C)	0.739	0.677
24. The staff use reflective approaches to evaluate and develop practice (e.g. action learning or clinical supervision) (E)(C)	0.671	0.622
25. The organisation's management has great respect for the employees' autonomy (PB)(E3)	0.730	0.612



1			
2			
3			
4	26. Cultural diversity is accepted and welcomed by the staff (RP)(E3)	0.598	0.542
5			
6	27. Healthcare staff have easy access to evidence-based knowledge (EIP)(L)	0.682	0.701
7			
8	28. Patients are able to participate in the assessment (CP)(C)	0.645	0.613
9			
10	29. Healthcare staff have the opportunity to consult specialists (EIP)(L)	0.672	0.591
11			
12	30. Healthcare staff feel well-equipped to develop clinical practice (PB)(E3)	0.776	0.633
13			
14	31. Nurse managers create an environment that promotes the development and exchange of ideas (CP)(C)	0.746	0.716
15			
16	32. Evidence-based clinical guidelines based on patients' experience, clinical experience and research are available (EIP)(E3)	0.666	0.733
17			
18	33. Patients are encouraged to give feedback on care and treatment, as well as departmental culture (CP)(C)	0.582	0.593
19			
20	34. Resources have been allocated for the provision of evidence-based care and treatment (EIP)(C)	0.746	0.779
21			
22	35. The organisation is non-hierarchical (EIP)(E3)	0.647	0.669
23			
24	36. Healthcare staff have a common goal for patient care and treatment (RP)(C)	0.718	0.648
25			
26	37. Structured training programs are available to all healthcare staff (EIP)(E3)	0.651	0.654
27			
28			
29			
30			
31			
32			
33			
34			
35			
36			
37			
38			
39			
40			
41			
42			
43			
44			
45			
46			
47			
48			
49			
50			
51			
52			
53			
54			
55			
56			
57			
58			
59			
60			

Peer Review