



## Reconstructing the last Newfoundland Ice Sheet, Canada.

Mc Henry, M., & Dunlop, P. (2015). *Reconstructing the last Newfoundland Ice Sheet, Canada. Geophysical Research Abstracts Vol. 17, EGU2015-2001-2, 2015.* Abstract from European Geosciences Union General Assembly, Vienna, Austria. <http://adsabs.harvard.edu/abs/2015EGUGA..17.2001M>

[Link to publication record in Ulster University Research Portal](#)

### Publication Status:

Published (in print/issue): 12/04/2015

### Document Version

Publisher's PDF, also known as Version of record

### General rights

The copyright and moral rights to the output are retained by the output author(s), unless otherwise stated by the document licence.

Unless otherwise stated, users are permitted to download a copy of the output for personal study or non-commercial research and are permitted to freely distribute the URL of the output. They are not permitted to alter, reproduce, distribute or make any commercial use of the output without obtaining the permission of the author(s).

If the document is licenced under Creative Commons, the rights of users of the documents can be found at <https://creativecommons.org/share-your-work/licenses/>.

### Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk)



## **Reconstructing the last Newfoundland Ice Sheet, Canada.**

Maureen McHenry and Paul Dunlop

Ulster University, Coleraine, N.Ireland (mc\_henry-m@email.ulster.ac.uk)

The Newfoundland Ice Sheet which formed part of the North American Ice Sheet Complex was situated on the margins of the northwest Atlantic Ocean during the Wisconsinan glaciation (~80ka BP to 10ka BP). This complex consisted of the Laurentide, the Cordilleran and Innuitian Ice Sheets, the Canadian Maritime Provinces Ice Cover and the Newfoundland Ice Sheet (NIS). Although all were confluent at the last glacial maximum, the NIS is known to have supported independent ice centres with advances from the Laurentide Ice Sheet being restricted to Newfoundland's northern and western margins. Given its distinctive position, it is likely the evolution of the NIS through the last glacial cycle was influenced by a number of external and internal drivers including configuration changes in the Laurentide Ice Sheet, ice stream initiation and shutdown, changes in oceanic circulation and fluctuating sea levels and climate signals from the wider Amphi-North Atlantic. As such Newfoundland is a key location for investigating ice sheet response to a number of internal and external forcing mechanisms during glacial cycles. An established technique for reconstructing former ice sheet behaviour is the mapping and spatial analysis of glacial landforms. This provides a valuable record of former ice sheet extent and behaviour through time as well as ice sheet retreat during deglaciation. Here we present new mapping based on our interpretation of SPOT satellite imagery and Digital Elevation Models of the entire Island of Newfoundland as well as swath bathymetric imagery from several locations offshore. Our new database consisting of ~150,000 individually mapped subglacial bedforms that includes drumlins, crag and tails, glacially moulded bedrock lineations and ribbed moraines significantly increases the known landform record in this region. The new database shows Newfoundland has a complex palimpsest landscape that records multiple ice sheet events across the island. Here we report our attempt at unravelling this new record using flowset analysis which separates discrete ice flow patterns into snapshots of ice sheet behaviour through time. Our initial flowset analysis shows the NIS was a dynamic ice sheet which was susceptible to configuration changes throughout the last glacial cycle including ice divide migration, regional configuration changes, ice stream activity and enhanced ice flow caused by marine drawdown.