



Demonstration of a subglacial bedform continuum: Is a unifying formation theory required?

Ely, J., Clark, C. D., Spagnolo, M., Stokes, C., Greenwood, S. L., Hughs, A. L. C., Dunlop, P., & Hess, D. (2014). *Demonstration of a subglacial bedform continuum: Is a unifying formation theory required? Geophysical Research Abstracts Vol. 16, EGU2014-5533, 2014.* Abstract from European Geosciences Union General Assembly, Vienna, Austria.

[Link to publication record in Ulster University Research Portal](#)

Publication Status:

Published (in print/issue): 27/04/2014

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.



Demonstration of a subglacial bedform continuum: Is a unifying formation theory required?

Jeremy C. Ely (1), Chris D. Clark (1), Matteo Spagnolo (2), Chris R. Stokes (3), Sarah L. Greenwood (4), Anna L. C. Hughes (5), Paul Dunlop (6), and Dale Hess (7)

(1) Department of Geography, University of Sheffield, Sheffield, UK (gga08je@shef.ac.uk; c.clark@shef.ac.uk), (2) School of Geosciences, University of Aberdeen, Aberdeen, UK (m.spagnolo@abdn.ac.uk), (3) Department of Geography, Durham University, Durham, UK (c.r.stokes@durham.ac.uk), (4) Department of Geological Sciences, Stockholm University, Stockholm, Sweden (sarah.greenwood@geo.su.se), (5) Department of Earth Science, University of Bergen, Bergen, Norway (anna.hughes@geo.uib.no), (6) School of Environmental Sciences, University of Ulster, Coleraine (p.dunlop@ulster.ac.uk), (7) Department of Earth and Environmental Sciences, The University of Rochester, Rochester, New York, United States (dalehess@gmail.com)

Ice sheet flow organises sediment into undulating landforms called subglacial bedforms, making their genesis pertinent to understanding ice flow mechanics. Currently, there is a lack of consensus regarding their formation. An array of different theories exist which seek to explain specific types of bedform, e.g. flutes, drumlins, ribbed moraine, mega-scale glacial lineations (MSGSL). In contrast, other theories appeal to the hypothesis that these landforms are actually the same phenomena that steadily vary on a scale and shape continuum and that possibly share the same mechanism of formation. Here we test the idea of a continuum by analysing the largest dataset of subglacial bedform morphometrics ever reported ($n \approx 100,000$). Scatter graphs of bedform size and shape metrics (length, width and elongation) reveal two separate clouds of data. The first is comprised of flutes, which form a separate distinct cluster to all other subglacial bedforms. The second, much larger, cloud of data comprises all remaining subglacial bedforms, which form a continuum of size and shape metrics. Hence, it is impossible to quantitatively differentiate previously assigned and commonly used landform labels based on their morphometry. Drumlins merge into MSGSL forming subglacial lineations (aligned with flow direction), and these are adjoined to subglacial ribs (orthogonal to flow) by quasi-circular bedforms which occur within transition zones between ribbed moraine and drumlins. We interpret this as showing that a separate mechanism is required for flute generation and that the other landforms sit along a subglacial bedform continuum of ribs-circles-lineations, consistent with unifying theories for their formation.