



## Surfactants: physicochemical interactions with biological macromolecules

Aguirre-Ramírez, M., Silva-Jimenez, H., Banat, I. M., & Diaz De Rienzo, M. A. (2021). Surfactants: physicochemical interactions with biological macromolecules. *Biotechnology Letters*, 43, 523–535. <https://doi.org/10.1007/s10529-020-03054-1>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Biotechnology Letters

**Publication Status:**  
Published (in print/issue): 31/03/2021

**DOI:**  
[10.1007/s10529-020-03054-1](https://doi.org/10.1007/s10529-020-03054-1)


**Document Version**  
Publisher's PDF, also known as Version of record

**General rights**  
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk).



# Surfactants: physicochemical interactions with biological macromolecules

M. Aguirre-Ramírez · H. Silva-Jiménez · I. M. Banat · M. A. Díaz De Rienzo 

Received: 24 January 2020 / Accepted: 7 December 2020  
© The Author(s) 2021

**Abstract** Macromolecules are essential cellular components in biological systems responsible for performing a large number of functions that are necessary for growth and perseverance of living organisms. Proteins, lipids and carbohydrates are three major classes of biological macromolecules. To predict the structure, function, and behaviour of any cluster of macromolecules, it is necessary to understand the interaction between them and other components through basic principles of chemistry and physics. An important number of macromolecules are

present in mixtures with surfactants, where a combination of hydrophobic and electrostatic interactions is responsible for the specific properties of any solution. It has been demonstrated that surfactants can help the formation of helices in some proteins thereby promoting protein structure formation. On the other hand, there is extensive research towards the use of surfactants to solubilize drugs and pharmaceuticals; therefore, it is evident that the interaction between surfactants with macromolecules is important for many applications which includes environmental processes and the pharmaceutical industry. In this review, we describe the properties of different types of surfactants that are relevant for their physicochemical interactions with biological macromolecules, from macromolecules–surfactant complexes to hydrophobic and electrostatic interactions.

---

M. Aguirre-Ramírez and H. Silva-Jiménez equally contributed to this work.

---

M. Aguirre-Ramírez  
Departamento de Ciencias Químico Biológicas, Instituto de Ciencias Biomédicas, Universidad Autónoma de Ciudad Juárez, Ciudad Juárez, Chihuahua, Mexico

H. Silva-Jiménez  
Área de Oceanografía Química, Instituto de Investigaciones Oceanológicas, Universidad Autónoma de Baja California, Ensenada, Baja California, Mexico

I. M. Banat  
School of Biomedical Sciences, University of Ulster, Coleraine BT52 1SA, Northern Ireland, UK

M. A. Díaz De Rienzo (✉)  
School of Pharmacy and Biomolecular Sciences, Liverpool John Moores University, James Parsons Building 10.05C, Byrom Street, Liverpool L3 3AF, UK  
e-mail: m.a.diaz@ljmu.ac.uk

**Keywords** Surfactants · Macromolecules · Biological systems · Molecular interactions

## List of symbols

- $\gamma_{11}$  Surface tension when two identical phases are considered
- $\gamma_1$  Surface tension of phase 1 i.e. liquid
- $\gamma_2$  Surface tension of phase 2 i.e. solid
- $\gamma_{12}$  Interfacial tension between two different phases
- $w_{11}$  Work of adhesion between two identical phases
- $w_{12}$  Work of adhesion between two different phases

## Introduction

Surfactants are amphiphilic molecules capable of reducing the surface tension between two immiscible phases (Otzen 2017). These molecules are either chemically produced (synthetic surfactants) or based on biological materials (biosurfactants). The reduction of surface tension is due to their amphiphilic properties, as their molecules consist of both hydrophilic and hydrophobic moieties (Li and Lee 2019). The hydrophilic part contains heteroatoms such as oxygen, sulphur, nitrogen and phosphorous, which appear in functional groups such as alcohol, thiol, ether, ester, acid, sulphate, sulfonate, phosphate, amine, amide, etc., while the hydrophobic part is typically a paraffin, cycloparaffin or aromatic hydrocarbon, which may contain halogens. Due to their dual affinity, amphiphilic molecules are not stable either in polar or in organic solvents. To meet both types of affinities, the hydrophilic moiety must be surrounded by a polar solvent, while the hydrophobic moiety must be in contact with an organic solvent. Such conditions exist only between two immiscible phases. The boundary between a condensed phase and a gaseous phase is referred to as a surface, and the boundary between two condensed phases such as two liquids or a liquid and a solid, is referred to as an interphase. Many properties of surfactants depend on this strong affinity for surfaces or interphases (Khan et al. 2015).

There are important properties that characterise each particular system. Surface tension is defined as the work required to increase the area of a surface isothermally and reversibly by unit amount (Ebnesaajjad 2014). Surface tension ( $\gamma$ ) is expressed as surface energy per unit area and alternatively as a force per unit length. If we consider two identical phases the surface tension ( $\gamma_1$ ) can be expressed by Eq. 1:

$$\gamma_1 = \frac{1}{2} w_{11} \quad (1)$$

where  $W_{11}$  represents the work of adhesion between the two identical phases, which is defined as the reversible thermodynamic work required to separate the interface from the equilibrium state of the two phases to a separation distance of infinity.

On the other hand, the interfacial tension between two different phases (1 and 2) can be given by Eq. 2:

$$\gamma_{12} = \gamma_1 + \gamma_2 - w_{12} \quad (2)$$

These characteristics are determinant in terms of the properties of the systems, such as the existence and persistence of emulsions or foams, where surfactants are responsible for the changes (reduction) in surface tension. Surfactants allow the mixing of hydrophilic molecules with hydrophobic ones, through the formation of structures called micelles which allow the association of both types of molecules in a single phase. This compatibility between molecules that do not have a natural affinity is also known as co-solubilisation (Poša et al. 2019) and can be used to establish different applications.

Surfactants are used in a wide range of industrial applications (Banat and Thavasi 2018). In agriculture, for example, phytosanitary agents are applied in the form of aerosol (surfactant) which, sometimes, contains a dispersed organic phase (emulsifier) to dissolve herbicides and insecticides (Marquez et al. 2018). While in food products, they contribute to the conditioning of creams, suspensions, emulsions, soluble or dispersible powders (Kralova and Sjöblom 2009). In mining processes, they play an important role in the flotation and leaching of metals like iron, zinc, uranium (Asselin and Ingram 2014; Diaz et al. 2015); as well as in the textile industry to improve the performance of different operations and to provide particular properties to the finished products (Pacífico and Giers 1995; Proffitt and Patterson 1988). In the oil industry, they have been used to help to solve problems caused by drilling operations to the conditioning of the finished products; in fact, extracted crude oil reaches the surface in the form of a water-in-oil emulsion, which makes it essential to remove or separate the water content (Marquez et al. 2019).

Chemical surfactants are derived from non-biodegradable components, and in some cases can cause serious problems to the environment, such as: (1) the formation of foams which inhibit or paralyze natural (or artificial) purification processes, concentrate impurities and can spread bacteria or viruses; (2) the increase of phosphate content in basins, from polyphosphates that are used in combination with surfactants (Santos et al. 2016).

Given the problems caused by synthetic surfactants, different studies have been carried out over the past years, seeking to find alternative products compatible with the environment and have demonstrated the feasibility of producing these compounds from

microorganisms (Akbari et al. 2018). Most microbial biosurfactants are typically biodegradable, biocompatible and have stable activities under extreme environmental conditions (Naughton et al. 2019). Hence the interest to study their production from fungi and bacteria, among which the genera *Bacillus* and *Pseudomonas* stand out. Many of these biosurfactants produced by *Pseudomonas aeruginosa* have been characterized and studied as agents capable of removing hydrophobic compounds from soil (Geetha et al. 2018), antimicrobials and biofilm disruptors (Elshikh et al. 2017; Diaz De Rienzo et al. 2016; Ceresa et al. 2020). Although the physicochemical properties of (bio) surfactants have been well documented through the years (Mankowich 1953; Behrens 1964; Van Os et al. 1993; Patino et al. 2007; Morais et al. 2017), their interaction with biological components has had less focus. This review therefore focuses on the properties of surfactants that are relevant for their physicochemical interactions with biological systems (Fig. 1), and when possible compare them with their biological counterparts.


### Surfactant–protein interactions

The study of the interactions between surfactants, both synthetic and microbial (biosurfactants), with proteins is of great interest in various biotechnology fields and industries such as food, cosmetics, pharmaceutical, biomedical, and environmental (Lee et al. 2011; Otzen 2011; Tucker et al. 2014; Malik 2015). In the biomedical industry, protein–surfactant systems are used for the production of hydrogels (Afinjuomo et al. 2019; Castelli et al. 2008). The hydrogels form the

base of fibrous proteins such as fibroin, which are used for tissue regeneration and drug delivery (Park et al. 2014; Dubey et al. 2018; Ohadi et al. 2020).

There are three main forces that drive the protein–surfactant interaction: (1) electrostatic, (2) hydrophobic and (3) Van der Waals (Mackie and Wilde 2005; Li and Lee 2019). The dominant interaction is determined by the nature of both molecules and their concentration (Mehan et al. 2015; Li and Lee 2019). These molecular interactions have an influence on the native structure of proteins promoting or preventing denaturation, aggregation and loss of enzymatic activity among other factors (Mehan et al. 2015). Surfactants of biological origin have an advantage over synthetic surfactants in terms of their ability to prevent denaturation of proteins and a reduction in their aggregation (Otzen 2011, 2017).

The protein–surfactant systems mainly studied are those that contain globular proteins such as bovine serum albumin (BSA),  $\alpha$ -lactoglobulin and  $\beta$ -glucosidase. In contrast, very few studies have been performed exploring the fibrous protein–surfactant systems. Type I collagen, silk fibroin, and keratin are fibrous proteins that have been studied in combination with ionic and non-ionic surfactants (Maldonado et al. 1991; Mandal and Kund 2008; Kezwon et al. 2016; Kezwoń and Wojciechowski 2016; Pan et al. 2016; Park et al. 2014; Dubey et al. 2018). A few studies suggest that the molecular interactions presented by fibrous proteins (collagen, fibroin, keratin) in combination with ionic and non-ionic surfactants are similar to the globular protein–surfactant systems (Lee et al. 2011; Khan et al. 2015; Kezwon et al. 2016; Kezwoń and Wojciechowski 2016; Pan et al. 2016).



**Fig. 1** Illustrative summary of the main types of interactions between (bio)surfactants and macromolecules

Type I collagen interacts with Sodium Dodecyl Sulphate (SDS), Cetyl Trimethyl Ammonium Bromide (CTAB), and Triton X-100 through hydrophobic and electrostatic molecular interactions. The predominance of a particular molecular interaction depends on the type of surfactant, i.e. surfactants could produce changes in collagen secondary structure (Maldonado et al. 1991; Kezwon et al. 2016; Kezwoń and Wojciechowski 2016).

The main physical parameters that have an effect on the surfactant–protein interactions are: (a) the surfactant concentration; (b) the chemical nature of surfactant (ionic or non-ionic surfactants); and (c) the secondary structure of the protein ( $\alpha$ -helix and  $\beta$ -sheets) (Díaz et al. 2003; Malik 2015).

#### Surfactant concentration


The effect on stabilization or destabilization mediated by a surfactant is dependent on the concentration of the surfactant (Mehan et al. 2015). In that sense, many surfactants (biological and synthetic ones), usually promote protein stabilization at concentrations far below Critical Micelle Concentration (CMC), while at concentrations higher than the CMC there is an opposite effect, they promote denaturation, aggregation, as well as loss of biological function of proteins (Díaz et al. 2003; Otzen 2011; Malik 2015). In general, the binding of the surfactant to the protein is carried out in three phases. In the binding phase (phase I), individual surfactant molecules bind to the protein without causing any structural change, and electrostatic interactions dominate over hydrophobic ones. In the cooperative phase (phase II), the increase in the surfactant concentration reaches a sub-CMC levels, triggering the formation of the hydrophobic clusters that start to bind to the hydrophobic regions of proteins leading to their denaturation and changes in the secondary structure. In this phase, hydrophobic interactions dominate over electrostatic; in addition, the unfolding process increases linearly (Otzen 2011; Malik 2015). Finally, the saturation phase (phase III) is where the protein binding sites are already saturated. In this phase, there are free surfactant molecules that interact with the protein-bound micelles and no longer cause further changes (Malik 2015).

#### Chemical nature of surfactants

Surfactants can be divided into two groups according to their chemical composition: ionic and non-ionic. The ionic surfactants, according to their charge, can be anionic or cationic (Otzen 2011; Khan et al. 2015). The hydrophilic group of the surfactant affects the stability of the protein because it can tightly bind to the protein causing its denaturation and contributes to the solubilization of the membrane proteins (Mehan et al. 2015). Anionic surfactants are typically protein-denaturing agents (Khan et al. 2015). Among the anionic surfactants, SDS is well known for having strong electrostatic interactions with proteins (Deep and Ahluwalia 2001; Otzen et al. 2009; Hansted et al. 2011; Otzen 2011). These interactions are generated between the positively charged amino acids present in the primary structure of the protein along with the interactions of the hydrocarbon chains of the surfactant, and the aliphatic regions of the amino acids arginine (Arg) and lysine (Lys) (Otzen et al. 2009).

Such properties have been used in some protein separation and/or solubilisation techniques. The interaction between SDS and several globular proteins has been previously reported, i.e. the denaturing effect of SDS on  $\alpha$ -lactalbumin occurs in different stages depending on the concentration of the surfactant. In the early stages, SDS monomers bind to the protein to form groups up to a critical concentration that results in the start of the denaturation process (Fig. 2). The binding of more monomers results in the loss of the secondary structure of the protein (Otzen et al. 2009). In the case of  $\beta$ -lactoglobulin, SDS has an opposite effect to the one observed with  $\alpha$ -lactalbumin, since this amphiphilic molecule reduces the aggregation of the protein at concentrations well below its CMC (Hansted et al. 2011).

Compared to anionic surfactants, cationic surfactants have a milder protein destabilization effect (Khan et al. 2015). These ionic surfactants interact with amino acids whose side chains are usually negatively charged like aspartate (Asp) and glutamate (Glu) (Otzen 2011). For example, Khan et al. (2019) reported that the interactions between CTAB and Hen egg white lysozyme are very hydrophobic and weakly electrostatic, which do not cause a change in the secondary structure of the protein but do cause a negative effect on the tertiary structure.



**Fig. 2** Representative scheme of the denaturation effect promoted by SDS over  $\alpha$ -lactalbumin. SDS monomers bind to the protein starting the denaturation process; at a high concentration of SDS monomers, the secondary structure of the protein is lost

In the case of non-ionic surfactants (i.e., dodecyl maltoside, polysorbates), they commonly minimize or prevent protein aggregation (Lee et al. 2011; Otzen 2011). According to various studies, the molecular interactions between proteins and non-ionic surfactants are very weak and the union of the biomolecule with the non-ionic surfactant is driven by hydrophobic interactions, which results in a tendency to solubilize proteins. These surfactants are used in the food industry and have biomedical applications in drug formulations (Lee et al. 2011; Campos et al. 2013; Tucker et al. 2014). Non-ionic surfactants usually have ethoxylate groups that interact with the hydrophobic moieties of proteins, exposing the hydrophilic groups present in both molecules, which results in the increase of the hydrophilicity of the non-ionic surfactant–protein complex, thereby reducing the aggregation of proteins (Rudolph and Jones 2002; Ruiz-Peña et al. 2010; Lee et al. 2011; Tucker et al. 2014). The chemical structure of this type of surfactant plays an important role in promoting or preventing protein denaturation, even if the structural differences are minor. Tween type surfactants (ethoxylated polysorbates) vary in the length of the fatty acid hydrocarbon chain and interact differently with BSA, as seen in the number of surfactant molecules that are able to bind to the protein as well as the type of binding (Ruiz-Peña et al. 2010).

Another type of surfactants known as dimeric or Gemini surfactants are constructed of two monomers of surfactants which are joined by a spacer close to the hydrophilic heads (Sinha et al. 2016). Despite their importance in several industrial fields, studies of Protein-Gemini surfactants interactions are limited, compared with those conducted with single chain surfactants (Sinha et al. 2016; Parray et al. 2018; Akram et al. 2019). Several studies have revealed that some interaction mechanisms of these new generation of surfactants with proteins are shared with their

corresponding monomers differing in the effects that they induce in the biomolecule, ranging from having stronger molecular interactions than their monomeric counterpart to changes or stabilization in the secondary and tertiary structures of proteins (Sinha et al. 2016; Sonu et al. 2017; Akram et al. 2019). Comparative studies of the interaction of BSA with the cationic surfactant Dodecyl Trimethyl Ammonium Bromide (DTAB) and with three Gemini-surfactants of the bis(dimethyldodecylammonium bromide) family; butanediyl-1,4-bis(dimethyldodecylammonium bromide) (12–4-12,2Br –), 2-butanol-1,4-bis(dimethyldodecylammonium bromide) (12–4(OH)-12,2Br –), 2,4-dibutanol-1,4-bis(dimethyldodecylammonium bromide) (12–4(OH)<sub>2</sub>-12,2Br –), showed that at lower concentrations of the surfactant the interaction in the surfactant–protein complex is managed by electrostatic forces and while the concentration of the surfactant increases.

The union of the protein with the surfactant is hydrophobic in nature, which is stronger with the Gemini-surfactant, causing greater denaturation of BSA compared to DTAB, which suggests that the spacer between the two monomers plays an important role (Sinha et al. 2016). Sonu et al. (2017) conducted a study on the effect of surfactant spacers [12-8-12, 2Br-], [12-4-12, 2Br-] and [12-4 (OH) -12, 2Br-] on the interaction with BSA and reported that the more hydrophobic the spacer is, the lower is the reduction in the number of  $\alpha$ -helices and denaturing effects. Akram et al. (2019) on the other hand, analysed the interaction of the BSA model protein with three members of a family of Gemini Cm-E20-Cm surfactants and demonstrated that the binding of these dimeric surfactants with the protein is considerably strong, without causing a significant loss of  $\alpha$ -helix (3–4%), keeping the secondary and tertiary structure of the BSA virtually intact. Other authors have reported that the effect caused by these Gemini-surfactants on the

various model proteins may be subject to changes at different temperatures, pH concentrations, ionic strength, and surfactant concentrations, among others (Faustino et al. 2009).

### Secondary structure of proteins

In some cases, the secondary structure of a protein could have an effect on the ability of a surfactant to promote its aggregation or denaturation activities, without necessarily being a specific surfactant–protein interaction. Zaragoza et al. (2012) showed that when the trehalolipid biosurfactant produced by a *Rhodococcus* sp. is present at a concentration lower than CMC, proteins with a high content of  $\alpha$ -helix in the secondary structure such as BSA and cytochrome *c* (Cyt-*c*) showed resistance to thermal unfolding and there was no alteration of the secondary structure. In addition, Isothermal Titration Calorimetry (ITC) investigations demonstrated that the interactions between trehalolipids and both proteins are not specific, suggesting the involvement of hydrophobic domains of proteins (Zaragoza et al. 2012). However, the biosurfactant mannosylerythritol lipid-A (MEL-A) has a different influence on the enzyme  $\beta$ -glucosidase. At CMC values, this biosurfactant promotes a secondary structure changes of  $\beta$ -glucosidase, causing a decrease in  $\beta$ -sheets content and an increase in  $\alpha$ -helices,  $\beta$ -turn, and random coil. These structural changes cause  $\beta$ -glucosidase to acquire thermal stability by increasing its midpoint temperature ( $T_m$ ) and unfolding enthalpy (Fan et al. 2018).

The above can be explained in thermodynamic and structural terms. On the one hand, at CMC values, MEL-A forms micelles, thereby increasing hydrophobic interactions. Thermodynamic data obtained by ITC, support the hypothesis that weak hydrophobic interactions are responsible for the union of MEL-A and  $\beta$ -glucosidase. On the other hand, the stability gained by  $\beta$ -glucosidase at CMC values can be given by the enzyme's secondary structural changes. The increase of  $\alpha$ -helix content is a potential factor which promotes, (1) the exposure of hydrophobic regions to amino acid residues that interact hydrophobically, (2) hydrogen bond formation with fatty acid chains, and (3) hydroxyl groups of glycosidic residues (Otzen 2011; Fan et al. 2018).

Based on various analytical methods, Zhang and Li (2018) reported that surfactin, a biosurfactant of the

lipopeptide type, induces changes in the conformations of the alkaline protease secreted by *Bacillus* sp., which results in weak hydrophobic interactions, hydrogen bonds and some electrostatic interactions. In addition, they found that the enzymatic activity of the alkaline protease may be affected positively or negatively at low or high concentrations of surfactin, respectively. In the first case, the low concentration of surfactin in the aqueous medium, allows the biosurfactant molecule to interact with the alkaline protease as a cofactor, thus causing an increase in enzymatic activity, while at high concentrations of surfactin, a decrease in enzymatic activity occurs. This is because the hydrophobicity of the alkaline protease is decreased by the high concentration of biosurfactant molecules present in the solution. Finally, the cases analysed in this review on the interactions between different surfactants with a model protein reveal that they are quite diverse, where the physicochemical characteristics of the interacting molecules play an essential role. Molecular interaction studies using various biophysical techniques, will allow us to understand the basis of interaction between surfactants and proteins.

### Surfactant–lipid interactions

The phase behaviour between surfactants–water and lipid–water is well documented (Chernik 2000; Koynova and Tenchov 2001; Ebnesajjad 2006), however the interaction between surfactants and lipids is not well reported with most studies have been carried out on temperature and enthalpy variables without a detailed description of the mechanisms involved (Koynova and Tenchov 2001). Surfactants are widely used as molecular tools, especially in studies of membrane biology for biomembrane solubilization, based on their ability to form mixed micelles with lipids and proteins (Koynova and Tenchov 2001) and as a liposome-mediated drug delivery system (Bnyan et al. 2018). Liposomes have been used as a model of biological membranes for a long time, due to their phospholipid structure. The structure of phospholipids has a hydrophilic head group and a hydrophobic tail group. When dispersed in an aqueous solution, the head is attracted by water, and the tail, including a long hydrocarbon chain, is repelled by water promoting the

formation of vesicles (Stryer 1981; Dua et al. 2012; Gunay and Ozer 2018).


The interaction between lipids and surfactants is derived in a different numbers of model systems (Helenius and Simons 1975; Lichtenberg et al. 1983). All these models show a general scheme for the interaction between lipids and surfactants (which displays the transition from vesicles to mixed micelles) and is described as a three-stage model (Fig. 3). The first stage is where the surfactant partition between the lipid bilayers and the aqueous phase and start reaching a level where the bilayers break into micelles; the second phase is where there is a mix between micelles and bilayers in a co-existent state and the last phase is characterized by an increase of the surfactant concentration leading to a phase where all the bilayers are solubilized and only lipid-rich micelles are present (Lichtenberg et al. 2013; Pizzirusso et al. 2017).

There are different studies that show the three-stage model applied to biological membranes, including homogenous phospholipids systems (phosphatidylcholine and phosphatidylserine),  $\text{Ca}^{2+}$ -ATPase membranes (Le Maire et al. 2000) and liposomes prepared from SR lipid (Langner and Hui 2000). The solubilisation of membranes generally occurs via the uptake of non-micellar surfactants monomers, which is why when a surfactant is added to solubilize a membrane preparation, if the surfactant concentration is below their CMC, then it is just the monomer fraction that interact with the biological membrane.

When it comes to the study of biosurfactants and membrane lipids interactions, few studies have been reported on molecular interactions (Ortiz et al. 2009; Aranda et al. 2007; Rodrigues et al. 2006; Malaspina

et al. 2017). The effect of trehalose lipids on membrane phospholipids was reported by Ortiz et al. (2008) showing that the biosurfactants exhibit a dehydrating effect on the interfacial region of saturated phosphatidylethanolamines promoting the formation of unsaturated phosphatidylethanolamines. The same research group evaluated the effect of trehalose lipid produced by *Rhodococcus* sp. on the structural properties of dimyristoyl phosphatidylserine (DMPS) membranes. They have showed that the biosurfactant incorporates into the DMPS membranes and increases the fluidity of the phosphatidylserine acyl chains making changes in the environment of the polar head group and, as a consequence, decreases the interfacial tension of the membrane, thereby decreasing the motional freedom of the phospholipids (Ortiz et al. 2009).

One of the most studied biosurfactant in terms of their effect on the plasma membrane is the Iturin produced by *Bacillus subtilis*. Iturin is an effective antifungal compound and its mechanisms of action is related to the disruption of the biological membrane by the formation of small vesicles and their aggregation in yeast cells (Peypoux et al. 1994; Rodrigues et al. 2006). Iturin was shown to pass through the cell wall and disrupt the plasma membrane with the formation of small vesicles and the aggregation of intramembranous particles, interacting with the nuclear membrane and probably with membranes of other cytoplasmic organelles affecting the morphology and membrane structure of yeast cells (Thimon et al. 1995). Recently, the studies in molecular surfactant-like peptides and lipids has become more focused and significant due to their excellent properties, such as versatility, biocompatibility and medicinal properties



**Fig. 3** Surfactants–lipids interaction: the three-stage model. Stage I: Surfactant molecules approach a bilayer. Stage II: Combination of micelles and lipid/surfactant aggregates. Stage III: Mixed micelles formation



(Cui et al. 2010; Hosseinkhani et al. 2013; Dehsorkhi et al. 2014; Du and Stenzel 2014; Malaspina et al. 2017; Doostmohammadi et al. 2019).

An important class of amphiphilic peptides called surfactant-like peptides (SLPs), present an intrinsic difference that can lead to different physical consequences namely composition and tail structure (Malaspina et al. 2017). Unlike conventional surfactants whose hydrophobic tails interact in all directions through hydrophobic interactions, the amphiphilic peptide tail contains not only hydrophobic groups but also hydrophilic sites (Colherinhas and Fileti 2014). This feature allows the SLPs to stabilize nanostructures in one direction through hydrophobic interactions and in the orthogonal direction by hydrogen bonds. These hydrogen bonds associated with hydrophobic interactions can stabilize at a high level, complex secondary structures such as helices and sheets. On the other hand, conventional lipids/surfactants with antimicrobials properties (Chen et al. 2010, 2012; Albada et al. 2012; Gaspar et al. 2013) are usually organized into micelles, vesicles, and nanotubes (Colherinhas and Fileti 2014; Malaspina et al. 2017). To understand the interaction between (bio)surfactants and lipids, it is necessary to be aware of the hydrodynamics of the molecules involved, their amphiphilic properties and how they play an important role when it comes to biological membranes. Nanoparticle models and the study of their properties could help us to understand the molecular basis of these interactions, which have remained unknown.

### Surfactant–polysaccharide interactions

Polysaccharides are monosaccharide (homo or hetero) built up biopolymers mainly produced by plants. Similar to surfactants, they could be classified based on their charge as non-ionic (o), cationic (+), and anionic (−) polymers (Kwak 1998). Polysaccharides and surfactant interactions are important to develop (a) emulsifiers; (b) flocculating agents; (c) stabilizing colloids; (d) or rheology controllers (Holmberg et al. 2002) in food, medicine and environmental applications.

Electrostatic, hydrophobic, dipole–dipole, and hydrogen bonding interactions along with the surfactant and polysaccharide characteristic are the main factors that affect the Polymer–Surfactant Systems

(PSS) (Grządka et al. 2019). These interactions have been summarised in Table 1 (Bao et al. 2008). These authors studied the interactions of ionic surfactants (SDS and CTAB) with neutral, positively, and negatively charged polysaccharides [Methyl cellulose (MC), chitosan (CS) and  $\kappa$ -carrageenan (KC)], respectively.

According to the surfactant–polysaccharide combination, molecular interactions change. Therefore, strong hydrophobic and weak ion–dipole interactions are present in MC–SDS mixture. Moreover, in KC–SDS and CS–SDS, ionic interactions drive the binding process between surfactant and the polymer. Hydrophobic interactions are weak in KC–SDS, while in CS–SDS, polymer hydrophobic moieties interact with alkyl chains of the SDS. In the case of CTAB with MC and CS, only hydrophobic interactions are present, and strong electrostatic interactions allow binding between KC and CTAB.

In the case of non-ionic polysaccharide and anionic surfactant, as ethyl hydroxyethyl cellulose (EHEC) and SDS, respectively, the hydrophobic interaction between the polymer and SDS alkyl chain drives their association. Accordingly, SDS plays an important role because its presence or absence promotes the extent of EHEC–SDS cluster formation. For example, if SDS concentration is below the critical aggregation concentration (CAC), surface tension is reduced depending on SDS molecules, but when SDS concentration increases to at or above the CAC, EHEC adsorption is accelerated. In diluted solutions, the surface activity is strong (12 ppm of EHEC and 2 mM SDS), making this PSS a vehicle for drug delivery (Nahringerbauer 1997).

Cationic surfactants such as DTAB, MTAB, and CTAB, interact with cellulose in the water interface. These cationic surfactants contain a different number of  $-\text{CH}_2-$  groups in the alkyl chain, and their CMC varies with respect to alkyl chain length (CTAB > MTAB > DTAB). The chain length of this kind of cationic surfactants influences interaction behaviour with non-ionic polysaccharides such as cellulose. For example, CTAB–cellulose interaction is driven by hydrophobic interactions, while electrostatic interactions are very significant in interactions of MTAB and DTAB with cellulose, respectively.

In the case of interactions of polysaccharides such as dextrin and carboxymethylcellulose with cationic surfactant groups (DTAB, MTAB, CTAB), the

**Table 1** Interactions between methyl cellulose, chitosan and  $\kappa$ -carrageenan with ionic surfactants, SDS and CTAB (Bao et al. 2008)

Polysaccharide	Surfactant	Interaction		
		Hydrophobic	Electrostatic	Ion–dipole
Methyl cellulose	SDS	Strong		Weak
Chitosan		Medium	Strong	
$\kappa$ -Carrageenan		Weak		
Methyl cellulose	CTAB	Strong		
Chitosan		Medium		
$\kappa$ -Carrageenan			Strong	

**Table 2** Interaction of cetyl trimethyl ammonium bromide family (CnTAB) with cellulose nanocrystals (C = carbon number in alkyl chain of surfactant) (Brinatti et al. 2016)

	CnTAB interaction with cellulose nanocrystals	Micelle formation	Flocculation
C = 12	Electrostatic	High concentration	
C = 14	Electrostatic–hydrophobic	Low concentration	High concentration
C = 16	Electrostatic–hydrophobic	Low concentration	High concentration

behaviour is different, for the interaction between dextrin and CTAB, hydrophobicity drives the interaction while in the case of carboxymethylcellulose and CTAB, electrostatic interactions are very significant (Biswas and Chatteraj 1997a, b).

Another example of PSS with an anionic surfactant, sodium stearyl lactylate (SSL), an anionic surfactant and  $\kappa$ -carrageenan (KC) polymer, both of which are important in the food industry, have a different behaviour in solutions and gels. SSL changes KC conformation due to electrostatic interactions and hindrance. In the gelation process (melting process), KC suffers a coil helix transition and, finally, helix–helix aggregation, modifying its melting enthalpy. SSL hinders KC helix–helix aggregation. But, at a high concentration of surfactant, SSL forms micelles (solutions and gels). The combination of hindrance and electrostatic repulsion promote conformational changes in KC, both in solutions and in gels. In solution, enthalpy decreases continuously at high SSL concentration range, while in gels, this parameter decreases at a specific SSL concentration (Ortiz-Tafoya et al. 2018).

In other cases, the interaction between a polysaccharide and surfactant depends on the alkyl chain length of the tensioactive molecule. Such is the case of CTAB homologues (CnTAB, where  $n$  is a carbon number in alkyl chain of surfactant) with cellulose

nanocrystals, a negatively charged polysaccharide. When  $n = 12$  and the surfactant concentration is high, electrostatic interactions are present and micelle formation occurs, while at  $n = 14–16$  and a low surfactant concentration, micelles are formed, and flocculation process occurs at high CnTAB concentration (Table 2). These processes are driven in first instance by electrostatic interactions and by the hydrophobic interactions (Brinatti et al. 2016).

#### Polysaccharides–biosurfactants interactions

Some biosurfactants contain sugars in their structure such as glycolipids (e.g. rhamnolipids) and also interact with polysaccharides. In the food and pharmaceutical industries, pickering/stabilizing high internal phase emulsions (HIPEs) are very important as they are used in bioactive delivery. In these HIPEs three kinds of molecules interact: proteins–polysaccharides–biosurfactants. For example, zein–propylene glycol alginate mixed with rhamnolipids stabilize pickering emulsion in the oil-in-water interface. This emulsion system is formed by a 3D network of adsorbed and non-adsorbed particles, however the basis of molecular interactions amongst these molecules is unclear (Dai et al. 2019).

## Conclusions

Recent works in this area highlight the importance of the interactions between surfactants and macromolecules and their role in biological membranes. The structures that form in solution are driven by molecular interactions. There are three main forces that drive the protein–surfactant interactions: electrostatic, hydrophobic, and Van der Waals, while the dominant interaction is controlled by the characteristics of both molecules and their concentration. The interactions between lipids and surfactants are described as a three-stage model, starting with the surfactant partition between the lipid bilayers and the aqueous phase, reaching a level where the bilayers break into micelles and ending with the solubilization of bilayers. The characteristics of PSS, such as polysaccharides–surfactants, can be controlled through the molecule design and their charge i.e. presence of electrostatic interaction at opposite charge PPS where hydrophobic interactions are predominant in o/– and o/+ PPS, and where o/– interaction is stronger than o/+ PSS; these are some of the most powerful parameters to take into account in order to obtain the desired structures to be used for different applications. By modifying the interaction type and strength, as well as the concentrations of the molecules involved, the final product can be used for a wide variety of industrial formulations.

**Acknowledgements** The authors acknowledge the funding support from the Faculty of Science, Liverpool John Moores University ECR Fellowship 2018–2019. We also acknowledge to CAC Biología Celular y Molecular, Universidad Autónoma de Ciudad Juárez for its host facilities.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Afinjuomo F, Fouladian P, Parikh A et al (2019) Preparation and characterization of oxidized inulin hydrogel for controlled drug delivery. *Pharmaceutics* 11:356
- Akbari S, Abdurahman NH, Yunus RM et al (2018) Biosurfactants—a new frontier for social and environmental safety: a mini review. *Biotechnol Res Innov* 2(1):81–90
- Akram M, Ansari F, Bhat IA et al (2019) Probing interactions of bovine serum albumin (BSA) with the biodegradable version of cationic gemini surfactant. *J Mol Liq* 276:519–528
- Albada HB, Prochnow P, Bobersky S et al (2012) Tuning the activity of a short Arg-Trp antimicrobial peptide by lipidation of a C- or N-terminal lysine side-chain. *ACS Med Chem Lett* 3(12):980–984
- Aranda FJ, Espuny MJ, Marqués A et al (2007) Thermodynamics of the interaction of a dirhamnolipid biosurfactant secreted by *Pseudomonas aeruginosa* with phospholipid membranes. *Langmuir* 23:2700–2705
- Asselin S, Ingram J (2014) Uranium leaching from contaminated soil utilizing rhamnolipid, EDTA, and citric acid. *Appl Environ Soil Sci* 2014:462514
- Banat IM, Thavasi R (2018) Microbial biosurfactants and their environmental and industrial applications. CRC Press, Boca Raton, USA
- Bao H, Lin L, Gan LH et al (2008) Interactions between ionic surfactants and polysaccharides in aqueous solutions. *Macromolecules* 41:9406–9412
- Behrens R (1964) The physical and chemical properties of surfactants and their effects on formulated herbicides. *Weeds* 12:255
- Biswas SC, Chatteraj DK (1997a) Polysaccharide–surfactant interaction. 1. Adsorption of cationic surfactants at the cellulose–water interface. *Langmuir* 13:4505–4511
- Biswas SC, Chatteraj DK (1997b) Polysaccharide–surfactant interaction. 2. Binding of cationic surfactants to carboxymethyl cellulose and dextrin. *Langmuir* 13:4512–4519
- Bnyan R, Khan I, Ehtezazi T et al (2018) Surfactant effects on lipid-based vesicles properties. *J Pharm Sci* 107(5):1237–1246
- Brinatti C, Huang J, Berry RM et al (2016) Structural and energetic studies on the interaction of cationic surfactants and cellulose nanocrystals. *Langmuir* 32:689–698
- Campos JM, Montenegro Stamford TL, Sarubbo LA, de Luna JM, Rufino RD, Banat IM (2013) Microbial biosurfactants as additives for food industries; a review. *Biotechnol Prog* 29(5):1097–1108
- Castelli F, Sarpietro MG, Micieli D et al (2008) Differential scanning calorimetry study on drug release from an inulin-based hydrogel and its interaction with a biomembrane model: pH and loading effect. *Eur J Pharm Sci* 35(1–2):76–85
- Ceresa C, Fracchia L, Williams M, Banat IM et al (2020) The effect of sophorolipids against microbial biofilms on medical-grade silicone. *J Biotechnol* 309:34–43
- Chen C, Pan F, Zhang S, Hu J et al (2010) Antibacterial activities of short designer peptides: a link between propensity for nanostructuring and capacity for membrane destabilization. *Biomacromolecules* 11(2):402–411

- Chen C, Hu J, Zhang S et al (2012) Molecular mechanisms of antibacterial and antitumor actions of designed surfactant-like peptides. *Biomaterials* 33(2):592–603
- Chernik GG (2000) Phase studies of surfactants–water systems. *Curr Opin Colloid Interface Sci* 4:381–390
- Colherinhas G, Fileti E (2014) Molecular dynamics study of surfactant-like peptide based nanostructures. *J Phys Chem B* 118(42):12215–12222
- Cui H, Webber M, Stupp SI (2010) Self-assembly of peptide amphiphiles: from molecules to nanostructures to biomaterials. *Biopolymers* 94(1):1–18
- Dai L, Yang S, Wei Y et al (2019) Development of stable high internal phase emulsions by pickering stabilizations: utilization of zein-propylene glycol alginate-rhamnolipid complex particles as colloidal emulsifiers. *Food Chem* 275:246–254
- Deep S, Ahluwalia JC (2001) Interaction of bovine serum albumin with anionic surfactants. *Phys Chem Chem Phys* 3:4583–4591
- Dehsorkhi A, Castelletto V, Hamley IW (2014) Self-assembling amphiphilic peptides. *J Pept Sci* 20(7):453–467
- Díaz X, Abuin E, Lissi E (2003) Quenching of BSA intrinsic fluorescence by alkylpyridinium cations: its relationship to surfactant–protein association. *J Photochem Photobiol* 155:157–162
- Diaz MA, De Ranson IU, Dorta B et al (2015) Metal removal from contaminated soils through bioleaching with oxidizing bacteria and rhamnolipid biosurfactants. *Soil Sediment Contam Int J* 24:16
- Diaz De Rienzo MA, Kamalanathan ID, Martin PJ (2016) Comparative study of the production of rhamnolipid biosurfactants by *B. thailandensis* E264 and *P. aeruginosa* ATCC 9027 using foam fractionation. *Process Biochem* 51:820–827
- Doostmohammadi M, Ameri A, Mohammadinejad R, Banat IM, Ohadi M, Dehghannoudeh G (2019) Hydrogels for peptide hormones delivery: therapeutic and tissue engineering applications. *Drug Des Dev Ther* 13:3405–3418
- Du AW, Stenzel MH (2014) Drug carriers for the delivery of therapeutic peptides. *Biomacromolecules* 15(4):1097–1114
- Dua JS, Rana AC, Bhandari AK (2012) Liposome: methods of preparation and applications. *Int J Pharm Stud Res* 3(2):14–20
- Dubey P, Sugam K, Ravindranathan S et al (2018) pH dependent sophorolipid assemblies and their influence on gelation of silk fibron protein. *Mater Chem Phys* 23:9–16
- Ebnesajjad S (2006) Surface treatment of materials for adhesive bonding. In: Ebnesajjad S (ed) *Surface tension and its measurement*, 2nd edn. Elsevier, USA, pp 9–28
- Ebnesajjad S (2014) Surface treatment of materials for adhesive bonding. In: Ebnesajjad S (ed) *Surface tension and its measurement*, 2nd edn. Elsevier, USA, pp 7–24
- Elshikh M, Moya-Ramírez I, Moens H et al (2017) Rhamnolipids and lactic acid sophorolipids: natural antimicrobial surfactants for oral hygiene. *J Appl Microbiol* 123:1111–1123
- Fan L, Xie P, Wang Y et al (2018) Biosurfactant–protein interaction: influences of mannosylerythritol lipids-A on  $\beta$ -glucosidase. *J Agric Food Chem* 66:238–246
- Faustino CMC, Calado ART, García-Río L (2009) Gemini-surfactant–protein interactions: effect of pH, temperature, and surfactant stereochemistry. *Biomacromolecules* 10:2508–2514
- Gaspar D, Veiga AS, Castanho MARB (2013) From antimicrobial to anticancer peptides. A review. *Front Microbiol* 4:294
- Geetha SJ, Banat IM, Joshi SJ (2018) Biosurfactants: production and potential applications in microbial enhanced oil recovery (MEOR). *Biocatal Agric Biotechnol* 14:23–32
- Grządka E, Matusiak J, Stankevič M (2019) Interactions between fluorocarbon surfactants and polysaccharides. *J Mol Liq* 283:81–90
- Gunay SM, Ozer Y (2018) Liposomes and micelles as nanocarriers for diagnostic and imaging purposes In: *Design of nanostructures for theranostics applications*. William Andrew Publishing, USA
- Hansted JG, Wejse PL, Bertelsen H et al (2011) Effect of protein–surfactant interactions on aggregation of  $\beta$ -lactoglobulin. *Biochim Biophys Acta* 1814:713–723
- Helenius A, Simons K (1975) Solubilization of membranes by detergents. *Biochim Biophys Acta* 415:29–79
- Holmberg K, Joensson B, Kronberg B et al (2002) Surfactants and polymers in aqueous solution. Wiley, Chichester, UK
- Hosseinkhani H, Hong PD, Yu DS (2013) Self-assembled proteins and peptides for regenerative medicine. *Chem Rev* 113:4837–4861
- Kezwon A, Góral I, Fraczyk T et al (2016) Effect of surfactants on surface activity and rheological properties of type I collagen at air/water interface. *Colloid Surf B* 148:238–248
- Kezwoń A, Wojciechowski K (2016) Collagen–surfactant mixtures at fluid/fluid interfaces. *Colloid Surf A* 509:390–400
- Khan TA, Mahler HC, Kishore RSK (2015) Key interactions of surfactants in therapeutic protein formulations: a review. *Eur J Pharm Biopharm* 97:60–67
- Khan JM, Malik A, Ahmed A et al (2019) Effect of cetyltrimethylammonium bromide (CTAB) on the conformation of a hen egg white lysozyme: a spectroscopic and molecular docking study. *Spectrochim Acta A Mol Biomol Spectrosc* 219:313–318
- Koynova R, Tenchov B (2001) Interactions of surfactants and fatty acids with lipids. *Curr Opin Colloid Interface Sci* 6:277–286
- Kralova I, Sjöblom J (2009) Surfactants used in food industry: a review. *J Dispers Sci Technol* 30(9):1363–1383
- Kwak JCT (1998) *Polymer–surfactant systems*. Marcel Dekker, USA
- Langner M, Hui S (2000) Effect of free fatty acids on the permeability of 1,2-dimyristoyl-sn-glycero-3-phosphocholine bilayer at the main phase transition. *Biochim Biophys Acta* 1463(2):439–447
- Le Maire M, Champeil P, Moller JV (2000) Interaction of membrane proteins and lipids with solubilizing detergents. *Biochim Biophys Acta* 1508:86–111
- Lee HJ, McAuley A, Schilke KF et al (2011) Molecular origins of surfactant-mediated stabilizations of protein drugs. *Adv Drug Deliv Rev* 63:1160–1171
- Li Y, Lee JS (2019) Staring at protein–surfactant interactions: fundamental approaches and comparative evaluation of their combinations: a review. *Anal Chim Acta* 1063:18–39

- Lichtenberg D, Robson RJ, Dennis EAW (1983) Solubilization of phospholipid by detergents structural and kinetic aspects. *Biochim Biophys Acta* 737:285–304
- Lichtenberg D, Ahyauch H, Alonso A et al (2013) Detergent solubilization of lipid bilayers: a balance of driving forces. *Trends Biochem Sci* 38(2):85–93
- Mackie A, Wilde P (2005) The role of interactions in defining the structure of mixed protein–surfactant interfaces. *Adv Colloid Interface Sci* 117:3–13
- Malaspina T, Colherinhas G, Outi FO et al (2017) Assessing the interaction between surfactant-like peptides and lipid membranes. *RSC Adv* 7:35973–35981
- Maldonado F, Almela M, Otero A et al (1991) The binding of anionic and nonionic surfactants to collagen through the hydrophobic effect. *J Protein Chem* 10(2):189–192
- Malik NA (2015) Surfactant–amino acid and surfactant–surfactant interaction in aqueous medium: a review. *Appl Biochem Biotechnol* 176:2077–2106
- Mandal BB, Kund SC (2008) A novel method for dissolution and stabilization of non-mulberry silk gland protein fibroin using anionic surfactant sodium dodecyl sulfate. *Biotechnol Bioeng* 99:1482–1489
- Mankowich AM (1953) Physicochemical properties of surfactants. *Ind Eng Chem* 45(12):2759–2766
- Marquez R, Forgiarini AM, Langevin D et al (2018) Instability of emulsions made with surfactant–oil–water systems at optimum formulation with ultralow interfacial tension. *Langmuir* 34:9252–9263
- Marquez R, Anton R, Vejar F et al (2019) New interfacial rheology characteristics measured using a spinning drop Rheometer at the optimum formulation. Part 2. Surfactant–oil–water systems with a high volume of middle-phase microemulsion. *J Surfactants Deterg* 22:177–188
- Mehan S, Aswal VK, Kohlbrecher J (2015) Tuning of protein–surfactant interaction to modify the resultant structure. *Phys Rev E* 92:032713
- Morais IMC, Cordeiro AL, Teixeira GS et al (2017) Biological and physicochemical properties of biosurfactants produced by *Lactobacillus jensenii* P6A and *Lactobacillus gasseri* P65. *Microb Cell Fact* 16(155):1–15
- Nahringbauer I (1997) Polymer–surfactant interaction as revealed by the time dependence of surface tension. The EHEC/SDS/water system. *Langmuir* 13:2242–2249
- Naughton PJ, Marchant R, Naughton V et al (2019) Microbial biosurfactants: current trends and applications in agricultural and biomedical industries. *J Appl Microbiol* 127:12–28
- Ohadi M, Shahravan A, Dehghannoudeh N, Eslaminejad T, Banat IM, Dehghannoudeh G (2020) Potential use of microbial surfactant in microemulsion drug delivery system: a systematic review. *Drug Des Dev Ther* 14:541–550
- Ortiz A, Teruel JA, Espuny MJ et al (2008) Interactions of a *Rhodococcus* sp. biosurfactant trehalose lipid with phosphatidylethanolamine membranes. *Biochim Biophys Acta* 1778:2806–2813
- Ortiz A, Teruel JA, Espuny MJ et al (2009) Interactions of a bacterial biosurfactant trehalose lipid with phosphatidylserine membranes. *Chem Phys Lipids* 158(1):46–53
- Ortiz-Tafoya MC, Rolland-Sabaté A, Garnier C et al (2018) Thermal, conformational and rheological properties of  $\kappa$ -carrageenan-sodium stearyl lactylate gels and solutions. *Carbohydr Polym* 193:289–297
- Otzen D (2011) Protein–surfactant interactions: a tale of many states. *Biochim Biophys Acta* 1814:562–591
- Otzen DE (2017) Biosurfactants and surfactants interacting with membranes and proteins: same but different? *Biochim Biophys Acta* 1859:639–649
- Otzen DE, Sehgal P, Westh P (2009)  $\alpha$ -Lactalbumin is unfolded by all classes of surfactants but by different mechanisms. *J Colloid Interface Sci* 329:273–283
- Pacifico C, Giers S (1995) Surfactants used in textile applications. *J Am Oil Chem Soc* 32:231–235
- Pan F, Lu Z, Tucker I et al (2016) Surface active complexes formed between keratin polypeptides and ionic surfactants. *J Colloid Interface Sci* 484:125–134
- Park JH, Kim MH, Jeong L et al (2014) Effect of surfactants on sol–gel transition of silk fibroin. *J Sol-Gel Sci Technol* 71:364–371
- Parray M, Mir MUH, Dohare N et al (2018) Effect of cationic gemini surfactant and its monomeric counterpart on the conformational stability and esterase activity of human serum albumin. *J Mol Liq* 260:65–77
- Patino JMR, Niño MRR, Sánchez CC (2007) Physico-chemical properties of surfactant and protein films. *Curr Opin Colloid Interface Sci* 12(4):187–195
- Peypoux F, Bonmatin JM, Labbe H et al (1994) [Ala4] surfactin, a novel isoform from *B. subtilis* studied by mass and NMR spectroscopies. *Eur J Biochem* 224:89–96
- Pizzirusso A, De Nicola A, Sevink GJA et al (2017) Biomembrane solubilization mechanism by Triton X-100: a computational study of the three stage model. *Phys Chem Chem Phys* 19:29780–29794
- Poša M, Pilipović A, Torović L et al (2019) Co-solubilisation of a binary mixture of isoflavones in a water micellar solution of sodium cholate or cetyltrimethylammonium bromide: influence of micelle structure. *J Mol Liq* 273:134–146
- Proffitt TJ, Patterson T (1988) Oleochemical surfactants and lubricants in the textile industry. *J Am Oil Chem Soc* 65:1682
- Rodrigues L, Banat IM, Teixeira J et al (2006) Biosurfactants: potential applications in medicine. *J Antimicrob Chemother* 57:609–618
- Rudolph TW, Jones LS (2002) Surfactant–protein interactions. *Pharm Biotechnol* 13:159–175
- Ruiz-Peña M, Oropesa-Nuñez TP, Louro SRW et al (2010) Physico-chemical studies of molecular interactions between non-ionic surfactants and bovine serum albumin. *Colloids Surf B Biointerfaces* 75:282–289
- Santos D, Rufino R, Luna J et al (2016) Biosurfactants: multifunctional biomolecules of the 21st century. *Int J Mol Sci* 17:401
- Sinha S, Tikariha D, Lakra J et al (2016) Interactions of bovine serum albumin with cationic monomeric and dimeric surfactants: a comparative study. *J Mol Liq* 218:421–428
- Sonu SH, Kumari S, Aggrawal R et al (2017) Study on interactions of cationic gemini surfactants with folded and unfolded bovine serum albumin: effect of spacer group of surfactants. *J Mol Liq* 245:369–379
- Stryer S (1981) *Kimball's biology pages, "cell membranes."* Biochemistry 2nd ed. Freeman, USA

- Tucker IA, Petkov JT, Penfold J et al (2014) Spontaneous surface self-assembly in protein–surfactant mixtures: interactions between hydrophobin and ethoxylated polysorbate surfactants. *J Phys Chem* 118:4867–4875
- Thimon L, Peypoux F, Wallach J et al (1995) Effect of lipopeptide antibiotic, iturin A, on morphology and membrane ultrastructure of yeast cells. *FEMS Microbiol Lett* 128:101–106
- Van Os NM, Haak JR, Rupert LAM (1993) Physico-chemical properties of selected anionic, cationic and nonionic surfactants. Elsevier, USA
- Zaragoza A, Teruel JA, Aranda FJ et al (2012) Interaction of a *Rhodococcus* sp. trehalose lipid biosurfactant with model proteins: thermodynamic and structural changes. *Langmuir* 28(2):1381–1390
- Zhang J, Li Y (2018) Study on the interaction between surfactin and alkaline protease in aqueous solution. *Int J Biol Macromol* 118:244–251

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.